

ORIGINAL ARTICLE

The safety and effectiveness of minor injuries telemedicine

J R Benger, S M Noble, J Coast, J M Kendall

Emerg Med J 2004;21:438–445. doi: 10.1136/emj.2002.004028

See end of article for authors' affiliations

Correspondence to:
Dr J Benger, Academic
Department of Emergency
Care, Emergency
Department, Bristol Royal
Infirmary, Bristol BS2
8HW, UK;
JB@sectae.org.uk

Accepted for publication
29 January 2003

Objectives: To determine the safety of minor injuries telemedicine compared with on-site specialist care, current practice, and a robust gold standard, and to assess the clinical effectiveness of this new technique.

Methods: Patients presenting to a peripheral hospital within 10 days of injury were separately assessed by each of: an emergency medicine specialist based at a district general hospital using telemedicine, a second on-site emergency medicine specialist, and an on-site general practitioner (representing current practice). The primary outcome measure was discrepancies between these three medical assessments and a gold standard. All patients were subsequently randomised to follow one of the independent treatment plans generated by the above assessments. Secondary outcomes were recovery and further use of healthcare services measured seven days after recruitment, and consultation duration.

Results: 600 patients were recruited over a 12 month period. Overall, 73 discrepancies were identified, with 12 important over-treatments and 11 important under-treatments. No consultation modality was clearly superior to any other, and there were no statistically significant differences in the secondary outcomes of clinical effectiveness measured at seven days. The mean duration of a telemedicine consultation (6.0 min) was almost twice as long as an on-site specialist (3.1 min) or on-site general practitioner consultation (3.4 min) ($p < 0.0001$ in both cases).

Conclusions: Minor injuries telemedicine is safe and clinically effective, providing care that is equivalent to specialist on-site assessment and the current practice of treatment by a general practitioner. There is no evidence that telemedicine provides superior care, and there are a number of process issues that may impede successful implementation of this new technique.

Literally translated, telemedicine means “medicine at a distance”. This definition includes telephone use,¹ although the term telemedicine is more commonly used to describe a consultation involving the transmission of visual images. In truly remote locations, such as Antarctica,² the benefits of telemedicine are obvious. Extrapolation to less remote settings offers the potential to provide urgent medical advice without the need for prolonged patient or staff travel.

Recent years have seen considerable pressure to rationalise emergency services in the United Kingdom. The Audit Commission's report of 1996 highlighted a shortage of medical staff, and proposed the closure of up to 30 emergency departments nationally.³ On the other hand, rural communities seek to keep care as local as possible.⁴ One approach has been to develop the emergency nurse practitioner.⁵ Such people reduce pressure on medical staff and are well suited to the management of minor injuries.⁶ An increasing number of units are employing emergency nurse practitioners, who have been shown to be as effective as junior doctors in ordering and interpreting radiographs.⁷

Telemedicine can support emergency nurse practitioners working in remote locations, continuing to provide a local service with reference to central expertise.⁸ This approach has been endorsed by the British Association for Accident and Emergency Medicine.⁹ While a handful of studies have suggested that telemedicine may be clinically effective,^{10–13} no large or scientifically rigorous trials have been undertaken in the United Kingdom.¹⁴

This paper reports the results of a study conducted to determine the safety of minor injuries telemedicine in comparison with the current practice of general practitioner care, on-site specialist management, and a gold standard. The study also aimed to assess the clinical effectiveness of this new technique through the medium of a prospective, randomised, and blinded clinical trial. An assessment of the

relative efficiency of the different options will be reported separately (unpublished data).

METHODS

Setting

The peripheral unit is a typical community hospital located in a Cotswold town. Four experienced nurses, supported by healthcare assistants, staff the emergency department, which sees about 10 000 new patients per annum. Sixteen local general practitioners (GPs) from four practices provide medical input. On-site radiography is available from 9 am to 4 pm on Monday and Tuesday, and 9 am to 1 pm on Wednesday to Friday. Under normal circumstances about 35% of patients presenting with minor injuries are treated exclusively by nursing staff, 62% are referred to a GP (often to arrange and interpret radiography), and about 3% are referred directly by nursing staff to the main hospital. The main hospital has an emergency department typical of a district general hospital, seeing about 35 000 new patients per annum. None of the staff in either hospital had any prior experience of telemedicine.

A PictureTel Swiftsite videoconferencing system was installed in the emergency department of each hospital. At the peripheral site there were two additional devices: a PictureTel 550 document camera for radiological transmission, and a Panasonic VX54B camera to provide close up views. The equipment was installed and optimised four weeks before the start of the study. Staff training ensured competency in equipment operation, with telemedicine procedures based on a previously published protocol.¹⁵

Participants

Participants were recruited from patients attending the peripheral site at times when the onsite specialist (JRB, a year three specialist registrar in emergency medicine), was

Box 1 Inclusion and exclusion criteria**Inclusion criteria**

- The patient required, in the view of the nurse assessing them, an opinion from a doctor.
- The onsite specialist was present at the peripheral site.
- The patient had sustained an injury within the previous 10 days.

Exclusion criteria

- Patients unable or unwilling to be followed up. This was required as a safety measure, and to establish a gold standard in each case.
- Patients unable to provide informed consent.
- Patients with major injuries who clearly required immediate transfer.
- Patients who had already participated in the study.

available. The onsite specialist played no routine part in the peripheral unit, and was present only for the purposes of the trial. The onsite specialist attended the peripheral unit from 8 am to 8 pm on a total of 156 days spread throughout a 12 month period. Recruitment was not conducted overnight as patient attendances were very infrequent, and there was no senior doctor immediately available for telemedicine at the main hospital. An experienced nurse, who was also able to request radiographs of the peripheral skeleton, assessed each patient on arrival according to normal practice. Patients fulfilling the inclusion and exclusion criteria (box 1) were referred to the onsite specialist for recruitment, with the nurse prospectively recording what would have happened to the patient had they not entered the trial.

Written consent was obtained from all patients, or from a parent or legal guardian for children. Approval for the study was gained from the East Gloucestershire Ethics Committee.

Study design

To evaluate safety the study directly compared three independent medical assessments with a gold standard. In addition, a prospective, randomised, and blinded trial was conducted to measure clinical effectiveness. Process measures relating to consultation duration, radiograph requests, and follow up were also collected prospectively. Figure 1 is a summary trial flow diagram.

Patients were independently assessed by each of: the onsite emergency medicine specialist, a second emergency medicine specialist (either a consultant, staff grade, or specialist registrar) from the main hospital using telemedicine, and an onsite GP (if available). GPs visited the peripheral unit on a rota basis from their individual surgeries, and were therefore not always immediately available to assess recruited patients. Participants were generally unwilling to wait more than one or two hours to see a GP if two emergency medicine specialists had already seen them, and GP assessment was therefore omitted if a GP was not readily available in the peripheral unit.

Assessment by telemedicine entailed an initial telephone call to the main emergency department to indicate that a teleconsultation was required. When a senior doctor became available they started the telemedicine link and undertook a remote consultation with the patient, facilitated by the nurse.

Radiographs requested by the nurse before medical consultation were made available to all doctors during their assessment. If any doctor felt that further radiographs were required they requested these as part of their treatment plan,

but these radiographs were not made available to the other doctors unless they had also independently requested them.

Each doctor made an independent written record of their diagnosis, radiological interpretation, and itemised treatment plan with follow up arrangements. The doctor did not discuss these with the patient. When all treatment plans had been recorded patients were randomised to follow one of these plans using one-to-one-to-one block randomisation with a variable block size. A second nurse opened a sealed, opaque envelope corresponding to the patient's study number. This indicated which plan should be followed. The treatment instructions from this plan were transcribed onto a loose sheet, which was returned to the first nurse to enact. This ensured that neither the treating nurse nor patient knew which plan was being followed. Where the GP treatment plan was indicated a second randomised option was also stated for use if the GP assessment had been omitted.

All treatments were carried out by nursing staff, with the patient referred to a GP or transferred to the main hospital if further medical intervention was indicated on the treatment plan.

Participants were reviewed seven days after their initial assessment, as substantial improvement is common by this time. All participating doctors were equally aware of the trial protocol and arrangements for review. Patients remained blinded to which treatment plan they had followed. A standardised questionnaire was completed by each participant (or consenting adult in the case of a younger child). A gold standard diagnosis was then established by direct clinical assessment of each patient by the principal investigator, the report of a consultant radiologist blinded to the study and referral to a consultant specialist or further follow up until complete resolution.

Outcome measures

Outcomes were assessed in terms of safety, process, and clinical effectiveness. Safety was assessed by comparing each arm of the trial to the gold standard. All discrepancies in diagnosis, radiological interpretation, and treatment plan were submitted to an expert panel comprising 10 consultants in emergency medicine who were not otherwise involved in the study. Panel members did not know which trial arm had given rise to each discrepancy, and were asked to score these according to the scale shown in box 2, which was developed specifically for this research. The scores allocated by each of the 10 panel members were used to calculate both a mean and median discrepancy score. The mean score was selected for further analysis because it served as a better discriminator between the 44 discrepancies that had a median score of +1 or -1. Process outcomes were the duration of each consultation and the frequency with which radiographs and follow up appointments were requested. Measures of clinical effectiveness were derived from the review questionnaires, and are listed in box 3.

Sample size

The sample size was calculated for the primary outcome measure of safety, rather than the secondary outcome measures of clinical effectiveness that were addressed in the randomised trial. We selected a sample size of 600 because the absence of a significant discrepancy in this number would allow us to state, within 95% confidence limits, that the risk of significant discrepancy in actual clinical practice would lie somewhere between zero and 1 in 200.¹⁶ This is comparable to existing and accepted risks in emergency medicine.¹⁷ With regards to the secondary outcomes of clinical effectiveness, this sample size is sufficient to detect a 12% difference between telemedicine and the onsite specialist ($\alpha = 0.05$, two sided, and power 0.8).




Figure 1 Summarising the trial protocol.

Data handling and statistical methods

Data management and analysis were performed using a computerised Stata database (Stata Corporation, Texas, USA). In the randomised trial data were analysed on an intention to treat basis incorporating the two stage randomisation procedure. For process measures, statistical testing took into account the non-independent nature of the data. The numbers available for each pairwise comparison are noted, because not all patients received all three alternatives. For categorical variables the Cochran Q test and McNemar’s χ^2 test were used.¹⁸ Paired *t* tests were used for normally distributed continuous variables. To take account of the multiple significance testing, a Bonferroni correction was applied.¹⁹ This meant that to maintain a 5% significance level *p* values from the three pairwise tests between the three groups were only considered significant if they were below 0.017.

For effectiveness data χ^2 testing and the Kruskal-Wallis equality of populations rank test were used. A two sided 5% significance level was used throughout, and 95% confidence intervals (CI) are presented where appropriate.

RESULTS

Recruitment and patient progression

Recruitment of patients took place between December 1999 and November 2000. Figure 2 summarises the flow of patients through the clinical effectiveness trial. A total of 713 patients met the inclusion criteria, of whom 103 were excluded and 10 declined to participate. The “other” exclusion was an 8 year old girl with a penetrating injury to the perineum, who was not considered an appropriate candidate for telemedicine. In 21 recruited patients (11 of whom were subsequently randomised to follow the telemedicine treatment plan) no telemedicine consultation took place because the main hospital was too busy with seriously ill patients to respond within two hours: this meant that for the primary outcome of safety, telemedicine could be compared with onsite specialist assessment in 579 patients, and compared with onsite GP assessment in 165 patients.

In the effectiveness trial, of the 600 patients recruited, 200 were initially randomised to receive the telemedicine treatment plan, 201 the onsite specialist plan and 199 the onsite GP plan. However, 135 patients randomised to the GP plan

Box 2 Expert panel discrepancy assessment scale**+3: Over (excessive) treatment leading to major inconvenience**

(unnecessary treatment restrictions preventing the performance of a normal occupation for greater than seven days, or a less severe restriction for longer)

+2: Over (excessive) treatment leading to moderate inconvenience

(unnecessary treatment restrictions preventing the performance of a normal occupation for three to seven days, or a less severe restriction for longer)

+1: Over (excessive) treatment leading to mild inconvenience

(unnecessary treatment restrictions preventing the performance of a normal occupation for one or two days, or an unnecessary visit to the main hospital)

0: No significant difference between the gold standard and trial treatment**-1: Under (inadequate) treatment with mild consequences**

(likely to cause avoidable symptoms for less than one week)

-2: Under (inadequate) treatment with moderate consequences

(likely to cause avoidable symptoms for between one and six weeks)

-3: Under (inadequate) treatment with severe consequences

(likely to cause avoidable symptoms for greater than six weeks)

had not seen a GP, because no GP was immediately available to assess the patient. Therefore secondary randomisation assigned 74 of these patients to telemedicine and 61 to the onsite specialist plan. This meant that 274 patients were allocated to telemedicine, 262 to the onsite specialist plan, and 64 to the GP treatment plan.

The mean age of recruited patients was 34 years with a range of 9 months to 92 years. A total of 475 patients (79.2%) were registered with a local GP. Table 1 shows the characteristics of all recruited patients by the treatment plan to which they were randomised.

Safety

There were 73 cases where a study diagnosis or radiological interpretation, or both, differed from the gold standard. Expert panel assessment of these gave:

- 12 discrepancies with a mean score greater than +1 ("over-treatments")
- 35 discrepancies with a mean score between +1 and 0
- 15 discrepancies with a mean score between 0 and -1
- 11 discrepancies with a mean score less than -1 ("under-treatments")

The 50 discrepancies with a mean score between +1 and -1 were excluded from further analysis as these were minor, and attributed to variations between clinicians rather than errors likely to have a significant effect on patient outcome.

The 12 "over-treated" and 11 "under-treated" cases are detailed further in tables 2 and 3. The significant under-treatment rate ranged from 0% (95% CI 0% to 1.7%) for general practitioners to 0.5% (95% CI 0.1% to 1.5%) for the onsite specialist, and 1.4% (95% CI 0.6% to 2.7%) for telemedicine. Telemedicine appears to have the highest rate of significant under-treatment, but this is not statistically significant on χ^2 testing. As none of the 579 telemedicine

Box 3 Clinical effectiveness outcomes derived from the review questionnaires

- Whether the problem had resolved.
- Whether the patient had been able to return to normal activity.
- Whether the patient had taken time off work (where applicable).
- Whether there had been further health services contact regarding the same condition.
- Whether the patient had sought further unscheduled care.
- Whether there had been any change in treatment since the patient entered the trial.

consultations received a score worse than -2 it is possible to state that for minor injuries telemedicine the risk of error leading to severe consequences has a 95% confidence interval between 0% and 0.52%.

Process measures

The mean delay between injury and presentation at the peripheral hospital was 1.3 days (95% CI 1.2 to 1.5 days). The mean wait to see a nurse at the peripheral hospital was 4.2 minutes (95% CI 3.4 to 5.0 minutes).

Table 4 shows the action that would have been taken by the assessing nurse, if the patient had not been recruited into the trial.

For all telemedicine consultations, the mean delay between a request for telemedicine being made and actual start of the teleconsultation was 7.1 minutes (95% CI 6.3 to 7.9 minutes). Technical failure occurred in two cases (0.3%).

The mean duration of all consultation types was as follows: telemedicine 6.0 minutes (95% CI 5.7 to 6.2 minutes), onsite specialist 3.1 minutes (95% CI 2.9 to 3.3 minutes), and onsite general practitioner 3.4 minutes (95% CI 3.2 to 3.6 minutes). Paired *t* testing showed no difference between the onsite specialist and GP consultations ($p=0.64$, $n=177$), but a significant difference between the time taken to perform a telemedicine consultation and either an onsite specialist ($n=578$) or GP consultation ($n=171$) ($p<0.0001$ in both cases). The average time taken to conduct a telemedicine consultation reduced as the study progressed, with a mean of 7.3 minutes (95% CI 6.2 to 8.4 minutes) for the first 100 patients and 5.0 minutes (95% CI 4.7 to 5.4 minutes) for the last 100. However, telemedicine consultations remained significantly ($p<0.0001$) longer than either the onsite specialist or the GP for the last 100 patients.

For each consultation type the number of patients who had radiographs and asked to attend for a further follow up appointment is shown in table 5. Overall there were no significant differences in the proportion who had radiographs by the three alternatives ($p=0.11$, $n=171$, Cochran Q test). There were, however, significant overall differences between the proportion followed up by the three alternatives ($p<0.0001$, $n=170$, Cochran Q test). GPs asked a significantly greater proportion of patients to attend for follow up than either telemedicine ($p<0.0001$, $n=170$, McNemar's χ^2 test), or the onsite specialist ($p<0.0001$, $n=177$ McNemar's χ^2 test). A significantly greater proportion of patients were asked to attend for follow up after a telemedicine consultation than after an onsite specialist consultation ($p<0.0001$, $n=578$, McNemar's χ^2 test). The proportion of patients asked to attend for follow up after a telemedicine consultation




Figure 2 Flow of patients through the trial of clinical effectiveness.

did not reduce during the study, being identical at 40% for both the first 100 and last 100 patients.

Clinical effectiveness

Eight patients (1.3%) were lost from the study. The mean time between recruitment and review was 7.0 days (SD

1.4 days, range 3 to 14 days). There was no significant difference in the time to review between the three randomisation groups ($p = 0.82$, Kruskal-Wallis test).

Table 6 summarises the main measures of clinical effectiveness. No significant differences were detected for any of these outcomes

Table 1 Baseline characteristics of participants by randomisation of treatment plan

Variable	Telemedicine (n = 274)	Onsite specialist (n = 262)	General practitioner (n = 64)
Sex			
Female	111 (40.5)	99 (37.8)	20 (31.2)
Male	163 (59.5)	163 (62.2)	44 (68.8)
Age			
<16	60 (21.9)	56 (21.4)	18 (28.1)
16–24	44 (16.1)	53 (20.2)	15 (23.4)
25–34	52 (19.0)	41 (15.7)	9 (14.1)
35–44	37 (13.5)	39 (14.9)	5 (7.8)
45–54	31 (11.3)	29 (11.1)	6 (9.4)
55–64	21 (7.7)	23 (8.8)	6 (9.4)
65+	29 (10.6)	21 (8.0)	5 (7.8)
In employment/full time education	210 (77.8)*	205 (79.2)†	51 (81)‡

*n = 270; †n = 259; ‡n = 63. Percentages are shown in parentheses.

Table 2 The 12 discrepancies scoring more than +1. The second column details the incorrect diagnosis made

Gold standard	Error made	Mean error score	Error source
Wrist sprain	Fracture diagnosed	1.8	Onsite specialist
Ankle sprain	Fracture diagnosed	1.7	Onsite specialist
Wrist sprain	Fracture diagnosed	1.6	General practitioner
Elbow bruising	Fracture diagnosed	1.4	Onsite specialist
Shoulder bruising	Fracture diagnosed	1.3	Telemedicine
Thumb sprain	Ulna collateral ligament rupture diagnosed	1.3	Onsite specialist
Ankle sprain	Fracture diagnosed	1.3	Telemedicine
Ankle sprain	Fracture diagnosed	1.2	Telemedicine
Finger lacerations and bruising	Compound fracture diagnosed	1.2	Telemedicine
Bruising to shin	Cellulitis diagnosed	1.1	Onsite specialist
Wrist sprain	Fracture diagnosed	1.1	Onsite specialist
Wrist sprain	Fracture diagnosed	1.1	Telemedicine

DISCUSSION

These results show that the safety of minor injuries telemedicine is similar to conventional practice. It is inevitable that errors will occur during any large series of consultations, and this research quantified and compared these errors for the three alternatives studied, finding no significant differences between them.

Only one randomised trial of telemedicine in a similar context has previously been published.¹³ This was a much smaller study, conducted in North America with just 104 participants, but showed similar results.

Over-treatments, radiographs, and follow up

We studied the over-treatment rate because this might be increased for telemedicine, particularly if medical staff were unsure of the diagnosis, but found no evidence that this was the case.

While the radiography rate was nearly identical for telemedicine and onsite specialist consultation (59.2% compared with 60.5%), a follow up appointment was arranged for a significantly greater proportion of telemedicine patients (35.8% compared with 27.5%: $p < 0.0001$ using McNemar's χ^2 test), perhaps reflecting a lack of confidence in telemedicine. This did not reduce during the course of the study. The highest rate of planned follow up was seen in GP consultations (65%). This may reflect the way in which GPs normally practice, as we observed that follow up appointments arranged by a GP were frequently an option that the patient could pursue only if problems persisted. This is supported by the finding that, at the

seven day study review, there were no significant differences between the randomisation groups in planned or unplanned use of healthcare services.

Under-treatments

Under-treatments are of greater importance as they may lead to long term effects and possible litigation. Several studies have attempted to quantify the incidence of error in emergency medicine, often concentrating on radiological interpretation. In the United Kingdom, a recent study quoted an error rate of 1.5%, falling to the more acceptable figure of 0.7% when a picture archiving and communications system (PACS) was introduced.²⁰ In emergency care it has been necessary to establish a risk below which a patient can be reasonably discharged, and this threshold is commonly set at 1%.²¹ In this study only one consultation, by the onsite specialist, received a mean under-treatment score worse than -2, and for telemedicine the risk of error leading to severe consequences was less than 1% (95% CI 0% to 0.52%).¹⁶

Process measures

Patients attending the peripheral hospital were usually seen by a nurse very promptly, and had to wait a mean of only 7.1 minutes for a telemedicine consultation, excepting those cases where the main hospital was too busy to respond. This suggests that patients attending the peripheral site were likely to be assessed by a senior doctor much more rapidly than if they had attended the main hospital in person. Indeed, medical staff commented that there was a possibility

Table 3 The 11 discrepancies with a mean score less than -1. The second column details the diagnosis missed

Gold standard	Error made	Mean error score	Error source
Conjunctival laceration	Laceration missed	-2.5	Onsite specialist
Undisplaced fracture of the distal fibula	Fracture missed	-1.8	Telemedicine
Compound finger fracture	Fracture missed	-1.6	Telemedicine
Greenstick fracture of the distal radius	Fracture missed	-1.6	Telemedicine
Infected hand laceration	Infection missed	-1.6	Telemedicine
Infected shoulder abrasion	Infection missed	-1.4	Telemedicine
Fractured clavicle	Fracture missed	-1.2	Telemedicine
Calf cellulitis	Infection missed	-1.2	Telemedicine
Fractured clavicle	Fracture misdiagnosed	-1.2	Onsite specialist
Avulsion fracture from the tip of the olecranon	Fracture missed	-1.2	Telemedicine
Knee laceration with an underlying patella fracture	Fracture missed	-1.1	Onsite specialist

Table 4 Action that would have been taken by the assessing nurse if the patient had not been recruited into the trial

Action	Number (n = 600)	Percentage
Telephone general practitioner and ask them to come to the peripheral emergency department to see patient	60	10.0
Arrange appointment for patient with a general practitioner	50	8.3
Bring patient back to see general practitioner later today or on a following day	282	47.0
Ask patient to wait until a general practitioner comes to the peripheral emergency department	147	24.5
Arrange for patient to be seen at the peripheral emergency department specialist clinic	11	1.8
Refer directly to the main emergency department	14	2.3
Refer to an inpatient team at the main hospital	3	0.5
Ring general practitioner for advice	31	5.2
Ring the main emergency department for advice	2	0.3
Total	600	100.0

that they would leave a patient with more urgent problems to respond to a request for minor injuries telemedicine.

If telemedicine acted to preferentially bypass the queue in the main emergency department, patients might choose to travel to the peripheral hospital for treatment. One approach to this problem would be to "triage" telemedicine cases as if they had come to the main site in person. This would cause practical difficulties at the peripheral site, and might also prove unacceptable to patients who are used to the prompt service that the peripheral hospital usually provides.

Only 17 (2.8%) of patients would have been referred directly to the main hospital if they had not been recruited into the trial, so the potential for telemedicine to reduce unnecessary patient transfers in this context is very limited. The nurse's assessment of what would have happened had the patient not been recruited could have been biased by the knowledge that the patient was about to enter the trial, but this transfer rate of 2.8% is very similar to the 3% which existed before the start of the study. The low transfer rate is attributable to the effective service already provided, and the fact that most of the injuries were minor. If the telemedicine plan had been followed for these 17 patients then eight would still have been transferred, with the remaining nine treated locally.

Telemedicine weaknesses

In 16 cases (2.8%) the doctor providing a telemedical opinion felt that telemedicine was inadequate in some way. This related either to problems in visualising and interpreting a radiograph (particularly chest radiographs) or to difficulties in assessing a wound (for example, burn depth). Nevertheless, no significant discrepancy resulted in any of these cases, suggesting that the problem was recognised and appropriately resolved.

Problems with telemedicine are often attributed to technical deficiencies that can be overcome by improving the performance of equipment.²² In this study, however, we were conforming (with the exception of using a television rather than a computer monitor) to a technical standard that achieves satisfactory results in a laboratory setting.²³ This suggests that

further improvements in technology may not provide complete solutions, and that some difficulties relate to the medium itself.

Study weaknesses

We assumed that treatment discrepancies were attributable to telemedicine rather than variations in clinician performance. As most participants had comparatively straightforward injuries, and the doctors involved were of comparable experience, this assumption appears reasonable. Furthermore, the use of a large expert panel and removal of discrepancies scoring between +1 and -1 should have prevented individual treatment variation from influencing the results.

It was not possible to achieve consecutive recruitment because of the requirement for the onsite specialist to be available. Nevertheless, minor injuries rarely presented overnight, when recruitment would have been inefficient, and there is no reason to suspect that our sample is not representative of the wider population of minor injuries patients.

The methodology of this study was unusual in that rather than being randomised to receive only one type of consultation, we elected to have each recruited patient undergo more than one consultation type, with strict blinding between the consultations. This allowed us to collect a large sample in which telemedicine and conventional consultation could be directly compared with high sensitivity. A disadvantage was that the sample size was not specifically calculated for the randomised trial of clinical effectiveness. Therefore the clinical outcomes, although supportive of our safety data, should be viewed with caution, particularly as only a comparatively small number of patients followed a GP treatment plan.

Each patient underwent two or three consecutive consultations, and the onsite assessment always preceded telemedicine. This could lead to an order effect, with refinement and shortening of the history given by the patient. If such an effect existed, however, it would tend to reduce the length of telemedicine consultations, in contrast with the results obtained. The GP consultations might also have been shortened by this effect, although some were performed before and some after telemedicine. The doctors involved were also careful to avoid

Table 5 Comparison of the radiography and follow up rates for each type of consultation

Consultation type	Number of patients radiographed	Number of patients asked to attend for follow up
Telemedicine (n = 579)	343 (59.2) (55.1% to 63.3%)	207 (35.8) (31.8% to 39.8%)
Onsite specialist (n = 600)	363 (60.5) (56.5% to 64.4%)	165 (27.5) (24.0% to 31.3%)
General practitioner (n = 177)	94 (53.1) (45.5% to 60.6%)	115 (65.0) (57.5% to 72.0%)

Data shown as numbers, percentages, and 95% CI.

Table 6 Summary of the main measures of clinical effectiveness

Outcome measure at review	Telemedicine (n = 270)	Onsite specialist (n = 259)	General practitioner (n = 63)	p Value
Completely better	67 (24.8) (19.8% to 30.4%)	61 (23.6) (18.5% to 29.2%)	10 (15.9) (7.9% to 27.3%)	0.32
Returned to normal activity	127 (47.0) (41.0% to 53.2%)	106 (40.9) (34.9% to 47.2%)	30 (47.6) (34.9% to 60.6%)	0.32
Taken time off work	91 (33.7) (28.1% to 39.7%)	95 (36.7) (30.8% to 42.9%)	18 (28.6) (17.9% to 41.3%)	0.45
Seen in an emergency department	89 (33.0) (27.4% to 38.9%)	74 (28.6) (23.2% to 34.5%)	25 (39.7) (27.6% to 52.8%)	0.20
Seen in an outpatient clinic	37 (13.7) (9.8% to 18.4%)	41 (15.8) (11.6% to 20.9%)	3 (4.8) (1.0% to 13.3%)	0.07
Seen in a general practitioner's surgery or at home	6 (2.2) (0.8% to 4.8%)	10 (3.9) (1.9% to 7.0%)	2 (3.2) (0.4% to 11.0%)	0.55
Sought further unscheduled care	14 (5.2) (2.9% to 8.5%)	15 (5.8) (3.3% to 9.4%)	4 (6.3) (1.8% to 15.5%)	0.91
Required a change in treatment	26 (9.6) (6.4% to 13.8%)	17 (6.6) (3.9% to 10.3%)	4 (6.3) (1.8% to 15.5%)	0.37

Data shown as numbers, percentages, and 95% CI.

giving information to the patient during each consultation, minimising the effect of consecutive consultations.

Application to practice

We have shown that telemedicine is safe and effective in minor injuries practice. However, there is very little evidence regarding the effectiveness of telemedicine in other aspects of emergency medicine, such as acute illness or more severe injury.

If telemedicine is used to support nursing staff a high consultation volume can be expected, particularly if it replaces an existing medical service. This is time consuming for the hospital providing the consultations, and also runs the risk that the transfer rate will be increased because of system failures or diagnostic doubt. Direct referral to the main hospital was a comparatively rare event (2.8%), but where it occurred transfer was avoided in 53% of cases. Therefore, telemedicine used to support doctors or as a supplement to an existing service may approximately halve the patient transfer rate, and is in keeping with the one previous paper that studied telemedicine to support remote general practitioners, in which transfer was avoided in 58% of cases.¹¹

While a comparatively low cost system is capable of delivering satisfactory clinical results in terms of safety and patient outcome, there are a number of potential problems that may be overlooked by enthusiasts. Many of these problems will not be influenced by further technological advances, and must be tackled in other ways: some are inherent to telemedicine itself, but many are related to service delivery and organisation. It is important that decision makers are not overly distracted by technical concerns but focus on the managerial and economic issues that underpin telemedicine as a change in the way that care is delivered.

Conclusion

In the management of minor injuries, telemedicine is capable of providing a satisfactory standard of care. Significant errors are rare and similar in frequency to conventional practice and accepted risks. There is no evidence that telemedicine is superior to the routine practice of GP consultation.

Telemedicine requires about twice as long as an onsite consultation, in addition to the time entailed in responding to a request and breaking away from other activities. While telemedicine may be effective for minor injuries work, emergency medicine encompasses a variety of other conditions that may be less amenable to this technology, and where research is still required.

ACKNOWLEDGEMENTS

Our thanks go to the patients and staff of Tewkesbury and Cheltenham Hospitals and all the general practitioners who participated in this study, and without whom it would not have been possible. Also our grateful acknowledgements go to Professor Gordon Wilcock for his academic supervision, Professor Richard Wootton for his advice and support, Mr Christopher Foy and Professor Tim Peters for their statistical advice, and Mr Iain Grant,

Mr Paul Howarth, Dr Tom Llewellyn, Mr Christian Oakland, Mr Sapal Tachakra, and Mr Paul Wilson for their assistance in the development of this study.

Authors' affiliations

J R Benger, East Gloucestershire NHS Trust, Tewkesbury Hospital, Tewkesbury, Gloucestershire, UK

S Noble, J Coast, Department of Social Medicine, University of Bristol, Bristol, UK

J Kendall, Emergency Department, Frenchay Hospital, Bristol, UK

Funding: this research was funded by a grant from the Department of Health under the Information and Communications Technology initiative (ICT/312).

Conflicts of interest: none declared.

REFERENCES

- 1 McLaren P, Ball CJ. Telemedicine: lessons remain unheeded. *BMJ* 1995;**310**:1390-1.
- 2 Hyer RN. Telemedical experiences at an Antarctic station. *J Telemed Telecare* 1999;**5**(suppl 1):87-9.
- 3 Audit Commission. *By accident or design*. London: HMSO, 1996.
- 4 FGH Consortium. *Keeping care local feasibility study*. Cardiff: Welsh Office Health Department, 1998.
- 5 Read SM, Jones NMB, Williams BT. Nurse practitioners in accident and emergency departments: what do they do? *BMJ* 1992;**305**:1466-70.
- 6 Brebner JA, Ruddick-Bracken H, Norman JN, et al. The nurse practitioner: management of minor trauma. *Accident and Emergency Nursing* 1996;**4**:43-6.
- 7 Freij RM, Duffy T, Hackett D, et al. Radiographic interpretation by nurse practitioners in a minor injuries unit. *Emerg Med J* 1996;**13**:41-3.
- 8 Benger JR, Wootton R. Minor injuries telemedicine. *J Telemed Telecare* 1999;**5**(suppl 3):1-50.
- 9 British Association for Accident and Emergency Medicine. *The way ahead policy document*. London: British Association for Accident and Emergency Medicine, 1998.
- 10 Darkins A, Dearden CH, Rocke LG, et al. An evaluation of telemedical support for a minor treatment centre. *J Telemed Telecare* 1996;**2**:93-9.
- 11 Armstrong IJ, Haston WS. Medical decision support for remote general practitioners using telemedicine. *J Telemed Telecare* 1997;**3**:27-34.
- 12 Tachakra S, Wiley C, Dawood M, et al. Evaluation of telemedical support to a free-standing minor accident and treatment service. *J Telemed Telecare* 1988;**4**:140-5.
- 13 Brennan JA, Kealy JA, Gerardi LH, et al. A randomized controlled trial of telemedicine in an emergency department. *J Telemed Telecare* 1998;**4**(suppl 1):18-20.
- 14 Benger JR. A review of telemedicine in accident and emergency: the story so far. *Emerg Med J* 2000;**17**:157-64.
- 15 Tachakra S, Sivakumar A, Hayes J, et al. A protocol for telemedical consultation. *J Telemed Telecare* 1997;**3**:163-8.
- 16 Hanley JA, Lippman-Hand A. If nothing goes wrong is everything all right? *JAMA* 1983;**249**:1743-5.
- 17 Guly HR. Risk Management. *Emerg Med J* 1997;**14**:408-10.
- 18 Siegel S, Castella NJ. *Nonparametric statistics for the behavioral sciences*. New York: McGraw-Hill, 1988.
- 19 Altman DG. *Practical statistics for medical research*. London: Chapman and Hall, 1991.
- 20 Weatherburn G, Bryan S, Nicholas A, et al. The effect of a picture archiving and communications system (PACS) on diagnostic performance in the accident and emergency department. *Emerg Med J* 2000;**17**:180-4.
- 21 Wears RL, Li S, Hernandez JD, et al. How many myocardial infarctions should we rule out? *Ann Emerg Med* 1989;**18**:953-63.
- 22 Wootton R, Dornan J, Fisk NM, et al. The effect of transmission bandwidth on diagnostic accuracy in remote fetal ultrasound scanning. *J Telemed Telecare* 1997;**3**:209-14.
- 23 Benger JR, Lock A, Cook J, et al. The effect of resolution, compression, colour depth and display modality on the accuracy of accident and emergency telemedicine. *J Telemed Telecare* 2001;**7**(suppl 1):6-7.



The safety and effectiveness of minor injuries telemedicine

J R Benger, S M Noble, J Coast, et al.

Emerg Med J 2004 21: 438-445

Updated information and services can be found at:

<http://emj.bmj.com/content/21/4/438.full.html>

These include:

References

This article cites 18 articles, 8 of which can be accessed free at:

<http://emj.bmj.com/content/21/4/438.full.html#ref-list-1>

Article cited in:

<http://emj.bmj.com/content/21/4/438.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Telemedicine](#) (99 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>