




# Genome Sequence of *Lecanicillium fungicola* 150-1, the Causal Agent of Dry Bubble Disease

Alice M. Banks,<sup>a\*</sup> Farhana Aminuddin,<sup>a</sup> Katherine Williams,<sup>a</sup> Thomas Batstone,<sup>a</sup> Gary L. A. Barker,<sup>a</sup> Gary D. Foster,<sup>a</sup>  
 Andy M. Bailey<sup>a</sup>

<sup>a</sup>School of Biological Sciences, University of Bristol, Bristol, United Kingdom

**ABSTRACT** The fungus *Lecanicillium fungicola* causes dry bubble disease in the white button mushroom *Agaricus bisporus*. Control strategies are limited, as both the host and pathogen are fungi, and there is limited understanding of the interactions in this pathosystem. Here, we present the genome sequence of *Lecanicillium fungicola* strain 150-1.

*Lecanicillium fungicola* (Preuss) Zare & Gams [synonym: *Verticillium fungicola* (Preuss) Hassebrauk] (1), an ascomycete fungus of the order Hypocreales, is the causal agent of dry bubble disease of the white button mushroom *Agaricus bisporus*, as well as of other commercially cultivated basidiomycetes (2). Dry bubble disease presents symptoms that include necrotic lesions on mushroom caps, stipe blowout, and undifferentiated tissue masses (2). Some factors involved in this interaction have been proposed based on suppression subtractive hybridization (SSH) and expressed sequence tag (EST) data (3). This disease is of economic importance, causing significant yield/quality losses in the mushroom industry (4). Control methods rely on rigorous hygiene procedures and targeted fungicide treatments; however, increased resistance against these fungicides has been reported (5, 6). Recent taxonomic revisions place *L. fungicola* close to several arthropod- and nematode-pathogenic fungi rather than to plant-pathogenic *Verticillium* spp. (1, 7).

*L. fungicola* strain 150-1, a UK isolate of medium virulence obtained from Warwick-HRI, UK (2), was maintained on potato dextrose agar. Genomic DNA was extracted from freeze-dried mycelium using the gentle toluene lysis method of Bonsch et al. (8), and DNA was fragmented to ~520 bp and sequenced using paired-end 100-base reads with an Illumina HiSeq 2500 instrument, generating 92,464,338 reads. Data were processed using Real-Time Analysis (RTA) 1.17.21.3 with default settings and were demultiplexed with CASAVA 1.8.2. CLC Genomics Workbench 6 (Qiagen Bioinformatics) was used for quality trimming (reads with Phred scores of >20 and reads shorter than 50 nucleotides [nt] were discarded), and an assembly comprising 781 contigs was obtained, spanning 44,574,141 bp and with an  $N_{50}$  value of 154,124 bp. The GC content was 49.8%.

Contig 84 was identified as the almost-complete mitogenome, and reanalysis of mapped reads to allow circularization of the contig revealed an extra 11 bases required to complete the circular genome, which spans 24,277 bp. The mitogenome shows synteny to that of the nematophagous fungus *Lecanicillium saksenae* (9), including identical gene order and just one intron in *rn1*, again reflecting the close phylogenetic relationship between these fungi. Manual annotation was performed by comparison to closely related mitogenomes (9, 10) with tRNAs identified using tRNAscan-SE (11).

Some of the symptoms of dry bubble, such as cap spotting, might be due to secondary metabolites. Analysis using fungiSMASH 5.0.0 (12) revealed the following 38 biosynthetic gene clusters for secondary metabolites: 8 polyketide synthases (PKSs), 21

**Citation** Banks AM, Aminuddin F, Williams K, Batstone T, Barker GLA, Foster GD, Bailey AM. 2019. Genome sequence of *Lecanicillium fungicola* 150-1, the causal agent of dry bubble disease. *Microbiol Resour Announc* 8:e00340-19. <https://doi.org/10.1128/MRA.00340-19>.

**Editor** Antonis Rokas, Vanderbilt University

**Copyright** © 2019 Banks et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Andy M. Bailey, [Andy.Bailey@bristol.ac.uk](mailto:Andy.Bailey@bristol.ac.uk).

\* Present address: Alice M. Banks, School of Natural and Environmental Sciences, Newcastle University, Newcastle upon Tyne, United Kingdom.

A. M. Banks and F. Aminuddin contributed equally to this work and should be considered joint first authors.

**Received** 22 March 2019

**Accepted** 14 April 2019

**Published** 9 May 2019

nonribosomal peptide synthase (NRPS) or NRPS-like clusters, 3 PKS-NRPS hybrids, 5 terpene synthases, and 1 indole cluster. Analysis of these gene clusters is ongoing.

The interaction between *L. fungicola* and its primary host, *A. bisporus*, remains far from fully characterized. While the disruption of  $\beta$ -1,6 glucanase resulted in reduced virulence (13), knockout of the *pmk1*-like mitogen-activated protein (MAP) kinase did not affect virulence (14), indicating that the models established for fungal plant pathogens might not readily apply to fungus-fungus interactions.

The genome sequence of *L. fungicola* provides a useful basis to uncover the molecular mechanisms underlying pathogenicity in the *L. fungicola*-*A. bisporus* interaction.

**Data availability.** This project has been deposited in DDBJ/ENA/GenBank under the accession number [FWCC00000000](https://www.ncbi.nlm.nih.gov/nuccore/FWCC00000000) and BioProject number [PRJEB19844](https://www.ncbi.nlm.nih.gov/bioproject/PRJEB19844). The genome assembly version is the first version and includes accession numbers [FWCC01000001](https://www.ncbi.nlm.nih.gov/nuccore/FWCC01000001) to [FWCC01000781](https://www.ncbi.nlm.nih.gov/nuccore/FWCC01000781). Raw sequencing data have been deposited under accession number [ERR3181828](https://www.ncbi.nlm.nih.gov/nuccore/ERR3181828). The annotated mitogenome has been deposited under accession number [LR536627](https://www.ncbi.nlm.nih.gov/nuccore/LR536627).

## ACKNOWLEDGMENTS

We acknowledge the generosity of Steve Lincoln, Warwick-HRI, United Kingdom, in providing strain 150-1. The genome was sequenced by the University of Bristol Genomics Facility.

K. Williams and T. Batstone were supported by grant MR/N029909/1 from the UK Medical Research Council, awarded to A. M. Bailey.

## REFERENCES

- Zare R, Gams W. 2008. A revision of the *Verticillium fungicola* species complex and its affinity with the genus *Lecanicillium*. *Mycol Res* 112: 811–824. <https://doi.org/10.1016/j.mycres.2008.01.019>.
- Mills PR, Thomas J, Sergeant MR, Costa A, Collopy PD, Bailey AM, Foster GD, Challen MP. 2008. Interactions between *Agaricus bisporus* and the pathogen *Verticillium fungicola*, p 1–17. In Avery SV, Stratford M, van West P (ed), *Stress in yeasts and filamentous fungi*, 1st ed, vol 27. Academic Press, London, United Kingdom.
- Bailey AM, Collopy PD, Thomas DJ, Sergeant MR, Costa AMSB, Barker GLA, Mills PR, Challen MP, Foster GD. 2013. Transcriptomic analysis of the interactions between *Agaricus bisporus* and *Lecanicillium fungicola*. *Fungal Genet Biol* 55:67–76. <https://doi.org/10.1016/j.fgb.2013.04.010>.
- Berendsen RL, Baars JJP, Kalkhove SIC, Lugones LG, Wosten HAB, Bakker PAHM. 2010. *Lecanicillium fungicola*: causal agent of dry bubble disease in white-button mushroom. *Mol Plant Pathol* 11:585–595. <https://doi.org/10.1111/j.1364-3703.2010.00627.x>.
- Chrysai-Tokousbalides M, Kastanias MA, Philippoussis A, Diamantopoulou P. 2007. Selective fungitoxicity of famoxadone, tebuconazole and trifloxystrobin between *Verticillium fungicola* and *Agaricus bisporus*. *Crop Prot* 26:469–475. <https://doi.org/10.1016/j.cropro.2006.02.016>.
- Grogan HM, Gaze RH. 2000. Fungicide resistance among *Cladobotryum* spp.—causal agents of cobweb disease of the edible mushroom *Agaricus bisporus*. *Mycol Res* 104:357–364. <https://doi.org/10.1017/S0953756299001197>.
- Amey RC, Athey-Pollard A, Mills PR, Foster GD, Bailey A. 2007. Investigations into the taxonomy of the mushroom pathogen *Verticillium fungicola* and its relatives based on sequence analysis of nitrate reductase and ITS regions. *Microbiology* 156:757–768. <https://doi.org/10.1134/S0026261707060161>.
- Bonsch B, Belt V, Bartel C, Duensing N, Koziol M, Lazarus CM, Bailey AM, Simpson TJ, Cox RJ. 2016. Identification of genes encoding squalestatin S1 biosynthesis and *in vitro* production of new squalestatin analogues. *Chem Commun* 52:6777–6780. <https://doi.org/10.1039/c6cc02130a>.
- Xin B, Lin R, Shen B, Mao Z, Cheng X, Xie B. 2017. The complete mitochondrial genome of the nematophagous fungus *Lecanicillium sak-senae*. *Mitochondrial DNA Part A*. 28:52–53. <https://doi.org/10.3109/19401736.2015.1110794>.
- Kouvelis VN, Ghikas DV, Typas MA. 2004. The analysis of the complete mitochondrial genome of *Lecanicillium muscarium* (synonym *Verticillium lecanii*) suggests a minimum common gene organization in mtDNAs of Sordariomycetes: phylogenetic implications. *Fungal Genet Biol* 41: 930–940. <https://doi.org/10.1016/j.fgb.2004.07.003>.
- Lowe TM, Chan PP. 2016. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res* 44: W54–W57. <https://doi.org/10.1093/nar/gkv413>.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Brucoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43:W237–W243. <https://doi.org/10.1093/nar/gkv437>.
- Amey RC, Mills PR, Bailey AM, Foster GD. 2003. Investigating the role of a *Verticillium fungicola*  $\beta$ -1,6-glucanase during infection of *Agaricus bisporus* using targeted gene disruption. *Fungal Genet Biol* 39:264–275. [https://doi.org/10.1016/S1087-1845\(03\)00061-6](https://doi.org/10.1016/S1087-1845(03)00061-6).
- Collopy PD, Amey RC, Sergeant MJ, Challen MP, Mills PR, Foster GD, Bailey AM. 2010. The *pmk1*-like mitogen-activated protein kinase from *Lecanicillium (Verticillium) fungicola* is not required for virulence on *Agaricus bisporus*. *Microbiology* 156:1439–1447. <https://doi.org/10.1099/mic.0.034439-0>.