

**Psychological distress during the acceleration phase of the COVID-19 pandemic: a survey of doctors practising in Emergency Medicine, Anaesthesia and Intensive Care Medicine in the United Kingdom and Republic of Ireland**

Tom Roberts<sup>1,2</sup>, Jo Daniels<sup>3</sup>, William Hulme<sup>4</sup>, Robert Hirst<sup>5</sup>, Daniel Horner<sup>1,6</sup>, Mark D Lyttle<sup>2,7</sup>, Katie Samuel<sup>5</sup>, Blair Graham<sup>8,9</sup>, Charles Reynard<sup>10</sup>, Michael J Barrett<sup>11,12</sup>, James Foley<sup>13</sup>, John Cronin<sup>12,14</sup>, Etimbuk Umana<sup>15</sup>, Joao Vinagre<sup>16</sup> and Edward Carlton<sup>1,17</sup> on Behalf of the collaborators of TERN, RAFT, PERUKI, ITERN, TRIC, and SATARN.

Affiliations:

- 1) Royal College of Emergency Medicine, London, UK
- 2) Emergency Department, Bristol Royal Hospital for Children, UK
- 3) Department of Psychology, University of Bath, UK
- 4) Statistical Consultant, Oxford, UK
- 5) Department of Anaesthesia, North Bristol NHS Trust, UK
- 6) Department of Intensive Care and Emergency Department, Salford Royal Hospital NHS Foundation Trust
- 7) Faculty of Health and Applied Sciences, University of the West of England, Bristol
- 8) Faculty of Health, University of Plymouth, Plymouth, UK.
- 9) Emergency Department, University Hospitals Plymouth, UK
- 10) Department of Cardiovascular Sciences, University of Manchester
- 11) Department of Emergency Medicine, Children's Health Ireland at Crumlin, Ireland
- 12) School of Medicine, Women's and Children's Health, University College Dublin, Ireland
- 13) Emergency Department, University Hospital Waterford, Waterford, Ireland
- 14) Department of Emergency Medicine, St Vincent's University Hospital, Dublin, Ireland
- 15) Emergency Department, Connolly Hospital Blanchardstown, Dublin, Ireland.
- 16) College of Anaesthesiologists of Ireland, Dublin, Ireland
- 17) Emergency Department, North Bristol NHS Trust

## Corresponding Author

Dr Tom Roberts

Address: 12 Hamilton Road, Bristol, BS3 1Pb

Email: [Tomkieranroberts@gmail.com](mailto:Tomkieranroberts@gmail.com)

Telephone: 07894234121

Word Count	3493
Figures	7
Tables	4
References	51
Supplementary Material	3

Keywords:

COVID-19

Psychological Distress

GHQ-12

Emergency Medicine

Anaesthetics

Intensive Care Medicine

## ABSTRACT

### Objective

To quantify psychological distress experienced by emergency, anaesthetic and intensive care doctors during the acceleration phase of COVID-19 in the UK and Republic of Ireland.

### Methods

Initial cross-sectional electronic survey distributed during acceleration phase of the first pandemic wave of COVID-19 in the UK and Republic of Ireland (United Kingdom: 18/03/2020 – 26/03/2020 and Ireland: 25/03/2020 – 02/04/2020). Surveys were distributed via established specialty research networks, within a three-part longitudinal study. Participants were Doctors working in emergency, anaesthetic and intensive medicine during the first pandemic wave of COVID-19 in acute hospitals across the United Kingdom and Republic of Ireland. Primary outcome measures were the General Health Questionnaire-12 (GHQ-12). Additional questions examined personal and professional characteristics, experiences of COVID-19 to date, risk to self and others, and self-reported perceptions of health and wellbeing.

**Results** 5440 responses were obtained, 54.3% (n=2955) from Emergency Medicine and 36.9% (n=2005) from Anaesthetics. All levels of doctor seniority were represented. For the primary outcome of GHQ-12 score, 44.2% (n=2405) of respondents scored >3, meeting the criteria for psychological distress. 57.3% (n=3045) had never previously provided clinical care during an infectious disease outbreak but over half of respondents felt somewhat prepared (48.6%, n=2653) or very prepared (7.6%, n=416) to provide clinical care to COVID-19 patients. However, 81.1% (n=4414) either agreed (31.1%, n=2709) or strongly agreed (31.1%, n=1705) that their personal health was at risk due to their clinical role.

**Conclusions** Findings indicate that during the acceleration phase of the COVID-19 pandemic almost half of front-line doctors working in acute care reported psychological distress as measured by the GHQ-12. Findings from this study should inform strategies to optimise preparedness and explore modifiable factors associated with increased psychological distress in the short and long-term.

Word Count: 284

## KEY SUMMARY

### What is known already?

- The COVID-19 outbreak has already placed exceptional demand on healthcare systems globally and is likely to continue to do so for the foreseeable future.
- Emergency and critical care doctors are responsible for the management of severely unwell COVID-19 patients. These doctors may be vulnerable to suffering recognised negative psychological effects associated with infectious disease outbreaks, including absenteeism, impaired occupational performance and long-term health conditions.

### What does this paper add?

- This paper presents key findings from the first phase of a cross-sectional longitudinal survey of practising emergency, anaesthetic and intensive care doctors in UK and Republic of Ireland during the acceleration phase of the first wave of the COVID-19 pandemic.
- The findings report a rate of psychological distress in responders of 44.2%. This work clarifies the extent and severity of cross specialty psychological impact during the early phase of a pandemic.
- These results could be used as a comparison for other studies analysing the psychological impact of infectious disease outbreaks at different timepoints or different regions.

## Introduction

On January 30<sup>th</sup>, 2020, the World Health Organisation (WHO) declared Coronavirus Infectious Disease 2019 (COVID-19) a Public Health Emergency of International Concern. Following subsequent acknowledgment of disease severity, COVID-19 was declared a global pandemic on March 11<sup>th</sup> 2020. [1] Clinical studies have consistently demonstrated high acuity amongst hospitalised patients, with approximately 17% requiring intensive care. [2] In addition, high infection rates have been registered in frontline clinicians, with over 106 fatalities reported in UK NHS healthcare workers by the 12<sup>th</sup> of April 2020 [3] and over 550,000 global deaths reported by July 2020. [4]

The rapidity, scale and severity of the pandemic has placed exceptional demands on acute care globally, and this workforce has faced unprecedented burden in workload intensity and personal health risk. Such demands are likely to impact upon psychological wellbeing, including an increased risk of traumatic stress in both the acute phase and at long term follow up. [5–8] Elevated psychological distress has significant consequences for health workers; increased sickness rates, absenteeism, impaired performance at work and the development of physical health problems are common examples. [9–11] However, studies to date that have focused on the disaster or infectious disease setting have been conducted during peak or following the occurrence of infectious outbreaks, making meaningful comparison with pre-peak incidence very difficult. In the wider literature the reported prevalence of distress during pre-COVID-19 times, has been reported as 28.5% in ICU doctors (n=627) and 44.4% Emergency Medicine Consultants (n=350). [12,13]

Establishing the prevalence of psychological distress, and the associated personal and professional factors, is essential to ensure adequate provision of support and mitigation of adverse effects. Several factors may be associated with poorer psychological outcomes, but these need to be established as relevant in the context of the COVID-19 pandemic. [5,14–17] Prospective longitudinal studies are needed to more fully assess the definitive impact of this major outbreak upon psychological wellbeing. [18]

The COVID-19 Emergency Response Assessment (CERA) study is a 3-part longitudinal study, designed to enhance our understanding of the impact of such events on the workforce, and underpin the development of policy and interventions to meet the needs of those affected. [19] The primary aim of this CERA phase one study is to quantify the degree of psychological distress in emergency medicine (EM), anaesthesia and intensive care medicine (ICM) doctors in the acceleration phase of the first wave of the COVID-19 pandemic in the UK and Republic of Ireland. The secondary aim is to provide a

descriptive synthesis of baseline personal and professional characteristics commonly associated with poorer outcomes related to psychological distress and trauma.

## **Materials and methods**

A quantitative online cross-sectional survey of acute care doctors practising EM, anaesthesia or ICM in the UK and Republic of Ireland. This was the first part of a longitudinal survey to be distributed at pre-planned phases aligned to the Centre for Disease Control and Prevention (CDC) pandemic model: (i) the acceleration phase, (ii) the pandemic peak, and (iii) the deceleration/ recovery phase of the initial COVID-19 pandemic wave in the UK and Ireland. [20] Data were gathered in the acceleration phase between the 18<sup>th</sup> - 28<sup>th</sup> March 2020 in the UK, and 25<sup>th</sup> March - 4<sup>th</sup> April 2020 in the Republic of Ireland. Results are presented in accordance with the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). [21] The protocol for the full three phase longitudinal study is published and available from (<http://dx.doi.org/10.1136/bmjopen-2020-039851>). [22] The study was prospectively registered on an open access platform, ISCTRN registry number 10666798 (<https://doi.org/10.1186/ISRCTN10666798>).

### *Participants and procedure*

Doctors of all grades working in EM, anaesthesia or ICM were invited to participate in the study. Responses excluded from analysis included those from other healthcare professional groups and doctors working outside of EM, anaesthesia and ICM, and doctors working in hospitals based outside of the UK or Republic of Ireland. Participants were invited through a multispecialty collaboration of established UK and Irish acute care research networks, led by the Trainee Emergency Research Network (TERN). These include Research and Audit Federation of Trainees (RAFT), Paediatric Emergency Research in the UK and Ireland (PERUKI), Trainee Research in Intensive Care (TRIC), Irish Trainee Emergency Research Network (I-TERN) and Irish Specialist Anaesthesiology Trainee Audit & Research Network (SATARN). The survey participation link was not shared on wider social media platforms in order to mitigate against duplicate completion and completion by respondents not meeting pre-specified inclusion criteria. Access to the survey link was distributed directly to individual participants in each department or hospital by members of the above research networks working within the same department or hospital. This was achieved using established communication links within departments and hospital. To supplement this strategy RAFT emailed members directly. All participants provided informed electronic consent prior to beginning the survey.

The survey was administered via the Research Electronic Data Capture (REDCap) online platform. [23,24] REDCap is fully compliant with Good Clinical Practice, GDPR and 20 ISO 27001. Data was held securely on secure online server hosted by the University Hospitals Bristol and Weston NHS Foundation Trust, UK. Participants were identifiable through their email address, but this data was only available to the Chief Investigator (TR) and data extracted for analysis was anonymised. Participants could exit the survey at any time if they no longer wished to participate. In this event, data from questions already completed were included for analysis, in line with consent. The recruitment process is detailed further in the protocol. [22]

### *Measures*

The General Health Questionnaire-12 (GHQ-12) is a brief, 12 item self-report measure devised to screen for psychological distress in the general population. [25] It has high specificity and sensitivity, with reliability demonstrated across a range of cultures and populations, [26,27] and has been used in similar studies measuring psychological impact of infectious outbreaks. The GHQ-12 was chosen due to its brevity and suitability for time-poor medical staff. [5,14] The measure assesses current state (rather than long-standing attributes) and asks participants to compare to their own baseline.

Data were also collected on personal and professional factors commonly associated with psychological distress in medical or disaster settings, [5,14–17] derived from a literature review and iterative discussion within the study steering group. Items were included where relevance has previously been established and replicated (for example, factors commonly associated with psychological distress) or where relevance was justified in the context of the current pandemic. Final inclusion was by consensus, underpinned by a requirement for the survey to be sufficiently brief to encourage full and repeated completion (see online supplementary material for a full report of included items, minus the GHQ-12, which has been removed for copyright reasons).

### *Ethical and regulatory approvals*

Ethical approval was provided by the University of Bath (UK), and Children's Health Ireland at Crumlin (Ireland). Regulatory approval was provided by the UK Health Research Authority (ref 218944).

### *Analysis*

Individual study records were checked and validated by the study chief investigator (TR) and statistician (WH) at survey completion; data was excluded in the event of duplicate entry (by email address), absence of consent or non-completion of a predetermined minimum required dataset for

analysis (completion of GHQ-12, grade, department, and hospital). Descriptive statistics relating to personal and professional characteristics are presented overall, and by department and geographic region.

GHQ-12 scores will be presented using two validated methods. [26] The first (bi-modal) method is used to identify a clinical cut off for psychological distress; the second method (Likert-type) is more sensitive to change in psychological distress over time and is most suitable for comparison between different time points. In the bi-modal method, item responses are assigned to the values 0, 0, 1, 1 (from the most positive to the most negative sentiment) and summed to form an aggregate score from zero (least distressed) to 12 (most distressed). A score of more than 3 is indicative of psychological distress. [26] The Likert-type 0-1-2-3 method is also presented. This forms an aggregate score from zero (least distressed) to 36 (most distressed). This method is more sensitive to changes within individuals over time and was included for consistency with subsequent longitudinal analyses using survey data from phases 2 and 3. Distribution of GHQ-12 aggregate scores were described using quartiles, and comparisons between different personal and professional characteristics were made. A descriptive synthesis was used to summarise key findings in relation to the personal and professional characteristics.

All analyses and statistical outputs were produced using the statistical programming language R v3.6.3. [28] Analysis scripts for this study are available on a GitHub repository:  
<https://github.com/wjchulme/TERN-CERA-study>

#### *Patient and Public Involvement*

The research team is primarily made up of frontline doctors from all represented specialties who undertook clinical work throughout the first wave of the COVID-19 pandemic on the frontline.

## Results

Enrolment is summarised in figure 1. The online survey link was accessed 8111 times, of which 5440 (67%) were suitable for analysis. This represents 15.9% of an estimated 34,188 doctors working across EM (11,843), Anaesthetics (20,556) and ICU (1789) in the UK and Ireland (data as per a freedom of information request to the General Medical Council UK and declared numbers by Ireland site leads). The GHQ-12 completion rate was 95.9% (n=5218 / 5440) of participants eligible for analysis. Online supplementary material outlines the adherence to the CHERRIES checklist.

### Sample characteristics

Demographics of the study population are summarised in Table 1 and were similar across all specialties. The median age was 31-35, 50.4% (n=2648) were male, and 37.4% (n= 2033) identified as a Junior Doctor grade.

	All (N=5440)	Anaesthetics (N=2005)	Emergency Medicine (N=2955)	Intensive Care (N=920)
<b>Age</b>				
20-25	204 (3.8%)	5 (0.2%)	182 (6.2%)	17 (1.9%)
26-30	1373 (25.3%)	355 (17.7%)	882 (29.9%)	221 (24.1%)
31-35	1313 (24.2%)	477 (23.8%)	702 (23.8%)	258 (28.1%)
36-40	865 (15.9%)	331 (16.5%)	458 (15.5%)	154 (16.8%)
41-45	659 (12.1%)	277 (13.8%)	337 (11.4%)	85 (9.3%)
46-50	447 (8.2%)	219 (10.9%)	203 (6.9%)	82 (8.9%)
51-55	315 (5.8%)	182 (9.1%)	108 (3.7%)	55 (6.0%)
56-60	174 (3.2%)	102 (5.1%)	56 (1.9%)	31 (3.4%)
61-65	72 (1.3%)	48 (2.4%)	20 (0.7%)	11 (1.2%)
66-70	8 (0.1%)	6 (0.3%)	1 (0.0%)	3 (0.3%)
>70	3 (0.1%)	1 (0.0%)	2 (0.1%)	0 (0.0%)
Missing	7	2	4	3
<b>Gender</b>				
Male	2648 (50.4%)	986 (50.8%)	1421 (49.8%)	490 (55.2%)
Female	2601 (49.5%)	953 (49.1%)	1427 (50.0%)	396 (44.6%)
Other	9 (0.2%)	2 (0.1%)	6 (0.2%)	1 (0.1%)
Missing	182	64	101	33
<b>Seniority</b>				
Junior Doctor	2033 (37.4%)	515 (25.7%)	1308 (44.3%)	327 (35.5%)
Middle Grade Doctor	1254 (23.1%)	463 (23.1%)	658 (22.3%)	248 (27.0%)
Senior Doctor (Consultant Grade)	1694 (31.1%)	892 (44.5%)	676 (22.9%)	284 (30.9%)
Other Senior Doctor	459 (8.4%)	135 (6.7%)	313 (10.6%)	61 (6.6%)
<b>Nation</b>				
England	4310 (79.2%)	1593 (79.5%)	2313 (78.3%)	738 (80.2%)
Northern Ireland	167 (3.1%)	83 (4.1%)	64 (2.2%)	39 (4.2%)
Republic of Ireland	416 (7.6%)	85 (4.2%)	317 (10.7%)	55 (6.0%)
Scotland	367 (6.7%)	120 (6.0%)	228 (7.7%)	47 (5.1%)
Wales	180 (3.3%)	124 (6.2%)	33 (1.1%)	41 (4.5%)
<b>Geographical Region (England)</b>				
East Midlands	303 (5.6%)	138 (6.9%)	133 (4.5%)	47 (5.1%)
East of England	327 (6.0%)	123 (6.1%)	179 (6.1%)	54 (5.9%)

London	818 (15.0%)	201 (10.0%)	560 (19.0%)	88 (9.6%)
North East	210 (3.9%)	73 (3.6%)	112 (3.8%)	47 (5.1%)
North West	596 (11.0%)	246 (12.3%)	270 (9.1%)	128 (13.9%)
South East	629 (11.6%)	196 (9.8%)	402 (13.6%)	84 (9.1%)
South West	686 (12.6%)	279 (13.9%)	318 (10.8%)	126 (13.7%)
West Midlands	340 (6.2%)	146 (7.3%)	161 (5.4%)	78 (8.5%)
Yorkshire and the Humber	401 (7.4%)	191 (9.5%)	178 (6.0%)	86 (9.3%)
<b>Geographical Region (Republic of Ireland)</b>				
Dublin	221 (4.1%)	37 (1.8%)	173 (5.9%)	27 (2.9%)
Rest of Ireland	195 (3.6%)	48 (2.4%)	144 (4.9%)	28 (3.0%)
<b>Redeployed</b>				
No	4920 (90.7%)	1628 (81.4%)	2830 (96.1%)	865 (94.2%)
Yes	506 (9.3%)	373 (18.6%)	116 (3.9%)	53 (5.8%)
Missing	14	4	9	2
<b>GHQ12 (0-1-2-3)</b>				
Median (Q1-Q3)	13 (10-17)	14 (11-18)	13 (10-16)	14 (11-17)
Mean	14.0	14.7	13.5	14.2
Missing	222	86	114	37
<b>GHQ12 (0-0-1-1)</b>				
≤3	2813 (53.9%)	913 (47.6%)	1663 (58.5%)	439 (49.7%)
>3	2405 (46.1%)	1006 (52.4%)	1178 (41.5%)	444 (50.3%)
Missing	222	86	114	37

### *Prevalence of psychological distress*

Analysis of GHQ-12 data indicated that 44.2% (n=2405) of respondents reached study threshold for psychological distress (>3 using 0-0-1-1 method) (figure 2). This was higher in both Anaesthetics (52.5%, n=1006) and ICM (50.3%, n=444) when compared to EM (41.5%, n=1178). The median aggregate GHQ-12 score (using the 0-1-2-3 method) was 13 (Q1-Q3: 10-17) (figure 3). Collated results to the individual GHQ-12 questions items are displayed in figure 4. From this visual representation, the domains of concentration, sleep, being under strain and day to day enjoyment of activities were negatively affected. The highest median GHQ-12 score by grade and department was 15 (Q1-Q3: 11-18) in 'other senior doctors' working in Anaesthetics, compared to the lowest median score of 13 (Q1-Q3: 10-16) found in all four grade cohorts working in EM (figure 5). The GHQ-12 was found to have good internal consistency in this population (Cronbach's alpha=0.846 (95% CI = 0.838–0.853)). GHQ-12 bar charts are available for all items in online supplementary material.

### *Professional Characteristics*

Professional characteristics are summarised in Table 2, with data on all items provided in the online supplementary material. Over half (57.3%, n=3045) reported no prior experience of providing care during infectious disease outbreaks. Although 39.5% (n=2073) reported having no education regarding the clinical care of patients with suspected COVID-19, 48.6% (n= 2643) felt 'somewhat prepared' to do so. A total of 56.2% (n=3058) of respondents reported zero (21%) or low (1-5 cases; 35.2%) direct clinical contact with suspected COVID-19 cases. Only 9.3% (n=506) of participants were redeployed to

other clinical areas, 73.7% (n=373) of those redeployed were from Anaesthetics and the majority of all those redeployed (70.9% (n= 359)) were redeployed to ICM. For those doctors redeployed to another clinical area, the median GHQ-12 was 14 (Q1-Q3: 11-18) compared to 13 (Q1-Q3: 10-17) in those not redeployed (figure 6). The location of redeployment did not make a substantial difference to median GHQ-12 scores (online supplementary material).

<b>Table 2 Professional Factors</b>			
<b>Training and experience</b>			
<b>Have you previously provided direct clinical care to any patients affected by these infectious diseases?*</b>	<b>n</b>	<b>% of total</b>	<b>% of non-missing</b>
None of the below	3045	57.3	48.3
Ebola virus	166	3.1	2.6
MERS-CoV	323	6.1	5.1
SARS	279	5.2	4.4
Chikungunya	152	2.9	2.4
Cholera	160	3.0	2.5
Influenza (swine, avian, zoonotic)	1996	37.5	31.6
Zika virus	80	1.5	1.3
Other	107	2.0	1.7
(Missing)	122	2.2	-
<b>How many suspected cases of Covid-19 have you had direct clinical contact with since March 1<sup>st</sup> 2020?</b>	<b>n</b>	<b>% of total</b>	<b>% of non-missing</b>
0	1144	21.0	22.0
1-5	1914	35.2	36.8
6-10	879	16.2	16.9
11-15	465	8.5	8.9
16-20	325	6.0	6.2
21-25	139	2.6	2.7
26-30	102	1.9	2.0
31-35	25	0.5	0.5
> 36	212	3.9	4.1
(Missing)	235	4.3	-
<b>How confident do you feel in the infection control training that has been provided to you?</b>	<b>n</b>	<b>% of total</b>	<b>% of non-missing</b>
Not confident at all	461	8.5	8.9
Somewhat not confident	1193	21.9	23.0
Neither not confident or confident	1118	20.6	21.5
Somewhat confident	2150	39.5	41.4
Very confident	274	5.0	5.3
(Missing)	244	4.5	-
<b>How prepared do you feel to provide direct care to suspected cases?</b>	<b>n</b>	<b>% of total</b>	<b>% of non-missing</b>
Completely unprepared	195	3.6	3.8
Somewhat unprepared	1365	25.1	26.3
Neither unprepared or prepared	577	10.6	11.1
Somewhat prepared	2643	48.6	50.9
Very prepared	416	7.6	8.0

(Missing)	244	4.5	-
<b>How do you feel the care received by patients who are NOT presenting with either symptoms or a diagnosis of COVID-19 is?</b>	<b>n</b>	<b>% of total</b>	<b>% of non-missing</b>
Significantly worse than before Covid-19	623	11.5	12.0
Slightly worse than before Covid-19	2018	37.1	38.9
The same as before Covid-19	2145	39.4	41.3
Slightly better than before Covid-19	345	6.3	6.6
Significantly better than before Covid-19	59	1.1	1.1
(Missing)	250	4.6	-
*= participants could select more than one option			

Provision of training for the *use of personal and protective equipment (PPE)* was variable (Table 3). 8.2% (n=433) did not receive training in donning and doffing, 17.1% (n=903) had not received formal fit testing for masks and 22.1% (n=1163) had not received PPE training for aerosol generating procedure. The modality of training was variable, with local departmental guidance the most common form of training. In relation to confidence in infection control, 30.4% reported feeling somewhat not confident (21.9%. n=1193) or not confident at all (8.5%, n=461) in their infection control training.

Table 3 PPE Training							
What training have you received in regard to personal protective equipment (PPE) since the COVID-19 outbreak was declared? (select all that apply)							
	No training	Formal instructional video	Written instruction	Simulation training	Departmental guidance	Other	Missing
<b>Donning and doffing (gloves, gown, facemask, eye protection)</b>	8.2% (n=433)	45.8% (n=2421)	42.9% (n=2267)	45.8% (n=2420)	57.8% (3145)	2.1% (n=109)	2.8% (n=155)
<b>Formal fit testing for mask</b>	17.1% (n=903)	14.1% (n=742)	11.3% (n=596)	38.7% (n=2038)	45.9% (n=2499)	9.9% (n=523)	3.2% (n=172)
<b>PPE training for exposure to aerosol generating procedure (e.g. intubation)</b>	22.1% (n=1163)	27.5% (n=1443)	35.0% (n=1838)	38.4% (n=2019)	46.3% (n=2519)	1.8% (n=97)	3.4% (n=185)
% = percentage of total. <b>Note</b> = Participants could select multiple options							

Participants reported highly variable use of information sources for COVID-19 related policy and clinical updates (figure 7). Government and institutional guidelines were the medium most frequently checked on a daily basis (online supplementary material). Social media was checked hourly by 16.3% (n=885) of respondents, whilst 12.8% (n=699) did not access this at all; no other source was characterised by interaction of this frequency. Online blogs and podcasts were checked less frequently; 17.7% (n=962) checked these daily, and 21.8% (n=1186) never used these sources.

### Personal Factors

Personal characteristics are summarised in Table 4, with data on all items provided in the online supplementary material. Of respondents who reported a physical health condition (42.0%, n=2284),

59.4% (n=1357) thought that COVID-19 could worsen their pre-existing condition. Of those with a pre-existing mental health condition (37% n= 2028), 49.0% (n=994) felt the pandemic would exacerbate their symptoms. In the full cohort, 81.1% (n=4414) agreed or strongly agreed that their personal health was at risk during the pandemic due to their clinical role (49.8% and 31.3% respectively). However, the greatest concern was the potential risk to families or loved ones due to their clinical role, with 35.3% (n=1921) “extremely worried” and 43.4% (n=2363) “generally worried”.

#### *Personal experience of COVID-19*

15.3% (n= 833) needed to self-isolate by the time of this first survey, the most common reasons being personal symptoms (55.4%, n=460), and symptomatic household contacts (35.8%, n=279). Only 5.2% (n= 43) of those who had to self-isolate missed more than 10 clinical shifts.

Table 4 <i>Personal Factors</i>			
Personal factors			
<b>Are you concerned that the exposure to the COVID-19 outbreak may increase symptoms of any established medical health condition?</b>	n	% of total	% of non-missing
Yes	1357	24.9	26.2
No	927	17.0	17.9
Prefer not to disclose	75	1.4	1.4
I do not have an established medical condition	2826	51.9	54.5
(Missing)	255	4.7	-
<b>Are you concerned that the exposure to the COVID-19 outbreak may increase symptoms of any established mental health conditions?</b>	n	% of total	% of non-missing
Yes	1034	19.0	20.0
No	994	18.3	19.2
Prefer not to disclose	93	1.7	1.8
I do not have an established mental health condition	3054	56.1	59.0
(Missing)	265	4.9	-
<b>I feel that my personal health is at risk during the COVID-19 outbreak due to my clinical role?</b>	n	% of total	% of non-missing
Strongly disagree	93	1.7	1.8
Disagree	216	4.0	4.2
Neither agree nor disagree	450	8.3	8.7
Agree	2709	49.8	52.4
Strongly agree	1705	31.3	33.0
(Missing)	267	4.9	-
<b>How worried are you about the potential risks to your family, loved one or others due to your clinical role in the COVID-19 outbreak?</b>	n	% of total	% of non-missing
Extremely worried	1921	35.3	37.1
Generally worried	2363	43.4	45.6
Neither worried or not worried	392	7.2	7.6
Generally not worried	414	7.6	8.0
Not worried at all	89	1.6	1.7
(Missing)	261	4.8	-
Personal experience of COVID-19			
<b>Have you had to self-isolate?</b>	n	% of total	% of non-missing

Yes	833	15.3	16.1
No	4339	79.8	83.9
(Missing)	268	4.9	-
<b>For what reason did you have to self-isolate?*</b>	<b>n</b>	<b>% of total</b>	<b>% of non-missing</b>
Personal symptoms	460	55.4	47.1
Personal diagnosis of COVID-19	39	4.7	4.0
Symptoms of a member of the household	297	35.8	30.4
Exposure to a positive case of COVID-19 in the work environment	99	11.9	10.1
Exposure to a positive case of COVID-19 in your personal environment	16	1.9	1.6
Other	65	7.8	6.7
(Missing)	3	0.4	-
<b>How many clinical shifts in your rota have you missed due to self-isolation?</b>	<b>n</b>	<b>% of total</b>	<b>% of non-missing</b>
0	81	9.7	9.8
1	77	9.2	9.3
2	119	14.3	14.3
3	131	15.7	15.8
4	124	14.9	14.9
5-7	196	23.5	23.6
8-10	59	7.1	7.1
>10	43	5.2	5.2
(Missing)	3	0.4	-

\*= participants could select more than one option

## Discussion

In this survey of frontline doctors across the UK and Ireland, over 40% met the criteria for psychological distress, measured by the GHQ-12, during the acceleration phase. These findings are higher than normative data in ICM Doctors and similar to rates found in EM Consultants. [12,13] However, comparison to previous research is limited by sample size, cohort differences and the historical nature of these studies.[29,30]. Figures are from early in the pandemic when clinical exposure and case fatality were low; by the end of the study period there had been 2825 reported COVID-19 cases and 436 reported COVID-19 deaths in the UK [31] and 4014 cases and 131 deaths in Ireland. [32] The data collection period fell during a period of unprecedented and escalating Government restrictions, culminating in a full UK lockdown on 23<sup>rd</sup> March 2020 and 27<sup>th</sup> March 2020 in Ireland – the effect of which cannot be fully accounted for in this work.

Despite efforts to ensure methodological rigour, typical limitations in keeping with survey studies will apply to this study such as response bias and social desirability bias. Whilst data have broadly been captured during the acceleration phase, substantial regional variation in COVID-19 activity was experienced during the survey period, meaning that participants' clinical experience is likely to vary by region. Future phases will attempt to account for this regional variation

As data has been collected during the acceleration phase of the pandemic, this data cannot be considered a true baseline. However, our data does provide findings from an early timepoint in an infection pandemic, which will inform longitudinal studies assessing the significance of psychological impact during peak and deceleration phases. These findings broadly support the role of several previously identified key (and potentially modifiable) stressors during pandemic medicine, including lack of preparedness and training with PPE; elevated concern in relation to risk to self and others, from provision of clinical care to patients with suspected infectious illness; the potential of moral injury through perceptions of worse care provision to other disease states; access to information and communication. [5,14–17]

This study highlights a large increase in rates of distress within the ICM cohort when compared to previous work. [12] Whereas in EM, the rates of distress are similar to a cohort of Consultants previously studied. [13] The comparisons to this research, conducted in 2002, may be limited by the significant changes of service design, delivery and pressures in the intervening years. However, with such a stark difference between the groups the reasons underlying this should be a priority for further research.

Findings are consistent with existing research in the field of infectious diseases and COVID-19. [33–37] Despite fairly low rates of exposure and self-isolation due to physical symptoms, between half and two-thirds of respondents expressed concern that exposure to COVID-19 would worsen their pre-existing physical and mental health conditions. This is unsurprising, given the prominently reported death rates of those with existing medical conditions. [38] Concern regarding infection of family and loved ones was highly prevalent and reported by over 80% of respondents, mirroring findings from a recent interview study examining the content of concerns in frontline healthcare workers. [39] Further research has also indicated that having a family member with COVID-19 may be a predisposing factor to psychological distress for healthcare workers themselves. [5] While concern for others and exposure to COVID-19 is unavoidable in frontline clinicians, the distress associated with it is not; psychological wellbeing warrants careful monitoring and intervention, in line with recommendations by the British Medical Association and British Psychological Society. [40,41]

During this survey, doctors expressed concern that the care of patients without COVID-19 would be negatively impacted. Such concerns have been realised in the literature, with reported increases in out of hospital cardiac arrest rates and anecdotal publications on reduced and increasingly late presentations of reversible disease. [42–44] This has also been observed in previous disease outbreaks, such as Ebola. [45,46] The emotional impact of this is likely to result in feelings such as

guilt, shame, and moral injury, [47] factors commonly associated with poorer psychological outcomes in the context of trauma [15] and worthy of further research in this context.

Availability of PPE to front-line clinicians during the COVID-19 outbreak has been a prominent concern internationally. [48,49] However even where PPE is available, a key driver for related psychological impact is the training, confidence and preparedness in its use. The majority of respondents received some form of training with regards to PPE at an early stage of the pandemic, but this was highly variable, and sometimes entirely documentary rather than practical. Given previous literature suggests poorer psychological outcomes with limited preparedness and confidence, [5] it is of note that the percentage of those respondents receiving no training for different PPE procedures, ranged from less than 10% to 22.1%.

Previous research has indicated that accessing social media as a primary source of information can be problematic and associated with acute and post-traumatic stress, particularly when information is conflicting. [35,50] Whilst it was beyond the scope of this analysis to evaluate any such causal impact, further research should seek to assess the relative impact of social media usage in this context. Findings from any research of this nature would be of potential benefit in informing guidance on content and delivery, and end-user insight, to benefit the psychological wellbeing of clinicians using this source of information.

Current research in the general population reflects our findings of increased distress in doctors. A UK study of 17,452 adults in April 2020, found the prevalence of significant distress (defined by a GHQ-12 of >4) to be 27.3 % (95% CI 26.4-28.2%). [51] This had increased from 18.8% (95% CI 17.8-20.0%) in the 2018-2019 cohort. Whilst comparison to our data is limited by the higher threshold for distress, the trends identified by Pierce *et al* place our results in the context of increased distress in the general population.

Whilst our findings reflect that many doctors struggled with sleep, concentration and feeling strained, many also reported feeling more useful than usual. General confidence, decision-making and sense of worth were reported by respondents to be either better or the same for the vast majority of respondents during the pandemic acceleration phase. Research examining resilience and post-traumatic growth in disaster settings have reported similar findings, particularly a sense of accomplishment and enhanced self-esteem. [14] Despite unprecedented restrictions on individual liberty and freedom of movement, most respondents reported feeling as happy as usual or more so, all things considered. It is a positive indicator to see this early on in the pandemic. Taken together, the findings reflect what may be reasonably expected at an early point in a developing crisis;

elevated psychological distress with a degree of impact on functioning, however protective factors such as increased feelings of worth and usefulness may mitigate against the full impact of the pandemic on mental health. The extent to which a high level of support from the general public towards healthcare professionals influenced feelings of positivity of resilience is unclear and warrants further investigation.

### *Conclusions*

High levels of psychological distress were present amongst UK and Ireland frontline emergency medicine, anaesthesia and intensive care doctors during the acceleration phase of the initial wave of the COVID-19 pandemic. These frontline staff experienced stress and strain, yet faced this with reasonable levels of confidence in preparedness, mobilisation of skills and increased self-worth. Future work will assess the degree and nature of the relationship between personal and professional factors and psychological distress within a longitudinal framework and consider implications for policy and practice.

## References:

- 1 WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020> (accessed 31 Mar 2020).
- 2 Docherty AB, Harrison EM, Green CA, *et al.* Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: Prospective observational cohort study. *BMJ* Published Online First: 2020. doi:10.1136/bmj.m1985
- 3 Cook T. Exclusive: deaths of NHS staff from covid-19 analysed | Comment | Health Service Journal. <https://www.hsj.co.uk/exclusive-deaths-of-nhs-staff-from-covid-19-analysed/7027471.article> (accessed 7 Jul 2020).
- 4 COVID-19 situation update worldwide, as of 10 July 2020. <https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases> (accessed 10 Jul 2020).
- 5 Kisely S, Warren N, McMahon L, *et al.* Occurrence, prevention, and management of the psychological effects of emerging virus outbreaks on healthcare workers: rapid review and meta-analysis. *BMJ* Published Online First: 2020. doi:10.1136/bmj.m1642
- 6 Maunder RG, Lancee WJ, Rourke S, *et al.* Factors associated with the psychological impact of severe acute respiratory syndrome on nurses and other hospital workers in Toronto. *Psychosom. Med.* 2004. doi:10.1097/01.psy.0000145673.84698.18
- 7 Pappa S, Ntella V, Giannakas T, *et al.* Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav Immun* Published Online First: 2020. doi:10.1016/j.bbi.2020.05.026
- 8 Allan SM, Bealey R, Birch J, *et al.* The prevalence of common and stress-related mental health disorders in healthcare workers based in pandemic-affected hospitals: a rapid systematic review and meta-analysis. 2020. doi:10.1101/2020.05.04.20089862
- 9 Arora M, Asha S, Chinnappa J, *et al.* Review article: Burnout in emergency medicine physicians. *EMA - Emerg. Med. Australas.* 2013;**25**:491–5. doi:10.1111/1742-6723.12135
- 10 Maunder RG, Lancee WJ, Balderson KE, *et al.* Long-term psychological and occupational effects of providing hospital healthcare during SARS outbreak. *Emerg Infect Dis* Published Online First: 2006. doi:10.3201/eid1212.060584

- 11 Halpern J, Maunder RG, Schwartz B, *et al.* Identifying risk of emotional sequelae after critical incidents. *Emerg Med J* Published Online First: 2011. doi:10.1136/emj.2009.082982
- 12 Coomber S, Todd C, Park G, *et al.* Stress in UK intensive care unit doctors. *Br J Anaesth* Published Online First: 2002. doi:10.1093/bja/aef273
- 13 Burbeck R, Coomber S, Robinson SM, *et al.* Occupational stress in consultants in accident and emergency medicine: A national survey of levels of stress at work. *Emerg Med J* 2002;**19**:234–8. doi:10.1136/emj.19.3.234
- 14 Brooks SK, Dunn R, Amlôt R, *et al.* A Systematic, Thematic Review of Social and Occupational Factors Associated with Psychological Outcomes in Healthcare Employees during an Infectious Disease Outbreak. *J Occup Environ Med* Published Online First: 2018. doi:10.1097/JOM.0000000000001235
- 15 Brooks SK, Dunn R, Sage CAM, *et al.* Risk and resilience factors affecting the psychological wellbeing of individuals deployed in humanitarian relief roles after a disaster. *J. Ment. Heal.* 2015. doi:10.3109/09638237.2015.1057334
- 16 Lancee WJ, Maunder RG, Goldbloom DS. Prevalence of psychiatric disorders among Toronto hospital workers one to two years after the SARS outbreak. *Psychiatr Serv* Published Online First: 2008. doi:10.1176/ps.2008.59.1.91
- 17 Ozer EJ, Best SR, Lipsey TL, *et al.* Predictors of posttraumatic stress disorder and symptoms in adults: A meta-analysis. *Psychol. Bull.* 2003;**129**:52–73. doi:10.1037/0033-2909.129.1.52
- 18 BMA survey finds doctors' lives still at risk despite PPE pledges.  
<https://www.bma.org.uk/news-and-opinion/bma-survey-finds-doctors-lives-still-at-risk-despite-government-pledges-on-ppe> (accessed 23 Apr 2020).
- 19 Holmes EA, O'Connor RC, Perry VH, *et al.* Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *The Lancet Psychiatry* 2020;**0**. doi:10.1016/S2215-0366(20)30168-1
- 20 Holloway R, Rasmussen SA, Zaza S, *et al.* Updated Preparedness and Response Framework for Influenza Pandemics. 2014.<https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6306a1.htm> (accessed 8 Apr 2020).
- 21 Eysenbach G. Improving the Quality of Web Surveys: The Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res* 2004;**6**:e34. doi:10.2196/jmir.6.3.e34

- 22 Roberts T, Daniels J, Hulme W, *et al.* COVID-19 emergency response assessment study: a prospective longitudinal survey of frontline doctors in the UK and Ireland: study protocol. *BMJ Open* Published Online First: 2020. doi:10.1136/bmjopen-2020-039851
- 23 Harris PA, Taylor R, Thielke R, *et al.* Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;**42**:377–81. doi:10.1016/J.JBI.2008.08.010
- 24 Harris PA, Taylor R, Minor BL, *et al.* The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 2019;**95**:103208. doi:10.1016/J.JBI.2019.103208
- 25 Goldberg D, Williams P. *A user's guide to the General Health Questionnaire*. London: : GL Assessment 1988.
- 26 Goldberg DP, Gater R, Sartorius N, *et al.* The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychol Med* Published Online First: 1997. doi:10.1017/S0033291796004242
- 27 Goldberg DP, Oldehinkel T, Ormel J. Why GHQ threshold varies from one place to another. *Psychol Med* Published Online First: 1998. doi:10.1017/S0033291798006874
- 28 R Core Team. *A Language and Environment for Statistical Computing*. R Found. Stat. Comput. 2018.
- 29 Imo UO. Burnout and psychiatric morbidity among doctors in the UK: A systematic literature review of prevalence and associated factors. *BJPsych Bull* Published Online First: 2017. doi:10.1192/pb.bp.116.054247
- 30 Kinman G, Teoh K. What could make a difference to the mental health of UK doctors? A review of the research evidence. 2018.
- 31 Coronavirus (COVID-19) in the UK. <https://coronavirus.data.gov.uk/#category=regions&map=rate> (accessed 17 Jun 2020).
- 32 Health Protection Surveillance Centre I. Epidemiology of COVID-19 in Ireland - daily reports, April 2020. <https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/casesinireland/epidemiologyofcovid-19inireland/april2020/> (accessed 26 Oct 2020).
- 33 Mukhtar S. Psychological health during the coronavirus disease 2019 pandemic outbreak. *Int.*

- J. Soc. Psychiatry. 2020. doi:10.1177/0020764020925835
- 34 Shevlin, M., McBride, O., Murphy, J., Miller, J. G., Hartman, T. K., Levita, L., ... & Bennett KM. Anxiety, Depression, Traumatic Stress, and COVID-19 Related Anxiety in the UK General Population During the COVID-19 Pandemic. *UK Popul Ment Heal Covid-19* 2020.
- 35 Torales J, O'Higgins M, Castaldelli-Maia JM, *et al.* The outbreak of COVID-19 coronavirus and its impact on global mental health. *Int J Soc Psychiatry* 2020;:002076402091521. doi:10.1177/0020764020915212
- 36 Wang C, Pan R, Wan X, *et al.* A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun* Published Online First: 2020. doi:10.1016/j.bbi.2020.04.028
- 37 Chen N, Zhou M, Dong X, *et al.* Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;**395**:507–13. doi:10.1016/S0140-6736(20)30211-7
- 38 Jeanne Ross, Weill Peter R, David. OpenSAFELY: factors associated with COVID-19-related hospital death in the linked electronic health records of 17 million adult NHS patients. *J Chem Inf Model* Published Online First: 2006. doi:10.1017/CBO9781107415324.004
- 39 Shanafelt T, Ripp J, Trockel M. Understanding and Addressing Sources of Anxiety among Health Care Professionals during the COVID-19 Pandemic. *JAMA - J Am Med Assoc* Published Online First: 2020. doi:10.1001/jama.2020.5893
- 40 BMA. The mental health and wellbeing of the medical workforce-now and beyond COVID-19. 2020. <https://www.bma.org.uk/media/2475/bma-covid-19-and-nhs-staff-mental-health-wellbeing-report-may-2020.pdf>
- 41 British Psychological Society Covid19 Staff Wellbeing Group. The psychological needs of healthcare staff as a result of the Coronavirus pandemic. [www.ics.ac.uk/ICS/Education/Wellbeing/ICS/Wellbeing.aspx](http://www.ics.ac.uk/ICS/Education/Wellbeing/ICS/Wellbeing.aspx) (accessed 7 Jul 2020).
- 42 Baldi E, Sechi GM, Mare C, *et al.* Out-of-Hospital Cardiac Arrest during the Covid-19 Outbreak in Italy. *N Engl J Med* Published Online First: 2020. doi:10.1056/nejmc2010418
- 43 Garcia S, Albaghdadi MS, Meraj PM, *et al.* Reduction in ST-Segment Elevation Cardiac Catheterization Laboratory Activations in the United States During COVID-19 Pandemic. *J. Am. Coll. Cardiol.* 2020. doi:10.1016/j.jacc.2020.04.011

- 44 Marijon E, Karam N, Jost D, *et al.* Out-of-hospital cardiac arrest during the COVID-19 pandemic in Paris, France: a population-based, observational study. *Lancet Public Heal* Published Online First: 2020. doi:10.1016/S2468-2667(20)30117-1
- 45 Ribacke KJB, Saulnier DD, Eriksson A, *et al.* Effects of the West Africa Ebola virus disease on health-care utilization - A systematic review. *Front Public Heal* Published Online First: 2016. doi:10.3389/FPUBH.2016.00222
- 46 Elston JWT, Cartwright C, Ndumbi P, *et al.* The health impact of the 2014–15 Ebola outbreak. *Public Health*. 2017. doi:10.1016/j.puhe.2016.10.020
- 47 Greenberg N, Docherty M, Gnanapragasam S, *et al.* Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ*. 2020. doi:10.1136/bmj.m1211
- 48 Rowan NJ, Laffey JG. Challenges and solutions for addressing critical shortage of supply chain for personal and protective equipment (PPE) arising from Coronavirus disease (COVID19) pandemic – Case study from the Republic of Ireland. *Sci Total Environ* Published Online First: 2020. doi:10.1016/j.scitotenv.2020.138532
- 49 Dyer C. Covid-19: doctors are warned not to go public about PPE shortages. *BMJ* Published Online First: 2020. doi:10.1136/bmj.m1592
- 50 Jones NM, Thompson RR, Schetter CD, *et al.* Distress and rumor exposure on social media during a campus lockdown. *Proc Natl Acad Sci U S A* Published Online First: 2017. doi:10.1073/pnas.1708518114
- 51 Pierce M, Hope H, Ford T, *et al.* Mental health before and during the COVID-19 pandemic: a longitudinal probability sample survey of the UK population. *The Lancet Psychiatry* Published Online First: 2020. doi:10.1016/S2215-0366(20)30308-4

#### Acknowledgements

The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, the Department of Health or the Royal Colleges involved in survey distribution.

The authors would like to acknowledge Mai Baquedano, at the University of Bristol, for her support with REDCap and GL Assessment for providing the licence for the GHQ-12 free of charge.

#### Author Contributions

Tom Roberts (TR) conceived the idea for the study. TR, Edd Carlton (EC), Jo Daniels (JD), Mark Lyttle (ML), and Blair Graham (BG) were responsible for the initial study design, which was refined with the

help of Katie Samuel (KS), Charles Reynard (CR), Robert Hirst (RH), Michael Barrett (MB) and William Hulme (WH). Expert advice on psychological assessment scores was provided by JD. WH provided the statistical plan. TR lead the dissemination of the study in UK Adult Emergency Departments (ED), ML lead the dissemination of the study in UK and Ireland Paediatric EDs, KS lead the dissemination of the study in UK Anaesthetic and ICU Departments, MB lead the dissemination of the study in Ireland EDs, along with John Cronin, James Foley and Etimbuk Umana. Joao Vinagre lead the dissemination in Ireland ICUs and Anaesthetic Departments. TR coordinated study set-up, finalisation of the study surveys and finalisations of study protocols. All authors contributed to the final study design and protocol development, critically revised successive drafts of the manuscript and approved the final version. The study management group is responsible for the conduct of the study.

## Funding

The Chief Investigator is directly funded as a research fellow by the Royal College of Emergency Medicine. The GHQ-12 is being used under licence from GL assessment; the fee for use of this instrument within all three surveys has been waived. Dr Carlton is a National Institute for Health Research Advanced Fellow.

The study has direct funding from RCEM. Grant code: G/2020/1

## Competing Interests

Many of the authors have been working as frontline clinicians during the COVID-19 pandemic. They have no competing interests to declare.

## Trainee Emergency Research Network (TERN) Collaborators:

L	Kane	Aberdeen Royal Infirmary	A	Hormis	Rotherham Hospital
L	Mackenzie	Addenbrooke's Hospital, Cambridge	R	Hannah	Royal Alexandra Children's Hospital
S	Sharma Hajela	Addenbrooke's Hospital, Cambridge	A	Corfield	Royal Alexandra Hospital, Scotland
J	Phizacklea	Addenbrooke's Hospital, Cambridge	J	Maney	Royal Belfast Hospital for Sick Children
K	Malik	Addenbrooke's Hospital, Cambridge	D	Metcalfe	Royal Berkshire Hospital
N	Mathai	Aintree University Hospital	S	Timmis	Royal Berkshire Hospital
A	Sattout	Aintree University Hospital	C	Williams	Royal Bolton Hospital
S	Messahel	Alder Hey Children's Hospital, Liverpool	R	Newport	Royal Bolton Hospital
E	Fadden	Alder Hey Children's Hospital, Liverpool	D	Bawden	Royal Cornwall Hospital
R	McQuillan	Altnagelvin Area Hospital, N. Ireland	A	Tabner	Royal Derby Hospital
B	O'Hare	Antrim Area Hospital, N. Ireland	H	Malik	Royal Devon and Exeter Hospital
S	Lewis	Arrowe Park Hospital, Merseyside	C	Roe	Royal Devon and Exeter Hospital
D	Bewick	Barnsley Hospital	D	McConnell	Royal Devon and Exeter Hospital
R	Taylor	Bath Royal United Hospital	F	Taylor	Royal Free London
I	Hancock Manthalapo	Bath Royal United Hospital	R	Ellis	Royal Glamorgam Hospital, Wales
D	Ramesh Babu	Bedford Hospital, Bedfordshire	S	Morgan	Royal Gwent Hospital, Wales
S	Hartshorn	Birmingham Children's Hospital	L	Barnicott	Royal Hampshire County Hospital

M	Williams	Birmingham Children's Hospital	S	Foster	Royal Hospital for Children, Glasgow
A	Charlton	Bradford Royal Infirmary	J	Browning	Royal Hospital for Sick Children Royal Hospital for Sick Children, Edinburgh
L	Somerset	Bristol Royal Hospital for Children	L	McCrae	Royal Infirmary Hospital, Edinburgh
C	Munday	Bristol Royal Hospital for Children	E	Godden	Royal Infirmary Hospital, Edinburgh
A	Turner	Bristol Royal Hospital for Children	A	Saunders	Royal Infirmary Hospital, Edinburgh
R	Sainsbury	Bristol Royal Infirmary	A	Lawrence-Ball	Royal Liverpool University Hospital
E	Williams	Bristol Royal Infirmary	R	House	Royal Liverpool University Hospital
S	Patil	Chelsea & Westminster Hospital	J	Muller	Royal London Hospital
R	Stewart	Chelsea & Westminster Hospital	I	Skene	Royal London Hospital
M	Winstanley	Chelsea & Westminster Hospital	M	Lim	Royal London Hospital
N	Tambe	Chesterfield Royal hospital	H	Millar	Royal Manchester Children's Hospital
C	Magee	City Hospital, Birmingham	A	Rai	Royal Manchester Children's Hospital
D	Raffo	Craigavon Area Hospital, N. Ireland	K	Challen	Royal Preston Hospital
D	Mawhinney	Craigavon Area Hospital, N. Ireland	S	Currie	Royal Preston Hospital
B	Taylor	Cumberland Infirmary, Cumbria	M	Elkanzi	Royal Stoke University Hospital
T	Hussan	Darlington Memorial Hospital	T	Perry	Royal Surrey County Hospital
G	Pells	Darlington Memorial Hospital	W	Kan	Royal Surrey County Hospital
F	Barham	Derriford Hospital, Plymouth	L	Brown	Royal Sussex County Hospital
F	Wood	Derriford Hospital, Plymouth	M	Cheema	Royal Sussex County Hospital
C	Szekeres	East Surrey Hospital	A	Clarey	Royal Victoria Hospital
R	Greenhalgh	East Surrey Hospital	A	Gulati	Royal Victoria Infirmary
S	Marimuthu	Eastbourne Hospital	K	Webster	Royal Victoria Infirmary
R	Macfarlane	Epsom and St Helier Hospitals	A	Howson	Salford Royal NHS Foundation Trust
M	Alex	Evelina Children's Hospital, London	R	Doonan	Salford Royal NHS Foundation Trust
B	Shrestha	Frimley Park Hospital	C	Magee	Sandwell Hospital
L	Stanley	Gloucester Royal Hospital	A	Trimble	Sheffield Children's Hospital
J	Gumley	Gloucester Royal Hospital	C	O'Connell	Sheffield Children's Hospital
K	Thomas	Gloucester Royal Hospital Great North Children's Hospital, Newcastle	R	Wright	Southampton General
M	Anderson	Gloucester Royal Hospital Great North Children's Hospital, Newcastle	E	Colley	Southmead Hospital, Bristol
C	Weegenaar	Great Western Hospital, Swindon	C	Rimmer	Southport Hospital, Merseyside
J	Lockwood	Harrogate Hospital	S	Pintus	Southport Hospital, Merseyside
T	Mohamed	Heartlands's Hospital, Birmingham	H	Jarman	St George's Hospital, London
S	Ramraj	Hillingdon Hospital, London	V	Worsnop	St George's Hospital, London
M	Mackenzie	Homerton Hospital, London	S	Collins	St Helier Hospital
A	Robertson	Homerton Hospital, London	M	Colmar	St John's Hospital, Livingston
W	Niven	Homerton Hospital, London	N	Masood	St John's Hospital, Livingston
M	Patel	Homerton Hospital, London	R	McLatchie	St John's Hospital, Livingston
S	Subramaniam	Horton General Hospital, Banbury	A	Peasley	Stepping Hill Hospital
C	Holmes	Huddersfield Royal Infirmary	S	Rahman	Stoke Mandeville Hospital South Tyneside and Sunderland NHS Foundation Trust.
S	Bongale	Inverclyde Royal Hospital	N	Mullen	Stoke Mandeville Hospital South Tyneside and Sunderland NHS Foundation Trust.
U	Bait	Ipswich Hospital	L	Armstrong	The Royal Berkshire Hospital, Reading
S	Nagendran	Ipswich Hospital	A	Hay	The Whittington Hospital, London
S	Rao	Ipswich Hospital	R	Mills	The Whittington Hospital, London
F	Mendes	James Paget Hospital, Great Yarmouth	J	Lowe	Torbay Hospital, Devon
P	Singh	John Radcliffe Hospital, Oxford	H	Raybould	Torbay Hospital, Devon

S	Subramaniam	John Radcliffe Hospital, Oxford	A	Ali	Torbay Hospital, Devon
T	Baron	John Radcliffe Hospital, Oxford	P	Cuthbert	Ulster Hospital Dundonald, Northern Ireland
C	Ponmani	King George Hospital	A	McKinney	Ulster Hospital Dundonald, Northern Ireland
M	Depante	King's College Hospital, London	S	Taylor	University College London Hospital
R	Sneep	King's College Hospital, London	V	Talwar	University College London Hospital
A	Brookes	King's College Hospital, London	Z	Al-Janabi	University Hospital Ayr, Scotland
S	Williams	King's College Hospital, London	C	Leech	University Hospital Coventry and Warwickshire
A	Rainey	King's College Hospital, London	J	Turner	
J	Brown	Kingston Hospital, London	L	McKechnie	University Hospital Crosshouse, Scotland
N	Marriage	Kingston Hospital, London	B	Mallon	University Hospital Crosshouse, Scotland
S	Manou	Leeds General Infirmary	J	McLaren	University Hospital Crosshouse, Scotland
S	Hart	Leeds General Infirmary	Y	Moulds	University Hospital Crosshouse, Scotland
M	Elsheikh	Leeds General Infirmary	L	Dunlop	University Hospital Hairmyres, Scotland
L	Cocker	Leicester Royal Infirmary	F	Burton	University Hospital Hairmyres, Scotland
M	Elwan	Leicester Royal Infirmary	M	Keers	University Hospital Lewisham, London
H	Vincent	Leicester Royal Infirmary	S	Robertson	University Hospital Lewisham, London
K L	Nunn	Leicester Royal Infirmary	L	Craver	University Hospital Lewisham, London
C	Sarja	Lister Hospital, Stevenage	D	Moultrie	University Hospital Monklands, Scotland
N	Viegas	Luton & Dunstable Hospital	N	Williams	University Hospital of North Tees
M	Graham	Mater Hospital, Belfast	O	Purvis	University Hospital of North Tees
S	Wooffinden	Manchester Royal Infirmary	S	Clark	University Hospital of North Tees
E	Reynard	Manchester Royal Infirmary	M	Davies	University Hospital of Wales, Cardiff
C	Cherian	Manchester Royal Infirmary	C	Foreman	University Hospital of Wales, Cardiff
N	Da-Costa	Medway NHS Foundation Trust	S	Ngua	University Hospital of Wales, Cardiff
A	Duckitt	Medway NHS Foundation Trust	C	Morgan	University Hospital of Wales, Cardiff
S	Bailey	Milton Keynes University Hospital	J	Hoskins	University Hospital of Wales, Cardiff
J	How	Milton Keynes University Hospital	N	Fryer	University Hospital Southampton
L	Hine	Milton Keynes University Hospital	J	Wright	University Hospital Southampton
T	Ihsan	Milton Keynes University Hospital	R	Frost	University Hospital Southampton
F	Abdullah	Milton Keynes University Hospital	L	Ellis	University Hospital Southampton
H	Bader	Milton Keynes University Hospital	P	Mackay	University Hospital Wishaw, Scotland
K	Pradhan	Milton Keynes University Hospital	A	Gray	Victoria Hospital, Kirkcaldy, Scotland
S	Manoharan	Milton Keynes University Hospital	K	Jacobs	Watford General Hospital
M	Kehler	Wolverhampton NHS Trust	M	Muslim	
L	Muswell	Newham University Hospital, London	I	Veettil Asif	West Middlesex university hospital
R	Bonsano	Newham University Hospital, London	P	Amiri	West Middlesex university hospital
M	Evans	Norfolk and Norwich University Hospitals	S	Shrivastava	West Middlesex university hospital
J	Christmas	North Hampshire Hospital, Basingstoke	F	Raza	West Middlesex university hospital
E	Knight	North Middlesex Hospital, London	S	Wilson	Wexham Park Hospital
K	O'Rourke	North Tees Hospital, Stockton on Tees	M	Riyat	Wexham Park Hospital
L	Adeboye	North Tees Hospital, Stockton on Tees	H	Knott	Wexham Park Hospital
K	Iftikhar	Northern General Hospital, Sheffield	M	Ramazany	Whiston Hospital, Merseyside
K	Evans	Northern General Hospital, Sheffield	S	Langston	Whiston Hospital, Merseyside
R	Darke	Northumbria Specialist Emergency Care Hospital	N	Abela	Whiston Hospital, Merseyside
R			L	Robinson	Whittington Hospital, London

R	Freeman	Northumbria Specialist Emergency Care Hospital	D	Maasdorp	Whittington Hospital, London
E	Grocholski	Northwick Park Hospital, London	H	Murphy	Whittington Hospital, London
K	Kaur	Peterborough City Hospital	H	Edmundson	Whittington Hospital, London
H	Cooper	Peterborough City Hospital	R	Das	Whittington Hospital, London
M	Mohammad	Princess Royal University Hospital, London	c	orjioke	Whittington Hospital, London
L	Harwood	Princess Royal University Hospital, London	D	Worley	Whittington Hospital, London
K	Lines	Queen Alexandra Hospital, Portsmouth	W	Collier	Whittington Hospital, London
C	Thomas	Queen Alexandra Hospital, Portsmouth	J	Everson	Whittington Hospital, London
D	Ranasinghe	Queen Alexandra Hospital, Portsmouth	N	Maleki	Whittington Hospital, London
S	Hall	Queen Elizabeth Hospital	A	Stafford	Whittington Hospital, London
J	Wright	Queen Elizabeth Hospital	S	Gokani	Whittington Hospital, London
S	Hall	Queen Elizabeth Hospital	M	Charalambos	Whittington Hospital, London
N	Ali	Queen Elizabeth Hospital	A	Olajide	Whittington Hospital, London
J	Hunt	Queen Elizabeth Hospital, Birmingham	C	Bi	Whittington Hospital, London
H	Ahmad	Queen Elizabeth Queen's mother hospital Margate	J	Ng	Whittington Hospital, London
C	Ward	Queen Elizabeth University Hospital, Glasgow	S	Naeem	William Harvey Hospital, Kent
M	Khan	Queens Medical Centre, Nottingham	A	Hill	Wythenshawe Hospital, Manchester
K	Holzman	Redhill Hospital, Surrey	C	Boulind	Yeovil District Hospital
J	Ritchie	Rotherham Hospital			

#### Ireland Trainee Emergency Research Network (I-TERN) Collaborators:

A	Patton	Cork University Hospital, Ireland
M	Jee Poh Hock	Galway Hospital, Ireland
J	Lynch	Tallaght University Hospital, Ireland
R	O'Sullivan	Bon Secours Hospital Cork, Ireland
S	Gilmartin	Children's Health Ireland at Crumlin, Ireland
S	Uí Bhroin	Children's Health Ireland at Tallaght, Ireland
P	Fitzpatrick	Children's Health Ireland at Temple Street, Ireland
A	Patton	Cork University Hospital, Ireland
S	Kukaswadia	Mercy University Hospital, Ireland
C	Prendergast	Midlands Regional Hospital Tullamore, Ireland
A	Ahmed	Sligo University Hospital, Ireland
C	Dalla Vecchia	St Vincent's University Hospital, Ireland
M	Grummell	Tallaght University Hospital, Ireland
I	Grossi	University Hospital Limerick, Ireland
B	MacManus	University Hospital Waterford, Ireland

Research and Audit Federation of Trainees (RAFT), Trainee Research in Intensive Care (TRIC) and Specialist Anaesthesia Trainee led Audit and Research Network (SATURN) Collaborators:

K	Samuel	North Bristol NHS Trust
A	Boyle	Royal Victoria Hospital, Belfast
A	Waite	Royal Liverpool University Hospital
B	Johnston	University of Liverpool
J	Vinagre	Children's Health Ireland at Temple Street
P	Turton	Arrowe Park Hospital, Merseyside
D	George	University Hospital of Wales, Cardiff
C	Battle	Morrison Hospital, Wales
J	Anandarajah	Wrexham Maelor Hospital, Wales