

1 **Distribution of the branched-chain  $\alpha$ -ketoacid dehydrogenase complex E1 $\alpha$**   
2 **subunit and glutamate dehydrogenase in the human brain and their role in**  
3 **neuro-metabolism<sup>†</sup>**

4 **Jonathon Hull<sup>#‡</sup>, Marcela Usmari Moraes<sup>#‡</sup>, Emma Brookes<sup>‡</sup>, Seth Love<sup>\*</sup>, and**  
5 **Myra E. Conway<sup>†</sup>**

6 <sup>‡</sup>Faculty of Health and Life Sciences, University of the West of England, Coldharbour  
7 Lane, Bristol, BS16 1QY, UK.

8 <sup>\*</sup>Dementia Research Group, Faculty of Medicine and Dentistry, University of Bristol,  
9 Bristol, BS16 1LE, UK.

10 <sup>†</sup>This study was supported by a bursary from BRACE (Bristol Research into  
11 Alzheimer's and Care of the Elderly) awarded to M.E.C. at the University of the West  
12 of England.

13 <sup>#</sup>These authors contributed equally to this publication.

14

15 <sup>\*</sup>Address correspondence: Myra E. Conway, Faculty of Health and Life Sciences,  
16 University of the West of England, Coldharbour Lane, Bristol, BS16 1QY, UK Tel: 0044  
17 117 328 3552; Fax: 0044 117 328 2904; email: [myra.conway@uwe.ac.uk](mailto:myra.conway@uwe.ac.uk).

18

1 **Abstract**

2 Glutamate is the major excitatory neurotransmitter of the central nervous system, with  
3 the branched-chain amino acids (BCAAs) acting as key nitrogen donors for *de novo*  
4 glutamate synthesis. Despite the importance of these major metabolites, their  
5 metabolic pathway in the human brain is still not well characterised. The metabolic  
6 pathways that influence the metabolism of BCAAs have been well characterised in rat  
7 models. However, the expression of key proteins such as the branched-chain  $\alpha$ -  
8 ketoacid dehydrogenase (BCKD) complex and glutamate dehydrogenase isozymes  
9 (GDH) in the human brain is still not well characterised. We have used specific  
10 antibodies to these proteins to analyse their distribution within the human brain and  
11 report, for the first time, that the E1 $\alpha$  subunit of the BCKD is located in both neurons  
12 and vascular endothelial cells. We also demonstrate that GDH is localised to  
13 astrocytes, although vascular immunolabelling does occur. The labelling of GDH was  
14 most intense in astrocytes adjacent to the hippocampus, in keeping with glutamatergic  
15 neurotransmission in this region. GDH was also present in astrocyte processes  
16 abutting vascular endothelial cells. Previously, we demonstrated that the branched-  
17 chain aminotransferase (hBCAT) proteins were most abundant in vascular cells  
18 (hBCATm) and neurons (hBCATc). Present findings are further evidence that BCAAs  
19 are metabolised within both the vasculature and neurons in the human brain. We  
20 suggest that GDH, hBCAT and the BCKD proteins operate in conjunction with  
21 astrocytic glutamate transporters and glutamine synthetase to regulate the availability  
22 of glutamate. This has important implications given that the dysregulation of glutamate  
23 metabolism, leading to glutamate excitotoxicity, is an important contributor to the  
24 pathogenesis of several neurodegenerative conditions such as Alzheimer's disease.

25

26 **Keywords:** BCKD, GDH, BCAA, glutamate.

27 **Running title:** BCKD and GDH in the human brain.

28 **Acknowledgements:** This study was supported by a grant from Bristol Research into  
29 Alzheimer's and Care of the Elderly (BRACE) to M.E.C. The University of the West of  
30 England, Bristol, BS16 1QY, UK. The authors declare there are no conflicts of interest  
31 involved in the study.

## 1 **Abbreviations**

2 AD, Alzheimer's disease; BCAA, branched-chain amino acids; hBCAT, human  
3 branched-chain aminotransferase; BCKA, branched-chain  $\alpha$ -keto acid; BCKD,  
4 branched-chain  $\alpha$ -keto acid dehydrogenase complex; CA, cornu ammonis area; DAB,  
5 3,3'-diaminobenzidine; DAPI, 4',6-Diamidino-2-Phenylindole; DPX, distyrene  
6 plasticiser xylene; EDTA, ethylenediaminetetraacetic acid; GABA, gamma-amino  
7 butyric acid; GDH, glutamate dehydrogenase; GFAP, glial fibrillary acidic protein;  
8 hBCATc, human cytosolic branched-chain aminotransferase; hBCATm, human  
9 mitochondrial branched-chain aminotransferase; PBS, phosphate buffered saline; RT,  
10 room temperature; VWF, Von Willebrand factor.


11

## 1 Introduction

2 The three-essential branched chain amino acids (leucine, isoleucine and valine) are  
3 key nitrogen contributors for the *de novo* synthesis of glutamate. Studies on the rat  
4 retina demonstrated that inhibition of the cytosolic branched-chain aminotransferase  
5 (BCATc) decreased *de novo* glutamate synthesis by 30% (LaNoue *et al.*, 2001; Lieth  
6 *et al.*, 2001). In humans, the first step in BCAA metabolism is catalysed by the  
7 branched-chain aminotransferases (hBCAT, EC 2.6.1.42), mitochondrial (hBCATm)  
8 and cytosolic (hBCATc). These proteins mediate the transfer of the  $\alpha$ -amino group  
9 from the BCAA to  $\alpha$ -ketoglutarate to produce glutamate and the respective branched-  
10 chain  $\alpha$ -keto acid (BCKA – ketoisocaproate, ketomethylvalerate and ketoisovalerate;  
11 Scheme 1, reaction 1). Studies in human tissue have demonstrated hBCATm  
12 expression within the vasculature, with hBCATc predominantly expressed in neurons  
13 (Hull *et al.*, 2012). The second step in BCAA metabolism is considered the rate-limiting  
14 step, as the BCKAs are irreversibly oxidised by the branched-chain  $\alpha$ -keto acid  
15 dehydrogenase (BCKD, EC 1.2.4.4) complex. This produces branched-chain acyl Co-  
16 A derivatives (Scheme 1, reaction 2), generating the Krebs' cycle substrate acetyl-  
17 CoA, and the Krebs' cycle intermediate succinyl-CoA (Harris *et al.*, 1986; Hutson *et*  
18 *al.*, 2005).

19 The BCKD complex is primarily regulated by phosphorylation and dephosphorylation  
20 of the E1 subunit by the BCKD kinase and BCKD phosphatase proteins, with  
21 phosphorylation inactivating the complex (Damuni *et al.*, 1984; Damuni *et al.*, 1987;  
22 Popov *et al.*, 1991; Popov *et al.*, 1992; Shimomura *et al.*, 1990; Shimomura *et al.*,  
23 2001; Wynn *et al.*, 2000). The distribution of the BCKD complex has yet to be  
24 determined in the human brain. However, work in rat has demonstrated BCKD-E1 $\alpha$   
25 within neurons, where it is thought to contribute to the BCAA-BCKA shuttle and energy  
26 production (Cole *et al.*, 2012). The activity of the BCKD complex is influenced by the  
27 activity of the two glutamate dehydrogenase isoforms (GDH1 and GDH2; EC 1.4.1.3),  
28 which catalyse the conversion of glutamate into  $\alpha$ -ketoglutarate (Scheme 1, reaction  
29 3) (Islam *et al.*, 2010).

### Scheme 1



1

2 GDH1 is expressed in the liver and was previously detected in astrocytes within the  
3 human brain. Conversely, the GDH2 protein is absent from liver and expressed in the  
4 human testes, kidney and brain, where it was also observed in astrocytes in the human  
5 cortex (Nissen *et al.*, 2016; Spanaki *et al.*, 2010; Spanaki *et al.*, 2014; Spanaki &  
6 Plaitakis, 2012). Both GDH isoforms are considered mitochondrial, however GDH  
7 activity has also been associated with the nucleus (di Prisco *et al.*, 1968; Lai *et al.*,  
8 1986) and the endoplasmic reticulum (Colon *et al.*, 1986; Lee *et al.*, 1999) of rat tissue.  
9 The GDH proteins link glutamate metabolism with the Krebs's cycle, and whilst the  
10 reaction is potentially reversible, it is accepted that in the brain this reaction is  
11 predominantly directed to metabolise glutamate (Lorin *et al.*, 2013; Yudkoff *et al.*,  
12 1994; Li *et al.*, 2012). GDH is regulated through many factors including the availability  
13 of purine nucleotides (ADP, ATP, GTP and NADH), steroid hormones and L-leucine.  
14 GTP is the predominant negative regulator of GDH1, activating the protein when cell  
15 energy production is reduced. GDH2 lacks such a regulator and is constitutively active  
16 in a metabolically active cell (Spanaki *et al.*, 2010). The predominant positive  
17 regulators of both enzymes are ADP and L-leucine (increasing activity by up to 14-fold  
18 and 9.7-fold, respectively), resulting in acute sensitivity of these proteins to the activity  
19 of hBCAT and the BCKD complex (Erecinska & Nelson, 1990; Mastorodemos *et al.*,  
20 2005; Plaitakis *et al.*, 2013; Smith & Stanley, 2008; Tomita *et al.*, 2011).

21 Previous work by Islam *et al.* (2007, 2010) using pure protein and rat models,  
22 demonstrated that hBCATm is able to form a protein complex with the BCKD subunit  
23 E1 $\alpha$  and GDH1. This complex allows channelling of the BCKAs to the E1 $\alpha$  subunit,  
24 increasing BCKA decarboxylation 12-fold. The GDH1 protein then binds to the  
25 pyridoxamine-5'-phosphate form of BCATm and increases the recycling of the BCAT  
26 cofactor pyridoxal-5'-phosphate, also replenishing  $\alpha$ -ketoglutarate (Islam *et al.*, 2007;

1 Islam *et al.*, 2010). Despite the importance of BCAA and glutamate metabolism in the  
2 human brain, it has not been established whether a similar process occurs within this  
3 tissue. Establishing a complete understanding of BCAA and glutamate metabolism is  
4 of particular relevance to diseases where these metabolites contribute to the  
5 pathological process, such as maple syrup urine disease and Alzheimer's disease,  
6 respectively.

7 Our aim in this study was to examine the distribution of the GDH protein and the BCKD  
8 complex in the human brain, to ascertain whether the previously described BCAA  
9 shuttle (Hull *et al.*, 2012) could be extended to include these key metabolic proteins.  
10 This previous work provided the first evidence of transaminases (hBCATm) in the  
11 endothelial layer operating as a support network to astrocytes mediating the fine-  
12 tuning of glutamate homeostasis. We further proposed that under normal  
13 physiological conditions, the BCAAs are taken up by neuronal cells, where hBCATc  
14 controlled metabolism acts as a pathway to replenish the glutamate pool.

15 For the first time, the BCKD complex E1 $\alpha$  subunit and the GDH proteins were mapped  
16 to the human brain. In agreement with findings of Cole *et al.*, (2012) in the rat, we  
17 report that BCKD complex E1 $\alpha$  is in neurons, widely distributed throughout the brain  
18 in a variety of neuronal types. However, in addition to observations in the rat, our  
19 findings highlight the expression of the BCKD complex E1 $\alpha$  in endothelial cells  
20 throughout the brain vasculature. In line with previous work (Spanaki *et al.*, 2014),  
21 immunolabelling for GDH was predominantly expressed in astrocytes; however,  
22 labelling was also evident within the vasculature with weak labelling of cerebellar  
23 Purkinje cells. We discuss the impact of these findings with respect to the human  
24 BCAA-BCKA shuttle and the regulation of glutamate production in the human brain.

# 1 **Materials and Methods**

## 2 **Materials**

3 **Antibodies.** Rabbit polyclonal antibodies to BCKD-E1 $\alpha$  (ab126173) and GDH1/GDH2  
4 (ab154027), goat polyclonal secondary antibody to rabbit Alexafluor® 488 (ab150077)  
5 and mouse Alexafluor® 568 (ab175473), mouse monoclonal antibody to glial fibrillary  
6 acidic protein (GFAP), a marker of astrocytes (ab10062) and  $\beta$ 3-tubulin, a neuronal  
7 marker (ab78078) were purchased from Abcam (Cambridge, UK). Mouse monoclonal  
8 antibody to von Willebrand factor (vWF), a marker of vascular endothelial cells  
9 (M0616), was purchased from Dako (Cambridge, UK).

10 **Immunohistochemistry.** Haematoxylin was purchased from SurgiPath  
11 (Peterborough, UK). 4',6-Diamidino-2-Phenylindole (DAPI)-containing hard-set  
12 mounting medium, peroxidase substrate 3,3'-diaminobenzidine (DAB), goat serum (S-  
13 1000), biotin-labelled secondary antibody raised to IgG, and avidin (Vectastain ABC  
14 kit) were purchased from Vector labs (Peterborough, UK). All other materials were  
15 purchased from Fisher Scientific (Loughborough, UK).

16

## 17 **Tissue preparation**

18 The study was approved by North Somerset and South Bristol Research Ethics  
19 Committee. All brain tissue used in this study was from brains donated to the South  
20 West Dementia Brain Bank, University of Bristol. The right hemi-brain had been fixed  
21 in 10% buffered formalin for 3 weeks before tissue blocks were cut and embedded in  
22 paraffin wax for detailed neuropathological assessment. For this study, we examined  
23 18 brains, from people who were older than 60 years of age, had no history of  
24 dementia, and minimal or no neuropathological abnormalities as determined by Braak  
25 stage (Table 1) and small vessel disease score (Supplementary Table 1).

26 Serial sections 7  $\mu$ m in thickness were taken from multiple regions of the brain,  
27 including the frontal, temporal, parietal and occipital lobes, basal ganglia, midbrain,  
28 cerebellum and pons, and collected on APES coated glass slides. Sections were

1 incubated at 60°C overnight to aid adhesion prior to immunolabelling as described  
2 below.

3

#### 4 **Immunohistochemistry**

5 The sections were dewaxed in histoclear (2 x 5 min) and dehydrated in 100% ethanol  
6 (2 x 3 min). Endogenous peroxidase was quenched in 0.09% hydrogen  
7 peroxide/methanol solution for 30 min at room temperature (RT). The slides were pre-  
8 treated with citrate buffer (10 mM sodium citrate, 0.05% Tween 20, pH 6.0). The slides  
9 were then washed (2 x 3 min) in phosphate buffered saline (PBS – 0.154 M NaCl, 1.86  
10 mM NaH<sub>2</sub>PO<sub>4</sub>·2H<sub>2</sub>O, 7.48 mM Na<sub>2</sub>HPO<sub>4</sub>·12H<sub>2</sub>O, pH 7.1) and non-specific binding sites  
11 were blocked with 10% horse serum in PBS (20 min, RT). Sections were incubated  
12 overnight (20 h, 4°C) with primary antibody (1:3000 for GDH1/GDH2 and BCKD-E1α)  
13 in PBS containing 3% Marvel (GDH) or 3% BSA (BCKD-E1α). The sections were  
14 washed in PBS (2 x 3 min) then incubated with biotinylated antibody to IgG for 20 min  
15 (Vectastain ABC kit). The slides were washed again in PBS (2 x 3 min) and incubated  
16 with the avidin-biotin complex in PBS for 20 min (Vectastain ABC kit). Slides were  
17 developed with DAB/H<sub>2</sub>O<sub>2</sub> in distilled water (DAB substrate kit) for 10 min prior to  
18 immersion in copper sulphate solution (16 mM CuSO<sub>4</sub>·5H<sub>2</sub>O, 0.123 M NaCl) for 4 min,  
19 and subsequently counterstained with Harris's haematoxylin (25% Gill haematoxylin).  
20 The slides were dehydrated in 100% ethanol (2 x 5 min), cleared in 100% clearene (2  
21 x 3 min) and mounted in Clearium. Sections were viewed and imaged on a Nikon  
22 Eclipse 80i.

23

#### 24 **Immunofluorescence**

25 The sections were dewaxed in histoclear (2 x 5 min) and dehydrated in 100% ethanol  
26 (2 x 5 min), followed by deionised water (2 x 5 min). Sections were pre-treated with  
27 EDTA (1 mM, pH 8.0) for 30 min at 95°C, and washed (2 x 5 min) in deionised water  
28 followed by PBS (2 x 5 min). Non-specific binding sites were blocked with 10% goat  
29 blocking serum in PBS for 20 min at RT. Slides were incubated overnight (20 h, 4°C)  
30 with primary antibodies (1:500 for β3-tubulin; 1:100 for BCKD-E1α; 1:500 for



1 GDH1/GDH2; 1:500 for GFAP; 1:25 for VWF) in PBS containing 1% goat serum. The  
2 slides were subsequently washed in PBS (2 x 5 min) and incubated with secondary  
3 antibody (1:100 for Alexafluor® 488 and Alexafluor® 568) for 1 h, washed in deionised  
4 water (2 x 5 min) and further blocked with Sudan black (0.3% Sudan black in 70%  
5 ethanol) for 2 min. Slides were rinsed 8 times with PBS and washed in PBS (2 x 5  
6 min), mounted with DAPI and allowed to set for 30 min. Images were acquired using  
7 Andor IQ software and a Nikon Eclipse TE300.

8

## 1 **Results**

2 Antibody specificity for BCKD-E1 $\alpha$  and GDH was tested by western blot analysis on  
3 pure protein and human brain homogenates (Figure 1A-B). No cross reactivity with  
4 unrelated proteins was detected at the antibody concentrations used in these  
5 experiments. For commercial antibodies raised to BCKD-E1 $\alpha$  and GDH, a negative  
6 antibody control was included with each experiment. For further confirmation, antigen  
7 absorption was also performed (at 200X molar excess) during IHC (Figure 1C-F).  
8 Labelling was completely prevented by prior incubation of the antibody with antigen.  
9 The figures are representative of the labelling observed in all 18 brains (Table 1).

10

### 11 **Distribution of BCKD-E1 $\alpha$ in the human brain**

12 Labelling of BCKD-E1 $\alpha$  was predominantly in neuropil and neurons throughout the  
13 brain (Figure 2A). However, endothelial cells were also labelled (Figure 2B). Certain  
14 astrocytic populations (notably Bergmann astrocytes) showed labelling but this was  
15 not a consistent feature (4/8 brains examined). BCKD-E1 $\alpha$  was detected in all regions  
16 of the brain examined, in both neurons and endothelial cells. Labelling of neurons was  
17 stronger than that of endothelial cells.

18 Within the hippocampus, neuronal labelling was strongest within the cornu ammonis  
19 area 4 (CA4), and weaker in successive CA regions through to CA1 (Figure 3A-C). In  
20 the cerebral cortex, larger pyramidal cells were strongly immunopositive (Figure 3D,  
21 large arrows). Vasculature labelling was observed throughout the cerebral cortex  
22 (Figure 3E) with only weak astrocytic labelling (9/18 brains examined), in the vicinity  
23 of the hippocampus and within the white matter (images not shown). The  
24 subependymal and the ependymal tissue were immunopositive (Figure 3F).

25 The molecular layer of the cerebellar cortex was diffusely immunopositive, with  
26 stronger labelling of the cell bodies of many Purkinje cells (Figure 4A-C). Weak  
27 labelling was observed within the granule cell layer (Figure 4C). The white matter was  
28 unlabelled apart from some immunopositive capillaries (Figure 4D-F) and small glial  
29 cells (probably oligodendrocytes). Strong, exclusively neuronal labelling was  
30 observed within the dentate and fastigial nuclei (Figure 5A-C). In the deep grey matter,

1 the neuropil was weakly labelled within the putamen, globus pallidus and  
2 hypothalamus. As previously noted with respect to hBCATc, neurons of the  
3 hypothalamus, raphe nuclei, nucleus basalis of Meynert, pontine nuclei and locus  
4 coeruleus were immunopositive for BCKD-E1 $\alpha$  (Figure 5D-F).

5

## 6 **Distribution of GDH in the human brain**

7 GDH immunoreactivity was expressed in astrocytes throughout the human brain  
8 (Figure 6A), with labelling of the astrocytic foot processes adjacent to the vasculature,  
9 in addition to the vasculature itself (Figure 6B-C). Whilst GDH and GFAP were co-  
10 localised, some astrocytic processes labelled for GDH but not GFAP. This might be  
11 expected, as GFAP is located primarily in the larger processes and not those  
12 enveloping synapses where you would find GDH (Kosaka & Hama, 1986). Most  
13 labelling for GDH was stippled/granular within the cytoplasm of the cell, in keeping  
14 with the mitochondrial restriction demonstrated in previous work (Spanaki *et al.*, 2014).  
15 Not all astrocytes labelled for GDH but labelling was detected in all sections and areas  
16 studied.

17 Within the hippocampus, GDH labelling was entirely astrocytic (Figure 7A-C). There  
18 was also labelling of astrocytes in the neocortex and white matter (Figure 7D-E), but  
19 labelling in the white matter was less dense than that in the cortex. The subpial  
20 feltwork of astrocytic processes was strongly immunopositive for GDH (Figure 7F) in  
21 all brain regions, not just in the cerebrum. Neurons were largely unlabelled (16/18  
22 brains examined), apart from occasional pyramidal neurons in the vicinity of blood  
23 vessels (Figure 8A). The labelling of perivascular astrocytic processes was evident  
24 throughout the cerebral cortex and cerebellum (Figure 8B); in places, the labelling also  
25 involved the adjacent endothelium (Figure 8C).

26 Within the cerebellum (Figure 9A), there was punctate labelling in the processes of  
27 cells within the Purkinje cell layer, most likely to be Bergmann astrocytes (Figure 9B-  
28 C). The Purkinje cells themselves were unlabelled (12/18 brains examined) or weakly  
29 immunopositive (Figure 9B-C, small arrows). Within the granule cell layer, rosette like  
30 structures (glomeruli) were immunopositive. Within the deep cerebellar nuclei, there

1 was strong granular immunopositivity in small glial cells in contrast to the largely  
2 immunonegative neurons (Figure 9D-E).

3 In addition to the granular/stippled cytoplasmic labelling of astrocytes in the cerebral  
4 cortex and deep grey matter structures, there was GDH immunopositivity of some glial  
5 nuclei within these regions (most notably in the putamen) (Figure 9F, arrows). Whilst  
6 none of the brains was from a donor with dementia, in one brain GDH labelling  
7 occurred in conjunction with what appeared to be an amyloid plaque.

8

### 9 **Summary of distribution of BCKD-E1 $\alpha$ and GDH in the human brain**

10 In summary, labelling of BCKD-E1 $\alpha$  was associated with the neuronal cells and the  
11 vasculature in all brain regions studied, reminiscent of the distribution of hBCATc and  
12 hBCATm proteins (Hull *et al.*, 2012). However, the labelling of blood vessels was  
13 weaker and less consistent than that of neurons. GDH was detected in all brain  
14 regions studied and was mainly within astrocytes. There was strong labelling of GDH  
15 in regions containing glutamatergic neurons – likely associated with  
16 neurotransmission within this region. Furthermore, close association of GDH positive  
17 astrocytic processes with the vasculature and in the deep nuclei of the cerebellum  
18 suggest additional roles for this protein.

19

## 1 Discussion

2 The BCKD complex and the GDH isoenzymes play a key role in brain glutamate  
3 metabolism. However, to date most of the information we have of their distribution  
4 within the brain is limited to data from small animal studies. Previous work from this  
5 group has already demonstrated key differences between the expression of BCATm  
6 within the human brain and that in lower mammals such as rats (Cole *et al.*, 2012; Hull  
7 *et al.*, 2012). Current findings indicate that within the human brain, BCKD-E1 $\alpha$  is  
8 neuronal, supporting its proposed role in neuronal BCAA metabolism. GDH was  
9 primarily localised to astrocyte populations of the human brain, as in rodents  
10 (Subbalakshmi & Murthy, 1985; Zaganas *et al.*, 2001; Zaganas *et al.*, 2012). However,  
11 we have shown that, BCKD-E1 $\alpha$  and GDH can also be detected within the vasculature  
12 of the human brain. These findings add complexity to our interpretation of the roles of  
13 these proteins in BCAA and glutamate metabolism.

14 Our understanding of the contribution of BCAAs to glutamate production is based on  
15 *ex vivo* and *in vivo* animal model [<sup>15</sup>N] studies, in which BCAAs (particularly leucine)  
16 contributed approximately 25% of the nitrogen for glutamate synthesis (Yudkoff *et al.*,  
17 1983; Kanamori *et al.*, 1998). We have previously demonstrated that hBCAT was  
18 strongly expressed within the human brain (Hull *et al.*, 2012). The hBCATc enzyme  
19 was localised to neurons, whereas hBCATm expression was associated with the  
20 vasculature. We proposed that, in the human brain, hBCATc metabolises BCAAs for  
21 the transamination of  $\alpha$ -ketoglutarate to glutamate, with hBCATm re-transaminating  
22 glutamate to cycle back into BCAAs. The hBCATc-mediated transfer of NH<sub>3</sub> for  
23 production of glutamate would require further metabolism of the BCKAs to prevent  
24 their re-transamination. In muscle, it is thought that the expression of minimally active  
25 BCKD results in up to 80% re-transamination of BCKAs, the remainder of which are  
26 predominantly exported rather than completely metabolised to Krebs cycle  
27 intermediates (Matthews *et al.*, 1981; Nissen & Haymond, 1981; Suryawan *et al.*,  
28 1998).

29 As in the rat study of Cole *et al.*, (2012), the majority of BCKD-E1 $\alpha$  labelled neurons  
30 were either glutamatergic (e.g. hippocampal granule cells), GABAergic (e.g. Purkinje  
31 cells) or cholinergic (e.g. motor neurons). Reamination of BCKAs is well defined in  
32 many organs (Harper *et al.*, 1984; Staten *et al.*, 1984); however, our work also raises

1 the potential for complete BCAA metabolism within neurons expressing BCKD. This  
2 is supported by previous documentation of very low levels (<1 mM) of BCKAs within  
3 the human brain (Keen *et al.*, 1993; Matsuo *et al.*, 1993), reflecting the formation of  
4 glutamate from BCAAs and the complete oxidation of BCKAs to the Kreb's cycle  
5 substrate acetyl-CoA, and the Kreb's cycle intermediate succinyl-CoA. The labelling  
6 of BCKD-E1 $\alpha$  varied in different parts of the cerebrum but was particularly strong within  
7 the hippocampus. The strong labelling of BCKD-E1 $\alpha$  in the CA4 region and weaker  
8 labelling towards the CA1 region mirrors the distribution of hBCATc (Hull *et al.*, 2012).  
9 BCKD-E1 $\alpha$  was primarily restricted to the somatic region of nerve cells, suggesting  
10 that the cell body is the primary site of BCKA oxidation. In the human brain, BCKD-  
11 E1 $\alpha$  expression was also noted within the vasculature, unlike in the rodent model. The  
12 metabolism of BCAAs within the vasculature may be similar to that in muscle, i.e.  
13 incomplete metabolism to BCKAs, which either re-transaminate glutamate or are  
14 released for further metabolism by other cells (e.g. neurons). However, the expression  
15 of hBCATm, BCKD and GDH within vascular cells raises the possibility that complete  
16 metabolism of BCAAs occur, which could generate ATP for the active transport  
17 required in this cell type. This would be in keeping with the reported active metabolon  
18 described by Islam *et al.*, (2007 & 2010), in which hBCATm, the BCKD complex and  
19 GDH1 work to metabolise BCAAs completely and restore the Kreb's cycle substrate  
20 acetyl-CoA, and the Kreb's cycle intermediate succinyl-CoA.

21 The absence of BCKD and hBCAT labelling within astrocytes indicates that BCAA  
22 oxidation is neuronal rather than astrocytic, and that this oxidation is largely complete.  
23 These findings do not rule out the possibility of translocation of BCKAs to neurons but  
24 make it unlikely that those BCKAs are produced by astrocytes. Skeletal muscle is  
25 thought to be the predominant source of BCKAs metabolised in the liver, due to the  
26 relatively reduced expression of hBCAT within this organ (Suryawan *et al.*, 1998).  
27 Suryawan *et al.*, (1998) also noted that the BCKD in the human brain is predominantly  
28 active, with 59% of the expressed protein maintained in the active state. Translocation  
29 of BCKAs from the bloodstream would depend on their transport across the blood-  
30 brain-barrier (BBB). The transport of ketoacids (e.g.  $\alpha$ -ketoglutarate) across the BBB  
31 occurs via the organic anion transporter family – and it is possible the BCKAs are  
32 transported in this manner (Riedmaier *et al.*, 2012).

1 Work by our group showed hBCATc expressed in magnocellular neurosecretory cells,  
2 the substantia nigra, the nucleus basalis of Meynert and the raphe nuclei (Hull *et al.*,  
3 2012). The expression of BCKD closely mirrors that of hBCATc within these cell types,  
4 supporting a role for hBCATc in cells that secrete a range of neurotransmitters and  
5 peptides. The action of leucine as a potent secretagogue has already been reported  
6 for the hormone insulin, where ketoisocaproate but not leucine, required  
7 transamination to stimulate insulin secretion (Xu *et al.*, 2001; Zhou *et al.*, 2010).  
8 Regulation of leucine levels is likely to be important in regulating neurosecretory  
9 function, and the contribution of BCAA metabolism to this regulation warrants further  
10 investigation.

11 Biochemical studies on mouse, rat and human tissue have detected GDH  
12 activity/immunoreactivity in both neurons and glia (Hohnholt *et al.*, 2017;  
13 Subbalakshmi & Murthy, 1985; Zaganas *et al.*, 2001; Zaganas *et al.*, 2012). Rothe *et*  
14 *al.*, (1994) estimated that GDH activity in neurons is approximately 15% of that in  
15 astrocytes and was crucial for neurons to metabolise glutamine-derived carbon during  
16 glucose deprivation (Hohnholt *et al.*, 2017). Our work confirms the expression of GDH  
17 within many astrocytes, accounting for the predominance of glial activity; the detection  
18 within neurons was more selective and the immunolabelling weaker. However,  
19 Burbaeva *et al.*, (2002) purified three isoforms of GDH protein from human brain  
20 tissue, and it is possible that additional GDH reactions are occurring within the brain  
21 in glia or neurons through the activity of a different GDH isoform.

22 Scant consideration has been given within the literature to GDH activity within the  
23 vasculature. The study of Helms *et al.*, (2012) was one of the first to consider the  
24 potential for glutamate metabolism within endothelial cells, corroborated by our study.  
25 The expression of both hBCATm and GDH suggests that the vasculature has the  
26 potential to metabolise BCAAs fully for energy, or to metabolise glutamate for recycling  
27 of  $\alpha$ -ketoglutarate and BCAAs. In health, the brain is protected from variations in  
28 plasma glutamate by restriction of entry of glutamate across the BBB. However,  
29 glutamate efflux does occur and is affected by blood glutamate level (Hosoya *et al.*,  
30 1999; Gottlieb *et al.*, 2003; Zlotnik *et al.*, 2008; Teichberg *et al.*, 2009; Uchida *et al.*,  
31 2011; Campos *et al.*, 2012; Zlotnik *et al.*, 2012). The ability of the vasculature to  
32 metabolise glutamate may occur as a supplementary mechanism to the glutamate-

1 glutamine cycle, whereby the hBCATm-BCKD-GDH metabolon works to metabolise  
2 BCAAs fully except in circumstances of excess glutamate (i.e. glutamate toxicity).  
3 Increased expression of hBCAT was previously observed in Alzheimer's disease,  
4 perhaps as a response to an increase in glutamate (Ashby *et al.*, 2015; Hull *et al.*,  
5 2015a). Conversely, if the metabolon exists only on the abluminal surface, it would  
6 function solely for the disposal of glutamate.

7 The direction of GDH metabolism will be determined by substrate concentration and  
8 the expression of other key metabolic proteins. The predominant action of GDH within  
9 the human brain is for the production of  $\alpha$ -ketoglutarate from glutamate, allowing  
10 glutamate to contribute carbon to the Krebs's cycle within the brain (McKenna *et al.*,  
11 2016; Nissen *et al.*, 2015). This is likely to be of particular importance within astrocytic  
12 processes surrounding the vasculature, where ATP generation through glutamate  
13 metabolism may provide the energy required for active transport across the BBB  
14 (Figure 10). Additionally, ammonia ( $\text{NH}_3$  - lipid soluble) produced in this reaction may  
15 then leave the brain; this process, together with glutamine export, constitutes a major  
16 mechanism by which the brain disposes of excess ammonia (Hawkins *et al.*, 2002).  
17 In astrocytes, the ammonia may also be incorporated into the structure of glutamate  
18 to form glutamine, through the work of the glutamine synthetase enzyme. This  
19 incorporation into glutamine is unlikely to be a major process at the astrocyte-  
20 vasculature boundary but is part of the glutamate-glutamine cycle in the astrocyte cell  
21 body (Figure 10).

22 The subcellular location of GDH is predominantly mitochondrial; however, we also  
23 found GDH immunopositivity within the nucleus of some cells. Lai *et al.*, (1986)  
24 previously observed GDH activity in both the mitochondrial and nuclear fractions of rat  
25 brain homogenate and our observations indicate that GDH is present within a subset  
26 of nuclei in the human brain as well. Nuclear GDH was observed in a small proportion  
27 of glial cells, mainly within the putamen. This nuclear presence raises interesting  
28 questions about additional functions of GDH, of which two have been proposed. Rajas  
29 *et al.*, (1996) noted that membrane-bound GDH in pig liver had microtubule-binding  
30 activity and suggested that GDH was involved in microtubule-dependent lysosome  
31 formation. Purohit *et al.*, (2013) reported that GDH acts as a histone H3-specific



1 protease in chicken liver, a process that relates directly to the regulation of gene  
2 expression.

3 Another proposed function of GDH is in the regulation of autophagy, for which the  
4 relationship between GDH and leucine is important. In a HeLa knockdown of GDH1,  
5 leucine failed to inhibit autophagy whilst inhibiting mTORC1 activity (Lorin *et al.*, 2013).  
6 The authors proposed that GDH1 regulates autophagy (at least in part) through the  
7 modulation of mTOR but may also contribute to the production of ROS scavengers  
8 and Krebs cycle intermediates (further inhibiting autophagy). We previously found  
9 hBCAT to be absent from astrocytes in the human brain, in contrast to findings in the  
10 rat (Hull *et al.*, 2012; Hull *et al.*, 2015b). It is of note that astrocytes in the human brain  
11 lack the capacity to metabolise BCAAs, including leucine, a key activator of GDH. The  
12 expression of glutamine-leucine counter-transporters on the surface of astrocytes  
13 (Bak *et al.*, 2006), raises the possibility that astrocytes can serve as a partial leucine  
14 sink. This would maintain high intracellular astrocyte levels of leucine to (1) supply  
15 neurons with leucine for glutamate production and (2) support astrocytic GDH activity  
16 for the continuous metabolism of glutamate for the production of free ammonia (to  
17 either be used in the production of glutamine or exported out of the brain) and  $\alpha$ -  
18 ketoglutarate (to be used for the production of ATP).

19

20 In summary, this work builds upon previous work from this group (Hull *et al.*, 2012)  
21 and provides further insight to the metabolism of BCAAs within the human brain and  
22 their association with the glutamate-glutamine cycle. We propose that under normal  
23 physiological conditions, neurons are the main site of BCAA metabolism, where  
24 glutamate is replenished through hBCATc transamination and BCKD activity controls  
25 re-transamination of the BCKAs. Metabolism of the BCAAs may also occur within the  
26 vasculature, likely contributing to the buffering of glutamate levels within the brain. We  
27 propose that the vasculature may also be a site at which the hBCATm-BCKD-GDH  
28 metabolon described by Islam *et al.*, (2007 & 2010) exists *in vivo*. Without the  
29 expression of hBCAT or the BCKD complex in astrocytes (Hull *et al.*, 2012), it is  
30 unlikely that BCAAs are metabolised within this cell type. Our findings have  
31 implications for the understanding of neurotransmitter metabolism within the human

1 brain, and consequently of neurological diseases in which there is dysregulation of  
2 glutamate or BCAA homeostasis.

3

## 1 **References**

- 2 Ashby, E.L., Kierzkowska, M., Hull, J., Kehoe, P.G., Hutson, S.M. and Conway, M.E.  
3 (2017) Altered expression of human mitochondrial branched chain aminotransferase  
4 in dementia with Lewy bodies and vascular dementia. *Neurochemical Research*,  
5 **42(1)**, pp. 306-319.
- 6 Bak, L.K., Schousboe, A. and Waagepetersen, H.S. (2006) The glutamate/GABA-  
7 glutamine cycle: aspects of transport, neurotransmitter homeostasis and ammonia  
8 transfer. *Journal of Neurochemistry*, **98(3)**, pp. 641-653.
- 9 Burbaeva, G.S., Turishcheva, M.S., Vorobyeva, E.A., Savushkina, O.K., Tereshkina,  
10 E.B. and Boksha, I.S. (2002) Diversity of glutamate dehydrogenase in human brain.  
11 *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, **26(3)**, pp. 427-  
12 435.
- 13 Campos, F., Sobrino, T., Ramos-Cabrer, P. and Castillo, J. (2012) Oxaloacetate: a  
14 novel neuroprotective for acute ischemic stroke. *The International Journal of*  
15 *Biochemistry & Cell Biology*, **44(2)**, pp. 262-265.
- 16 Cole, J.T., Sweatt, A.J. and Hutson, S.M. (2012) Expression of mitochondrial  
17 branched-chain aminotransferase and  $\alpha$ -keto-acid dehydrogenase in rat brain:  
18 implications for neurotransmitter metabolism. *Frontiers in Neuroanatomy*, **6(18)**, p. 1-  
19 18.
- 20 Colon, A.D., Plaitakis, A., Perakis, A., Berl, S. and Clarke, D.D. (1986) Purification and  
21 characterization of a soluble and a particulate glutamate dehydrogenase from rat  
22 brain. *Journal of Neurochemistry*, **46(6)**, pp. 1811-1819.
- 23 Damuni, Z., Merryfield, M.L., Humphreys, J.S. and Reed, L.J. (1984) Purification and  
24 properties of branched-chain alpha-keto acid dehydrogenase phosphatase from  
25 bovine kidney. *Proceedings of the National Academy of Sciences*, **81(14)**, pp. 4335-  
26 4338.
- 27 Damuni, Z.A.H.I. and Reed, L.J. (1987) Purification and properties of the catalytic  
28 subunit of the branched-chain alpha-keto acid dehydrogenase phosphatase from  
29 bovine kidney mitochondria. *Journal of Biological Chemistry*, **262(11)**, pp. 5129-5132.

- 1 Di Prisco, G., Banay-Schwartz, M. and Strecker, H.J. (1968) Glutamate  
2 dehydrogenase in nuclear and mitochondrial fractions of rat liver. *Biochemical and*  
3 *Biophysical Research Communications*, **33(4)**, pp. 606-612.
- 4 Erecińska, M. and Nelson, D. (1990) Activation of glutamate dehydrogenase by  
5 leucine and its nonmetabolizable analogue in rat brain synaptosomes. *Journal of*  
6 *Neurochemistry*, **54(4)**, pp. 1335-1343.
- 7 Gottlieb, M., Wang, Y. and Teichberg, V.I. (2003) Blood-mediated scavenging of  
8 cerebrospinal fluid glutamate. *Journal of Neurochemistry*, **87(1)**, pp. 119-126.
- 9 Harper, A.E., Miller, R. and Block, K.P. (1984) Branched-chain amino acid metabolism.  
10 *Annual Review of Nutrition*, **4(1)**, pp. 409-454.
- 11 Harris, R.A., Paxton, R., Goodwin, G.W., Kuntz, M.J., Shimomura, Y. and Han, A.  
12 (1986) Regulation of branched-chain amino acid metabolism. *Biochemical Society*  
13 *Transactions*, **14(6)**, pp. 1005-1008.
- 14 Hawkins, R.A., Peterson, D.R. and Vina, J.R. (2002) The Complementary Membranes  
15 Forming the Blood-Brain Barrier. *The International Union of Biochemistry and*  
16 *Molecular Biology Life*, **54(3)**, pp. 101-107.
- 17 Helms, H.C., Madelung, R., Waagepetersen, H.S., Nielsen, C.U. and Brodin, B. (2012)  
18 *In vitro* evidence for the brain glutamate efflux hypothesis: Brain endothelial cells  
19 cocultured with astrocytes display a polarized brain-to-blood transport of glutamate.  
20 *Glia*, **60(6)**, pp. 882-893.c
- 21 Hohnholt, M.C., Andersen, V.H., Andersen, J.V., Christensen, S.K., Karaca, M.,  
22 Maechler, P. and Waagepetersen, H.S. (2017) Glutamate dehydrogenase is essential  
23 to sustain neuronal oxidative energy metabolism during stimulation. *Journal of*  
24 *Cerebral Blood Flow & Metabolism*, accepted manuscript.
- 25 Hosoya, K.I., Sugawara, M., Asaba, H. and Terasaki, T. (1999) Blood-Brain Barrier  
26 Produces Significant Efflux of L-Aspartic Acid but Not D-Aspartic Acid. *Journal of*  
27 *Neurochemistry*, **73(3)**, pp. 1206-1211.

- 1 Hull, J., Hindy, M.E., Kehoe, P.G., Chalmers, K., Love, S. and Conway, M.E. (2012)  
2 Distribution of the branched chain aminotransferase proteins in the human brain and  
3 their role in glutamate regulation. *Journal of Neurochemistry*, **123(6)**, pp. 997-1009.
- 4 Hull, J., Patel, V., El Hindy, M., Lee, C., Odeleye, E., Hezwani, M., Love, S., Kehoe,  
5 P., Chalmers, K. and Conway, M. (2015a) Regional increase in the expression of the  
6 BCAT proteins in Alzheimer's disease brain: Implications in glutamate toxicity. *Journal*  
7 *of Alzheimer's Disease*, **45(3)**, pp. 891-905.
- 8 Hull, J., Patel, V.B., Hutson, S.M. and Conway, M.E. (2015b) New insights into the  
9 role of the branched-chain aminotransferase proteins in the human brain. *Journal of*  
10 *Neuroscience Research*, **93(7)**, pp. 987-998.
- 11 Hutson, S.M., Sweatt, A.J. and LaNoue, K.F. (2005) Branched-chain amino acid  
12 metabolism: implications for establishing safe intakes. *The Journal of Nutrition*,  
13 **135(6)**, pp. 1557S-1564S.
- 14 Islam, M.M., Wallin, R., Wynn, R.M., Conway, M., Fujii, H., Mobley, J.A., Chuang, D.T.  
15 and Hutson, S.M. (2007) A novel branched-chain amino acid metabolon: protein-  
16 protein interactions in a supramolecular complex. *Journal of Biological Chemistry*,  
17 **282(16)**, pp. 11893-11903.
- 18 Islam, M.M., Nautiyal, M., Wynn, R.M., Mobley, J.A., Chuang, D.T. and Hutson, S.M.  
19 (2010) Branched-chain amino acid metabolon: interaction of glutamate  
20 dehydrogenase with the mitochondrial branched-chain aminotransferase (BCATm).  
21 *Journal of Biological Chemistry*, **285(1)**, pp. 265-276.
- 22 Kanamori, K., Ross, B.D. and Kondrat, R.W. (1998) Rate of glutamate synthesis from  
23 leucine in rat brain measured *in vivo* by <sup>15</sup>N NMR. *Journal of Neurochemistry*, **70(3)**,  
24 pp. 1304-1315.
- 25 Keen, R.E., Nissenson, C.H. and Barrio, J.R. (1993) Analysis of femtomole  
26 concentrations of  $\alpha$ -ketoisocaproic acid in brain tissue by precolumn fluorescence  
27 derivatization with 4, 5-dimethoxy-1, 2-diaminobenzene. *Analytical Biochemistry*,  
28 **213(1)**, pp. 23-28.

- 1 Kosaka, T. and Hama, K. (1986) Three-dimensional structure of astrocytes in the rat  
2 dentate gyrus. *Journal of Comparative Neurology*, **249(2)**, pp. 242-260.
- 3 Lai, J.C., Sheu, K.F.R., Kim, Y.T., Clarke, D.D. and Blass, J.P. (1986) The subcellular  
4 localization of glutamate dehydrogenase (GDH): is GDH a marker for mitochondria in  
5 brain? *Neurochemical research*, **11(5)**, pp. 733-744.
- 6 LaNoue, K.F., Berkich, D.A., Conway, M.E., Barber, A.J., Hu, L., Taylor, C. and  
7 Hutson, S. (2001) Role of specific aminotransferases in *de novo* glutamate synthesis  
8 and redox shuttling in the retina. *Journal of Neuroscience Research*, **66**, pp. 914-922.
- 9 Lee, W.K., Shin, S., Cho, S.S. and Park, J.S. (2000) Purification and characterization  
10 of glutamate dehydrogenase as another isoprotein binding to the membrane of rough  
11 endoplasmic reticulum. *Journal of Cellular Biochemistry*, **76(2)**, pp. 244-253.
- 12 Li, M., Li, C., Allen, A., Stanley, C.A., and Smith, T.J. (2012) The structure and  
13 allosteric regulation of mammalian glutamate dehydrogenase. *Archives of*  
14 *Biochemistry and Biophysics*, **519(2)**, pp. 69-80.
- 15 Lieth, E., LaNoue, K.F., Berkich, D.A., Xu, B., Ratz, M., Taylor, C. and Hutson, S.M.  
16 (2001) Nitrogen shuttling between neurons and glial cells during glutamate synthesis.  
17 *Journal of Neurochemistry*, **76(6)**, pp. 1712-1723.
- 18 Lorin, S., Tol, M.J., Bauvy, C., Strijland, A., Poüs, C., Verhoeven, A.J., Codogno, P.  
19 and Meijer, A.J. (2013) Glutamate dehydrogenase contributes to leucine sensing in  
20 the regulation of autophagy. *Autophagy*, **9(6)**, pp. 850-860.
- 21 Mastorodemos, V., Zaganas, I., Spanaki, C., Bessa, M. and Plaitakis, A. (2005)  
22 Molecular basis of human glutamate dehydrogenase regulation under changing  
23 energy demands. *Journal of neuroscience research*, **79(1-2)**, pp. 65-73.
- 24 Matsuo, Y., Yagi, M. and Walser, M. (1993) Arteriovenous differences and tissue  
25 concentrations of branched-chain ketoacids. *Journal of Laboratory and Clinical*  
26 *Medicine*, **121**, pp. 779-784.

- 1 Matthews, D.E., Bier, D.M., Rennie, M.J., Edwards, R.H., Halliday, D., Millward, D.J.  
2 and Clugston, G.A. (1981) Regulation of leucine metabolism in man: a stable isotope  
3 study. *Science*, **214(4525)**, pp. 1129-1131.
- 4 McKenna, M.C., Stridh, M.H., McNair, L.F., Sonnewald, U., Waagepetersen, H.S. and  
5 Schousboe, A. (2016) Glutamate oxidation in astrocytes: roles of glutamate  
6 dehydrogenase and aminotransferases. *Journal of Neuroscience Research*, **94(12)**,  
7 pp. 1561-1571.
- 8 Nissen, J.D., Lykke, K., Bryk, J., Stridh, M.H., Zaganas, I., Skytt, D.M., Schousboe, A.,  
9 Bak, L.K., Enard, W., Pääbo, S. and Waagepetersen, H.S. (2016) Expression of the  
10 human isoform of glutamate dehydrogenase, hGDH2, augments TCA cycle capacity  
11 and oxidative metabolism of glutamate during glucose deprivation in astrocytes. *Glia*,  
12 **65(3)**, pp. 474-488.
- 13 Nissen, J.D., Pajęcka, K., Stridh, M.H., Skytt, D.M. and Waagepetersen, H.S. (2015)  
14 Dysfunctional TCA-cycle metabolism in glutamate dehydrogenase deficient  
15 astrocytes. *Glia*, **63(12)**, pp. 2313-2326.
- 16 Nissen, S., and Haymond, M.W. (1981) Effects of fasting on flux and interconversion  
17 of leucine and alpha-ketoisocaproate *in vivo*. *American Journal of Physiology-  
18 Endocrinology and Metabolism*, **241(1)**, pp. E72-E75.
- 19 Plaitakis, A., Zaganas, I. and Spanaki, C. (2013) Deregulation of glutamate  
20 dehydrogenase in human neurologic disorders. *Journal of Neuroscience Research*,  
21 **91(8)**, pp. 1007-1017.
- 22 Popov, K.M., Shimomura, Y. and Harris, R.A. (1991) Purification and comparative  
23 study of the kinases specific for branched chain  $\alpha$ -ketoacid dehydrogenase and  
24 pyruvate dehydrogenase. *Protein Expression and Purification*, **2(4)**, pp. 278-286.
- 25 Popov, K.M., Zhao, Y., Shimomura, Y., Kuntz, M.J. and Harris, R.A. (1992) Branched-  
26 chain alpha-ketoacid dehydrogenase kinase. Molecular cloning, expression, and  
27 sequence similarity with histidine protein kinases. *Journal of Biological Chemistry*,  
28 **267(19)**, pp. 13127-13130.

- 1 Purohit, J.S., Tomar, R.S., Panigrahi, A.K., Pandey, S.M., Singh, D. and Chaturvedi,  
2 M.M. (2013) Chicken liver glutamate dehydrogenase (GDH) demonstrates a histone  
3 H3 specific protease (H3ase) activity *in vitro*. *Biochimie*, **95(11)**, pp. 1999-2009.
- 4 Rajas, F., Gire, V. and Rousset, B. (1996) Involvement of a membrane-bound form of  
5 glutamate dehydrogenase in the association of lysosomes to microtubules. *Journal of*  
6 *Biological Chemistry*, **271(47)**, pp. 29882-29890.
- 7 Riedmaier, A.E., Nies, A.T., Schaeffeler, E. and Schwab, M., (2012) Organic anion  
8 transporters and their implications in pharmacotherapy. *Pharmacological Reviews*,  
9 **64(3)**, pp. 421-449.
- 10 Rothe, F., Brosz, M. and Storm-Mathisen, J. (1994) Quantitative ultrastructural  
11 localization of glutamate dehydrogenase in the rat cerebellar cortex. *Neuroscience*,  
12 **62(4)**, pp. 1133-1146.
- 13 Shimomura, Y., Nanaumi, N., Suzuki, M., Popov, K.M. and Harris, R.A. (1990)  
14 Purification and partial characterization of branched-chain  $\alpha$ -ketoacid dehydrogenase  
15 kinase from rat liver and rat heart. *Archives of Biochemistry and Biophysics*, **283(2)**,  
16 pp. 293-299.
- 17 Shimomura, Y., Obayashi, M., Murakami, T. and Harris, R.A. (2001) Regulation of  
18 branched-chain amino acid catabolism: nutritional and hormonal regulation of activity  
19 and expression of the branched-chain  $\alpha$ -keto acid dehydrogenase kinase. *Current*  
20 *Opinion in Clinical Nutrition & Metabolic Care*, **4(5)**, pp. 419-423.
- 21 Smith, T.J. and Stanley, C.A. (2008) Untangling the glutamate dehydrogenase  
22 allosteric nightmare. *Trends in Biochemical Sciences*, **33(11)**, pp. 557-564.
- 23 Spanaki, C., Zaganas, I., Kleopa, K.A. and Plaitakis, A. (2010) Human GLUD2  
24 glutamate dehydrogenase is expressed in neural and testicular supporting cells.  
25 *Journal of Biological Chemistry*, **285(22)**, pp. 16748-16756.
- 26 Spanaki, C. and Plaitakis, A. (2012) The role of glutamate dehydrogenase in  
27 mammalian ammonia metabolism. *Neurotoxicity Research*, **21(1)**, pp. 117-127.



- 1 Spanaki, C., Kotzamani, D., Petraki, Z., Drakos, E. and Plaitakis, A. (2014)  
2 Heterogeneous cellular distribution of glutamate dehydrogenase in brain and in non-  
3 neural tissues. *Neurochemical Research*, **39(3)**, pp. 500-515.
- 4 Staten, M.A., Bier, D.M. and Matthews, D.E. (1984). Regulation of valine metabolism  
5 in man: a stable isotope study. *The American Journal of Clinical Nutrition*, **40(6)**, pp.  
6 1224-1234.
- 7 Subbalakshmi, G.Y.C.V. and Murthy, C.R. (1985) Differential response of enzymes of  
8 glutamate metabolism in neuronal perikarya and synaptosomes in acute  
9 hyperammonemia in rat. *Neuroscience Letters*, **59(1)**, pp. 121-126.
- 10 Suryawan, A., Hawes, J.W., Harris, R.A., Shimomura, Y., Jenkins, A.E. and Hutson,  
11 S.M. (1998) A molecular model of human branched-chain amino acid metabolism.  
12 *The American Journal of Clinical Nutrition*, **68(1)**, pp. 72-81.
- 13 Teichberg, V.I., Cohen-Kashi-Malina, K., Cooper, I. and Zlotnik, A. (2009)  
14 Homeostasis of glutamate in brain fluids: an accelerated brain-to-blood efflux of  
15 excess glutamate is produced by blood glutamate scavenging and offers protection  
16 from neuropathologies. *Neuroscience*, **158(1)**, pp. 301-308.
- 17 Tomita, T., Kuzuyama, T. and Nishiyama, M. (2011) Structural basis for leucine-  
18 induced allosteric activation of glutamate dehydrogenase. *Journal of Biological*  
19 *Chemistry*, **286(43)**, pp. 37406-37413.
- 20 Uchida, Y., Ohtsuki, S., Katsukura, Y., Ikeda, C., Suzuki, T., Kamiie, J. and Terasaki,  
21 T. (2011) Quantitative targeted absolute proteomics of human blood–brain barrier  
22 transporters and receptors. *Journal of Neurochemistry*, **117(2)**, pp. 333-345.
- 23 Wynn, R.M., Ho, R., Chuang, J.L., AEvarsson, A., Hol, W.G.J. and Chuang, D.T.  
24 (2000) Structure and function of human mitochondrial branched-chain  $\alpha$ -ketoacid  
25 dehydrogenase. *Biochemical Society Transactions*, **28(5)**, pp. A424-A424.
- 26 Xu, G., Kwon, G., Cruz, W.S., Marshall, C.A. and McDaniel, M.L. (2001) Metabolic  
27 regulation by leucine of translation initiation through the mTOR-signaling pathway by  
28 pancreatic  $\beta$ -cells. *Diabetes*, **50(2)**, pp. 353-360.

- 1 Yudkoff, M., Daikhin, Y., LIN, Z., Nissim, L., Stern, J., Pleasure, D. and Nissim, I.  
2 (1994) Interrelationships of leucine and glutamate metabolism in cultured astrocytes.  
3 *Journal of Neurochemistry*, **62(3)**, pp. 1192-1202.
- 4 Yudkoff, M., Nissim, I., Kim, S., Pleasure, D., Hummeler, K. and Segal, S. (1983) [<sup>15</sup>N]  
5 leucine as a source of [<sup>15</sup>N] glutamate in organotypic cerebellar explants. *Biochemical*  
6 *and Biophysical Research Communications*, **115(1)**, pp. 174-179.
- 7 Zaganas, I., Waagepetersen, H.S., Georgopoulos, P., Sonnewald, U., Plaitakis, A. and  
8 Schousboe, A. (2001) Differential expression of glutamate dehydrogenase in cultured  
9 neurons and astrocytes from mouse cerebellum and cerebral cortex. *Journal of*  
10 *Neuroscience Research*, **66(5)**, pp. 909-913.
- 11 Zaganas, I., Spanaki, C. and Plaitakis, A. (2012) Expression of human GLUD2  
12 glutamate dehydrogenase in human tissues: functional implications. *Neurochemistry*  
13 *International*, **61(4)**, pp. 455-462.
- 14 Zhou, Y., Jetton, T.L., Goshorn, S., Lynch, C.J. and She, P. (2010) Transamination is  
15 required for  $\alpha$ -ketoisocaproate but not leucine to stimulate insulin secretion. *Journal*  
16 *of Biological Chemistry*, **285(44)**, pp. 33718-33726.
- 17 Zlotnik, A., Gurevich, B., Cherniavsky, E., Tkachov, S., Matuzani-Ruban, A., Leon, A.,  
18 Shapira, Y. and Teichberg, V.I. (2008) The contribution of the blood glutamate  
19 scavenging activity of pyruvate to its neuroprotective properties in a rat model of  
20 closed head injury. *Neurochemical Research*, **33(6)**, pp. 1044-1050.
- 21 Zlotnik, A., Sinelnikov, I., Gruenbaum, B.F., Gruenbaum, S.E., Dubilet, M., Dubilet, E.,  
22 Leibowitz, A., Ohayon, S., Regev, A., Boyko, M. and Shapira, Y. (2012) Effect of  
23 glutamate and blood glutamate scavengers oxaloacetate and pyruvate on neurological  
24 outcome and pathohistology of the hippocampus after traumatic brain injury in rats.  
25 *The Journal of the American Society of Anesthesiologists*, **116(1)**, pp. 73-83.

1 **Figure legends**

2

3 **Figure 1. Specificity of the GDH and BCKD-E1 $\alpha$  antibodies.** Western blot analysis  
4 of pure protein (50 ng) and human brain homogenate (10  $\mu$ g). A: Western blot analysis  
5 using anti-GDH. B: Western blot analysis using anti-BCKD (E1 $\alpha$  subunit). C:  
6 Immunohistochemistry BCKD staining of the cerebellum. D: Antigen incubation of  
7 serial section of panel C, at 200X molar excess showing complete removal of  
8 immunoreactivity. E: Immunohistochemistry GDH staining of the cerebellum. F:  
9 Antigen incubation of serial section of panel E, at 200X molar excess showing  
10 complete removal of immunoreactivity. Scale: C, D, E and F, 25  $\mu$ m, 40x magnification.

11

12 **Figure 2. Co-localisation of BCKD-E1 $\alpha$  to cell type within the cerebellum.** A: (From  
13 left to right). Green immunofluorescence labelling BCKD-E1 $\alpha$ , red immunofluorescence  
14 labelling  $\beta$ 3 tubulin (a marker of neuronal cells) and finally a merge showing BCKD-E1 $\alpha$   
15 is present in neurons. B: (From left to right). Green immunofluorescence labelling  
16 BCKD-E1 $\alpha$ , red immunofluorescence labelling VWF (a marker of vessels) and finally a  
17 merge showing BCKD-E1 $\alpha$  is present in vessels. Scale bar: A and B, 25  $\mu$ m, 40x  
18 magnification.

19

20 **Figure 3. Staining of BCKD-E1 $\alpha$  in the frontal and temporal lobe.** A: The  
21 hippocampal formation showing immunopositive neurons within the cornu ammonis  
22 area 4 (CA4) and CA3 regions. B: Hippocampal region CA4 showing positive neuronal  
23 cells (small arrows) and a positive vessel (large arrow). C: Hippocampal region CA2  
24 showing positive neuronal cells. D: Small immunoreactive neurons (small arrows) and  
25 large pyramidal neurons (large arrows) within the temporal cortex. E: Small  
26 immunoreactive neurons and a small vessel within the temporal cortex. F: Ependymal  
27 tissue showing immunopositive cells within the frontal cortex. Scale bar: A, 200  $\mu$ m,  
28 4x magnification; B, C, E and F, 25  $\mu$ m, 40x magnification; D, 100  $\mu$ m, 10x  
29 magnification.

1

2 **Figure 4. Staining of BCKD-E1 $\alpha$  in the cerebellum.** A: The cerebellum structure  
3 showing immunopositive neurons within the molecular layer. B-C: Immunopositive  
4 Purkinje neurons within the molecular layer of the cerebellum with weak labelling of  
5 the granule cell layer. D-F: The white matter of the cerebellum showing  
6 immunoreactive capillaries. Scale bar: A, 200  $\mu$ m, 4x magnification; B, C, E and F, 25  
7  $\mu$ m, 40x magnification; D, 100  $\mu$ m, 10x magnification.

8

9 **Figure 5. Staining of BCKD-E1 $\alpha$  in the cerebellum and brainstem.** A: The dentate  
10 nucleus within the cerebellum. B-C: Increased magnification of the dentate nucleus  
11 showing staining of small neurons. D: The Raphe nuclei (large arrows) showing  
12 immunopositive neurons adjacent to the 4<sup>th</sup> ventricle (\*) and medial longitudinal  
13 fasciculus (MLF). E: Immunopositive neurons within the nucleus basalis of Meynert.  
14 F: The locus coeruleus showing immunopositive neurons. Scale bar: A and D, 200  
15  $\mu$ m, 4x magnification; B, C, E and F, 25  $\mu$ m, 40x magnification.

16

17 **Figure 6. Co-localisation of GDH to cell type within the cerebellum.** A: (From left  
18 to right). Green immunofluorescence labelling GDH, red immunofluorescence labelling  
19 GFAP (a marker of astrocytes) and finally a merge showing GDH is detected in  
20 astrocytes (small arrows). B: (From left to right). Green immunofluorescence labelling  
21 GDH, red immunofluorescence labelling GFAP (a marker of astrocytes) and finally a  
22 merge showing GDH is present in astrocytes surrounding vessels (small arrows). C:  
23 (From left to right). Green immunofluorescence labelling GDH, red  
24 immunofluorescence labelling VWF (a marker of vessels) and finally a merge showing  
25 GDH is present within certain vessels. Scale bar: A, B and C, 25  $\mu$ m, 4x magnification.

26

27 **Figure 7. Staining of GDH in the frontal and temporal lobe.** A: The hippocampal  
28 formation showing supportive immunopositive astrocytes within the molecular layer.  
29 B: Increased magnification of the granule cell layer of the dentate gyrus (DG) showing

1 small immunopositive astrocytes. C: Increased magnification of astrocyte staining  
2 showing both cell body and process labelling. D: Immunoreactive astrocytes within  
3 the temporal white matter. E: Increased magnification of immunoreactive astrocytes.  
4 F: Ependymal and subependymal tissue showing immunopositive cells within the  
5 frontal cortex. Scale bar: A, 200  $\mu\text{m}$ , 4x magnification; B, C, E and F, 25  $\mu\text{m}$ , 40x  
6 magnification; D, 50  $\mu\text{m}$ , 40x magnification.

7

8 **Figure 8. Staining of GDH in the temporal lobe and cerebellum.** A: Temporal  
9 cortex with labelling of large pyramidal neurons (small arrows) adjacent to an  
10 immunopositive vessel (\*). B: Cerebellar white matter showing perivascular astrocytic  
11 labelling. C: Cerebellar white matter showing vessel staining. Scale bar: A, 100  $\mu\text{m}$ ,  
12 10x magnification; B and C, 25  $\mu\text{m}$ , 40x magnification.

13

14 **Figure 9. Staining of GDH in the cerebellum and putamen.** A: The cerebellum  
15 formation showing immunopositive cells within the molecular and nuclear layers. B-  
16 C: Granule cell layer with immunopositive purkinje cell (small arrow) and labelling  
17 within the nuclear layer (large arrow) likely representing a vessel or astrocytic  
18 processes. Also present are immunopositive Bergman astrocytes at the level of the  
19 golgi cell layer (arrow head). D: Dentate nuclei immunopositive for GDH. E: Increased  
20 magnification of the dentate nuclei demonstrating negative neuronal staining with  
21 immunopositive astrocytes and astrocytic processes. F: Nuclear presence of GDH  
22 (small arrow) within the putamen. Scale bar: A and D, 200  $\mu\text{m}$ , 4x magnification; B,  
23 C, E and F, 25  $\mu\text{m}$ , 40x magnification.

24

25 **Figure 10. The glutamate-glutamine cycle and the supporting role of the hBCAT,**  
26 **BCKD and GDH proteins.** Glutamate is released from neuronal cells during  
27 excitatory neurotransmission. Post-excitation, glutamate is taken up via excitatory  
28 amino acid transporters on astrocytes. In astrocytes much of the glutamate undergoes  
29 amidation to glutamine by glutamine synthetase. Glutamine is then released for  
30 uptake by neuronal cells, to replenish the glutamate pool through the action of

1 glutaminase. However, much glutamate can be lost through oxidation via glutamate  
2 dehydrogenase, or generation of glutathione/purines in astrocytes. The glutamate-  
3 glutamine cycle must work with other anapleurotic pathways to regenerate the  
4 neuronal pool of glutamate. BCAA metabolism is thought to participate in nitrogen  
5 shuttling in the *de novo* synthesis of glutamate. The BCAAs, particularly leucine, can  
6 pass easily through the blood brain barrier through an L system transporter. Uptake  
7 by neuronal cells results in transamination by hBCATc with  $\alpha$ -ketoglutarate forming  
8 glutamate, contributing to the neuronal pool. The BCKAs are completely metabolised  
9 via BCKD to produce Kreb's cycle intermediates and avoid re-transamination of  
10 glutamate. In addition to astrocytes, the endothelial cells of the vasculature play a key  
11 role in removing excess glutamate and regenerating neuronal pools of glutamate. As  
12 levels of hBCATm are strongly expressed and GDH is also present in many vessels,  
13 we propose that glutamate is either transaminated to the BCAAs or metabolised by  
14 GDH. The BCAAs can then re-enter the pool for glutamate generation, and the  
15 ammonia ( $\text{NH}_3$ ) generated by GDH activity can be exported from the brain.  
16 Abbreviations: BCAA, branched chain amino acid; BCKD, branched chain keto acid  
17 dehydrogenase complex; GDH, glutamate dehydrogenase; Gln, glutamine; Glu,  
18 glutamate; hBCAT, human branched chain aminotransferase.

19

20 **Table 1. Clinical outcome for patients used in this study.**

21

22 **Supplementary figure 1. Specificity of the GDH and BCKD-E1 $\alpha$  antibodies.**

23 Immunohistochemical analysis of human brain sections, including antigen incubations.  
24 A: Immunohistochemistry BCKD staining of the cerebellum. B: Antigen incubation of  
25 serial section of panel A, at 200X molar excess showing complete removal of  
26 immunoreactivity. C: Immunohistochemistry GDH staining of the cerebellum. D:  
27 Antigen incubation of serial section of panel C, at 200X molar excess showing  
28 complete removal of immunoreactivity. Scale: C, D, E and F, 50  $\mu\text{m}$ , 10x magnification.

29

30

1 **Supplementary table 1. Clinical outcome for patients used in this study.**

2