

1 Holocene alluvial fan evolution, Schmidt-hammer exposure-age dating  
2 and paraglacial debris floods in the SE Jostedalbreen region, southern  
3 Norway

4  
5  
6 JOHN A. MATTHEWS, LINDSEY J. MCEWEN, GERAINT OWEN AND SIETSE  
7 LOS

8  
9  
10 Matthews, J.A., McEwen, L.J., Owen, G. & Los, S. 2020. Holocene alluvial fan  
11 evolution, Schmidt-hammer exposure-age dating and paraglacial debris floods in the  
12 SE Jostedalbreen region, southern Norway. *Boreas*, Vol.

13  
14 The evolution of several subalpine alluvial fans SE of the Jostedalbreen ice cap was  
15 investigated based on their geomorphology and Schmidt-hammer exposure-age dating  
16 (SHD) applied to 47 boulder deposits on the fan surfaces. A debris-flood rather than  
17 debris-flow or water-flow origin for the deposits was inferred from their morphology,  
18 consisting of low ridges with terminal splays up to 100 m wide without lateral levees.  
19 This was supported by fan, catchment, and boulder characteristics. SHD ages ranged  
20 from  $9480 \pm 765$  to  $1955 \pm 810$  years. The greatest number of boulder deposits, peak  
21 debris-flood activity and maximum fan aggradation occurred between  $\sim 9.0$  and  $8.0$   
22 ka, following regional deglaciation at  $\sim 9.7$  ka. The high debris concentrations  
23 necessary for debris floods were attributed to paraglacial processes enhanced by  
24 unstable till deposits on steep slopes within the catchments. From  $\sim 8.0$  ka, fan  
25 aggradation became progressively less as the catchment sediment sources tended  
26 towards exhaustion, precipitation decreased during the Holocene Thermal Maximum,  
27 and tree cover increased. After  $\sim 4.0$  ka, some areas of fan surfaces stabilized, while  
28 Late-Holocene climatic deterioration led to renewed fan aggradation in response to  
29 the neoglacial growth of glaciers, culminating in the Little Ice Age. These changes are  
30 generalized within a conceptual model of alluvial fan evolution in this recently-  
31 deglaciated mountain region and in glacierized catchments. This study highlights the  
32 potential importance of debris floods, of which relatively little is known, especially in  
33 the context of alluvial fan evolution.

34  
35 *John A. Matthews ([J.A.Matthews@Swansea.ac.uk](mailto:J.A.Matthews@Swansea.ac.uk)), Geraint Owen, Sietse Los,*  
36 *Department of Geography, College of Science, Swansea University, Singleton Park,*  
37 *Swansea SA2 8PP, Wales, UK; Lindsey J. McEwen, Centre for Water Communities*  
38 *and Resilience, University of the West of England, Coldharbour Lane, Bristol BS16*  
39 *1QY, UK; received*

40  
41  
42 Alluvial fans are fan-shaped depositional landforms created where steep, high-  
43 powered channelized flows deposit their material load on entering a zone of flow  
44 expansion and reduced power (Harvey 2004; Owen 2014; Ventra & Clarke 2018).  
45 Typically, they are classified according to the predominant depositional process, into  
46 (i) fluvial fans, where stream flows (water flow or water flood) deposit bedload, and  
47 (ii) colluvial fans dominated by mass movement processes, particularly debris flow

48 (also known as gravity-flow fans) (Rachocki & Church 1990; Crosta & Frattini 2004;  
49 Harvey *et al.* 2005; De Haas *et al.* 2015, 2019; Bowman 2019). Whereas most studies  
50 have emphasised these two types of alluvial fans, there is increasing recognition of the  
51 existence of a continuum of landforms, which reflect interactions between processes  
52 and intermediate-type flows (Wells & Harvey 1987; Hungr *et al.* 2001; Germain &  
53 Ouellet 2014). Terms for the flows that are intermediate in character between water  
54 floods and debris flows include fluid (wet or watery) debris flows (Sletten & Blikra  
55 2007; Harvey *et al.* 2013), hyperconcentrated flows (Matthews *et al.* 1999; Pierson  
56 2005; Sletten & Blikra 2007; Calhoun & Clague 2018), debris torrents (Slaymaker  
57 1988) and debris floods (Hungr *et al.* 2001; Wilford *et al.* 2004; Mayer *et al.* 2010;  
58 D'Agostino 2013; Ouellet & Germain 2014). However, the nature of these flows,  
59 which are characterised by sediment concentrations of 40–70% by weight according  
60 to Costa (1984), and their role in fan development, are still poorly understood.

61

62 In order to understand better the development of alluvial fans, the long-  
63 standing problem of precise numerical dating (of the fan surface) needs to be  
64 overcome. Several techniques ranging from historical analysis to dendrochronology  
65 and lichenometry have been applied to the dating of fan development over annual to  
66 decadal timescales (e.g. D'Agostino 2013; Jomelli 2013; Schneuwly-Bollschweiler &  
67 Stoffel 2013; Stoffel 2013). Far fewer techniques, including those based on  
68 radiocarbon, optically stimulated luminescence (OSL) and terrestrial cosmogenic  
69 nuclides are applicable over longer, centennial to millennial timescales (e.g. Harvey *et*  
70 *al.* 2005; Schneuwly-Bollschweiler *et al.* 2013; Schürch *et al.* 2016). Here we apply  
71 the relatively new technique of Schmidt-hammer exposure-age dating (SHD) to fan  
72 surfaces. SHD is appropriate for providing numerical ages for boulders exposed  
73 during the Lateglacial and Holocene (see, for example, Winkler 2009; Matthews &  
74 Owen 2010; Shakesby *et al.* 2011; Matthews *et al.* 2013, 2015, 2018; Stahl *et al.*  
75 2013; Tomkins *et al.* 2016, 2018; Wilson & Matthews 2016; Winkler *et al.* 2016;  
76 Wilson *et al.* 2019).

77

78 Various temporal patterns and activity phases have been recognised in records  
79 of floods, debris flows and other colluvial processes (ranging from snow flows to rock  
80 falls) in southern Norway (Blikra & Nesje 1997; Blikra & Nemeč 1998; Blikra &  
81 Selvik 1998; Sletten *et al.* 2003; Bøe *et al.* 2006; Sletten & Blikra 2007; Matthews *et*

82 *al.* 2009, 2018; Vasskog *et al.* 2011). Detailed case studies of two alluvial fans have,  
83 moreover, revealed contrasting histories. Radiocarbon dating and lichenometry show  
84 that development of the subalpine Nystølen fan in the Jostedalsgreen region (Lewis &  
85 Birnie 2001; McEwen *et al.* 2011) was dominated by deposition in the Little Ice Age  
86 of the last few centuries, whereas SHD shows that the alpine Illåe fan in Jotunheimen  
87 is largely a relict paraglacial landform that developed before ~8.0 ka (McEwen *et al.*  
88 2020). Differences in the evolution of these two fans were accounted for largely by  
89 the extent to which their catchments were glacierized in the past.

90

91 In this study, the aim is to generalize further by analysing the development of  
92 subalpine alluvial fans in the SE Jostedalsgreen region of southern Norway (Fig. 1),  
93 based on their geomorphology and the exposure age of their surface boulder deposits.  
94 There are three main objectives: (i) To date the numerous boulder deposits on the  
95 fan surfaces using SHD and hence provide a firm chronology; (ii) To assess the  
96 origin of the boulder deposits with reference to processes of debris flow, water flow  
97 (floods) and debris floods; and (iii) To reconstruct the evolution of several fans and  
98 hence develop a regional conceptual model of fan evolution in recently-deglaciated  
99 mountain catchments.

100

101

## 102 Study sites and environment

103

104 The alluvial fans are located to the SE of the Jostedalsgreen ice cap on valley floors at  
105 300–400 m above sea level at the foot of steep tributary valleys descending from a  
106 glacierized plateau at >1600 m a.s.l. (Fig. 2). Four fans, from south to north, are  
107 termed here: (i) the Erikstølsdalen fan; (ii) the Kvamsdalen fan; (iii) the Snøskreda  
108 fan; and (iv) the Kupejelet fan, the latter two being located in Austerdalen.  
109 Kupejelet fan (Fig. 3), in many ways similar to the others, was previously  
110 investigated by Innes (1985a, b). Data from a fifth fan (Nystølen fan) in lower  
111 Langedalen (Fig. 2), previously investigated by Lewis & Birnie (2001) and McEwen  
112 *et al.* (2011) are included in some of our analyses. The four fans (Fig. 4) were selected  
113 because of their extensive boulder deposits suitable for dating by SHD, using field  
114 and aerial photographic evidence. Other neighbouring fans were unsuitable: those

115 south of Veitastromd have been greatly modified by land clearance, while those further  
116 north in Austerdalen (e.g. at the mouth of Røysedalen) have been eroded by the  
117 historical advance of the glacier Austerdalsbreen.

118

119 All five fans are subalpine in character: the Kvamsdalen and Kupegelet fans  
120 are largely covered in *Betula pubescens* woodland, whereas the Erikstølen, Snøskreda  
121 and Nystølen fans have much larger areas of grassland, which are partly the result of  
122 snow-avalanche activity, and partly a response to grazing animals associated with the  
123 agricultural settlement of Veitastromd and sæters such as Tungestølen. Climatic data  
124 from the meteorological station Bjørkhaug, in the neighbouring valley of Jostedal  
125 (324 m a.s.l.), indicate a mean annual air temperature of +3.7 °C, with a July mean of  
126 +13.4 °C, a January mean of -4.9 °C and a mean annual precipitation of 1380 mm  
127 (Aune 1993; Førland 1993). The local lithology is predominantly granite with some  
128 areas of granitic gneiss (Lutro & Tveten 1996).

129

130 Morphometric data from the fans, their catchments, and their surface boulder  
131 deposits are summarised in Table 1. The five catchments are small (1.17–3.44 km<sup>2</sup>),  
132 high relief, steep and rugged, with a Melton ratio (relief/ $\sqrt{\text{area}}$ ; e.g. Melton 1965) of  
133 0.70–1.08. The fans are correspondingly small (0.16–0.51 km<sup>2</sup>) with gradients of 9–  
134 17°, but fan toes have been truncated by the main river or obscured by the growth of  
135 peat mires. The boulder deposits on the fan surfaces form broad, irregular ridges, up  
136 to 200 m in length with a mean width of 24–34 m (maximum width 100 m), most with  
137 terminal splays, some with finger-like extensions (Fig. 5).

138

139 Three of the investigated catchments are currently 8–24% glacierized by the  
140 Kvitikoll ice cap (Fig. 1) which, together with the Tverrdalsbreen glacier, occupy the  
141 plateau and extend onto the lee-slopes to the east. The catchment of the Nystølen fan  
142 is 56% glacierized. However, all catchments have late-lying snowbeds on their upper  
143 slopes, and are likely to have been affected by expanded plateau glaciers during Late-  
144 Holocene neoglaciation and especially in the Little Ice Age.

145

146 Rapid Early-Holocene deglaciation of the main valleys of SE Jostedalsbreen  
147 occurred during the Preboreal, and by ~10.1-9.7 ka glaciers had receded to the valley  
148 heads, close to their Little Ice Age limits (Dahl *et al.* 2002; see also Mottershead &

149 Collin 1976; Aa 1982; Nesje 1991, 2009). Subsequently, further rapid warming and  
150 glacier shrinkage resulted in the total melting of the Jostedalbreen ice cap by ~7.3 ka  
151 (Nesje & Kvamme 1991; Nesje *et al.* 2000, 2001). Centennial- to millennial-scale  
152 glacier variations interrupted neoglacial re-growth of glaciers after ~6.1 ka, which  
153 culminated in the Little Ice Age maximum of extant glaciers around AD 1750 (Grove  
154 1988; Bickerton & Matthews 1993). Latero-terminal moraines indicate the down-  
155 valley limits of several of these glaciers in the Little Ice Age (Fig. 2).

156

157

## 158 Methodology

159

160 Field research focused on 47 boulder deposits, which are located on Fig. 4A-D. These  
161 represent integral geomorphological units each of which can be attributed to single  
162 depositional events. They also represent the entire statistical population of boulder  
163 deposits from each fan. SHD was carried out on these deposits, supplemented by  
164 lichenometric dating and measurements of boulder roundness and boulder size.

165

### 166 *SHD*

167

168 As a basis for SHD dating, R-values were recorded from a minimum of 100 boulders  
169 on each deposit (one impact per boulder) using a mechanical N-type Schmidt hammer  
170 (Proceq 2004). Use of one impact per boulder ensured that the R-value frequency  
171 distribution approximates the boulder-age distribution (Matthews *et al.* 2014). In  
172 order to minimise variability and measurement errors, small or unstable boulders,  
173 edges, joints and cracks, and lichen-covered or wet boulder surfaces were avoided,  
174 and measurements were confined to near-horizontal surfaces and granitic lithologies  
175 (cf. Shakesby *et al.*, 2006; Matthews & Owen, 2010; Viles *et al.*, 2011). No cleaning  
176 or artificial abrading of the boulder surfaces was carried out prior to measurement as  
177 this would have removed age-related weathering effects. The Schmidt hammer was  
178 regularly tested on the manufacturer's test anvil during the fieldwork to ensure no  
179 deterioration in instrument performance following prolonged use (cf. McCarroll 1987,  
180 1994).

181

182 Calibration of R-values followed the approach developed by Matthews &  
183 Owen (2010), Matthews & Winkler (2011) and Matthews & McEwen (2013), full  
184 details of which are given in Matthews et al. (2018). The calibration equation is a  
185 linear regression of surface age ( $y$ ) on mean R-value ( $x$ ) derived from two local  
186 control points: ‘old’ and ‘young’ surfaces of known age. Use of a linear relationship  
187 has been specifically tested over the Holocene timescale (Shakesby *et al.* 2011), and is  
188 justified also by comparison over similar relatively short timescales with terrestrial  
189 cosmogenic nuclide dating both in southern Norway (Wilson *et al.* 2019) and  
190 elsewhere (e.g. Tomkins *et al.* 2016, 2018). A linear or near-linear relationship, which  
191 results from the slow rate of chemical weathering of rock surfaces, is therefore  
192 considered appropriate, particularly in alpine and subalpine environments over the last  
193 ~10 ka.

194

195 Confidence intervals (95%) for SHD age ( $C_t$ ) are based on combining the  
196 relatively small error term associated with the calibration equation ( $C_c$ ) with the larger  
197 sampling error associated with the dated surfaces ( $C_s$ ). Uncertainty associated with  $C_s$   
198 is relatively small provided: (i) very large R-value sample sizes are used for control  
199 points; and (ii) control-point ages are accurately known. Here we used 600-750 R-  
200 values for each control point and hence can justify using precise ages for the control  
201 points.

202

203 The ‘young’ control point involves R-values from 600 boulders (one impact  
204 per boulder) deposited on the Erikstølsdalen and Snøskreda fans (Fig. 2) during a  
205 flash flood following intense rainfall on 14 August 1979 (cf. Gjessing & Wold 1980;  
206 Drageset 2001). Both the geomorphological integrity and lichen sizes associated with  
207 the flood deposits leave no doubt that the surface boulders sampled are representative  
208 of a synchronous surface and that their age is very tightly constrained. The rockfall  
209 deposits used previously by Matthews & Wilson (2015) as their ‘young’ control point  
210 were deemed unsuitable for the present study due to the roughness characteristics of  
211 such colluvial boulders (cf. Matthews & McEwen 2013; Matthews *et al.*, 2015; Olsen  
212 *et al.*, 2020). In contrast, the 1979 flood deposits, being characterised by relatively  
213 smooth boulders, have similar roughness to the boulder deposits on the fans, and are  
214 therefore appropriate for a study of alluvial fans.

215

216 The 'old' control point, involving 750 R-values recorded from three glacially-  
217 scoured bedrock outcrops near Tungastølen and at the mouth of Kvamsdalen (Fig. 2),  
218 was used previously by Matthews & Wilson (2015). The precise date of ~9.7 ka used  
219 for this control point is based on the age of moraine ridges deposited by  
220 Jostedalsbreen outlet glaciers in valleys on both sides of the ice cap. Evidence for the  
221 age of these moraines comes from both radiocarbon (Nesje 1984; Dahl *et al.* 2002)  
222 and cosmogenic nuclide dating (Matthews *et al.* 2008) in Erdalen on the NW side of  
223 the ice cap, and by radiocarbon dating near Nigardsbreen in Jostedalen on the SE side  
224 (Dahl *et al.* 2002). The moraines, which are of a similar size to Little Ice Age  
225 moraines and located up to ~1 km beyond the Little Ice Age limits of Erdalsbreen and  
226 Nigardsbreen, relate to the Erdalen Event, an Early-Holocene centennial-scale glacier  
227 and climatic fluctuation that involved two glacier re-advances dated by Dahl *et al.*  
228 (2002) to ~10.1 and 9.7 ka.

229

230 Although no similar moraines dating from the Erdalen Event occur in  
231 Austerdalen or Langedalen, the glaciers in these valleys are assumed to have  
232 fluctuated broadly synchronously with other outlet glaciers of Jostedalsbreen, as has  
233 been demonstrated for the Little Ice Age interval (cf. Bickerton & Matthews 1992,  
234 1993). We attribute the absence of Erdalen Event moraines downvalley of the Little  
235 Ice Age glacier limits in Austerdalen or Langedalen to the presence of relatively large  
236 ice bodies in these valleys and correspondingly large glacier re-advances during the  
237 Erdalen Event. Combined with the occurrence of this event during an otherwise  
238 prolonged period of rapid glacier retreat, we conclude that the 'old' control surfaces in  
239 the study area were deglaciated closely following the termination of the Erdalen Event  
240 (i.e. ~9.7 ka).

241

242 Probability density function analyses were used to understand the SHD age-  
243 frequency distributions over the Holocene timescale. Separate analyses were carried  
244 out for each fan and for the combined data set. Probability density was calculated at  
245 100-year intervals using the mean and standard deviation for each fan (R Core Team  
246 2019). Calculation assumed a normal distribution of the data. Probability density  
247 functions for each of the four alluvial fans were obtained by averaging the density  
248 values of the relevant individual boulder deposits. A regional density function was

249 obtained by averaging the density functions of all 47 boulder deposits from the four  
250 fans.

251

### 252 *Supplementary measurements*

253

254 Three types of supplementary measurements were made from the boulder deposits.  
255 First, the maximum diameter (longest axis) of the 5 largest lichens of the *Rhizocarpon*  
256 subgenus from each of the 47 deposits was measured in order to perform  
257 lichenometric dating. This had been previously attempted by Innes (1985a) for a fan  
258 in Austerdalen. We used updated lichenometric dating curves from the neighbouring  
259 glacier foreland of Nigardsbreen (Bickerton & Matthews 1992, 1993). Second, in  
260 order to assess potential sediment sources, the boulder roundness distribution and a  
261 numerical index of mean boulder roundness were derived from a subsample of 25  
262 boulders from 37 of the boulder deposits using the Powers (1953) roundness chart (cf.  
263 Matthews 1987). Finally, the maximum intermediate-axis clast size ( $d$ ) from each  
264 boulder deposit was measured to allow the calculation of palaeohydrological  
265 parameters associated with the flows that deposited the sediment (Williams 1983):

266

$$267 \text{ Unit stream power } (\omega) = 0.079 d^{1.3} (10 \leq d \leq 1500 \text{ mm}) \quad (1)$$

$$268 \text{ Bed shear stress } (\tau) = 0.17 d^{1.0} (10 \leq d \leq 3300 \text{ mm}) \quad (2)$$

$$269 \text{ Mean flow velocity } (V) = 0.065 d^{0.50} (10 \leq d \leq 1500 \text{ mm}) \quad (3)$$

270

271

## 272 **Results**

273

### 274 *R-values from control surfaces and calibration equations*

275

276 Combined data for the ‘old’ and ‘young’ control points (Table 2) show excellent  
277 agreement between each pair of ‘old’ and ‘young’ control surfaces, which justifies  
278 treating each pair of surfaces sampled from different locations as replicates drawn  
279 from the same statistical population. The R-value distributions of the control points  
280 (Fig. 6A) exhibit the symmetrical, unimodal characteristics of synchronous surfaces.  
281 Furthermore, the small standard deviations ( $\sigma = 6\text{--}8$ ) relative to the standard  
282 deviations associated with the fan surfaces ( $\sigma = 8\text{--}11$ ; Table 3), together with wide



283 separation of the mean R-values, signal the potential for dating using the calibration  
284 equation shown in Fig. 6B.

285

#### 286 *R-values and SHD ages from the boulder deposits*

287

288 R-value distributions for 47 boulder deposits are generally symmetrical and unimodal,  
289 which is again indicative of synchronous surfaces produced here by single  
290 depositional events (Fig. 6). Whereas mean R-values (Table 3) vary widely between  
291 38.5 (Sa 4) and 58.8 (En 7) most are closer to the characteristic of the ‘old’ control  
292 point than to those of the ‘young’ control point. SHD ages are correspondingly wide  
293 ranging but with a large majority of the boulder deposits dating from early in the  
294 Holocene (>70% before ~6.0 ka) and only two dating from the last 2.0 ka (Table 3).  
295 The sampling error (Cs) resulting from the high natural variability of weathered  
296 boulder surfaces is the dominant control on the 95% confidence intervals for age,  
297 which range from ~700-900 years.

298

299 SHD ages and probability density distributions indicate significant clustering  
300 of events and notable similarities and differences between the chronology of boulder  
301 deposits from each fan (Fig. 8), which are discussed below. Amalgamation of the age  
302 data from the four fans in the combined record emphasises the overall concentration  
303 of dates shortly after deglaciation and the long-term declining frequency of  
304 depositional events through the Holocene.

305

#### 306 *Lichen sizes and lichenometric ages*

307

308 Mean lichen size on the boulder deposits varies from ~50–300 mm, which  
309 corresponds to a lichenometric age of ~70–1500 years (Fig. 9). At three sites from  
310 Kupegelet and one from Snøskreda, single largest lichens reached 300–320 mm,  
311 which are comparable to the largest lichens (270–290 mm) measured from the same  
312 sites by Innes (1985). As a result of using the up-dated calibration equation of  
313 Bickerton & Matthews (1992, 1993), our results suggest that >50% of deposits are  
314 characterised by mean lichen sizes >150 mm and date from pre-Little Ice Age times.  
315 In contrast, Innes (1985) concluded that all the deposits fell within the Little Ice Age.  
316 However, as there is no correlation between our SHD and lichenometric ages, it can

317 be deduced that most of the latter are large underestimates of the true age of the  
318 deposits, resulting from extrapolation of surface ages beyond the range of  
319 lichenometric dating, combined with the limited lifespan of lichens of the  
320 *Rhizocarpon* subgenus in this environment (cf. Matthews & Trenbirth 2011).

321

322

### 323 *Boulder characteristics and palaeohydrological parameters*

324

325 Mean boulder roundness from boulder deposits across all four fans lies consistently  
326 between values characteristic of sub-angular and sub-rounded clasts (Fig. 10).  
327 Furthermore, the majority of sites (70%) have a sub-rounded modal class with a  
328 negligible proportion of angular and very angular clasts. These roundness values are  
329 consistent with a till source for the boulders and are consistent with a relatively small  
330 degree of abrasion during transportation in relatively small catchments (cf. Boulton  
331 1978; Matthews 1987; Evans & Benn 2004).

332

333 Maximum boulder size and the median size ( $D_{50}$ ; intermediate axis) of the 10  
334 largest boulders from the boulder deposits (Fig. 11) approximate to 2.0 m and 1.0 m,  
335 respectively. The largest boulder (2.5 m) occurred at Erikstølsdalen (En 1) while the  
336 largest  $D_{50}$  (1.7 m) was recorded from Kupegelet (Kt 8). Such sizes require high  
337 competent flows and imply major floods as large as the 1979 flash-flood event, the  
338 deposits of which, on the Snøskreda fan, involved maximum boulder sizes and  $D_{50}$   
339 values of 2.1 m and 1.2 m, respectively. The flash flood had a return period of ~1 in  
340 100 years, estimated from its magnitude on the main river in Jostedal (Gjessing &  
341 Wold 1980; Drageset 2001), but this may have increased to ~1 in 1000 years in  
342 smaller catchments within the region (Matthews & McEwen 2013).

343

344 Minimum boulder-transport conditions for the largest boulder in deposits for  
345 each fan are summarised in Table 4. Lowest unit stream power for entrainment ( $\omega$ )  
346 varies from 1646 W m<sup>-2</sup> (Kvamsdalen) to 2065 W m<sup>-2</sup> (Erikstølsdalen), lower than  
347 values for the coarsest debris-flood deposits on the upper Illåe fan, Jotunheimen (up to  
348 2850 W m<sup>-2</sup>; McEwen *et al.* 2020). A large number of deposits (47%) have largest  
349 clasts beyond the upper range of clast sizes used by Williams (1983). Lowest bed  
350 shear stresses for entrainment ( $\tau$ ) of the largest clast ranged from 357 to 425 N m<sup>-2</sup>.

351

352

## 353 Discussion

354

355 *Water floods, debris flows or debris floods?*

356

357 The morphology and sedimentary characteristics of the boulder deposits, and the  
358 characteristics of the fans and their catchments, tend to be intermediate in terms of  
359 established criteria for recognising the products of water flow and debris flow (Table  
360 5). Using all these criteria, the boulder deposits can be attributed with some  
361 confidence to debris floods, which are now recognised in the most widely-used  
362 genetic classification of landslide types (Hungr *et al.*, 2014).

363

364 Previous studies by Innes (1985a, 1985b) assumed that these boulder deposits  
365 were debris-flow lobes, which also tend to have boulder concentrations in their  
366 terminal areas and on lateral levées. However, their morphologies differ from debris-  
367 flow lobes in several respects. They are commonly irregular, broad ridges, which are  
368 raised above the general level of the adjacent fan surfaces by up to several metres  
369 (Fig. 5A, B). With a mean width of 24-34 m and a maximum width of up to 100 m  
370 (Table 1), they are considerably wider than typical debris-flow deposits and, crucially,  
371 levées are absent. They terminate in several different plan shapes ranging from  
372 simple, steep-fronted tongues (similar to debris-flow lobes) to single or multiple  
373 splays (less thick as well as wider than debris-flow lobes), the latter sometimes with  
374 finger-like extensions (Fig. 5C-F). Similar broad ridges without levées occur in  
375 Iceland, where they were described as ‘debris flow-like’ (Decaulne *et al.* 2007).  
376 Debris flood ridges and splays also differ from the thin gravel sheets with bars and  
377 braided channels deposited on fans by water floods.

378

379 Neither the size nor slope of each of our fans, nor their catchment  
380 characteristics, are typical of either debris-flow fans, which are smaller and steeper  
381 with very small rugged catchments, or fluvial fans, which are generally larger with  
382 gentler slopes and much larger catchments. Although no sections were available  
383 through these deposits, the surface sediments of the ridges appear intermediate

384 between unsorted diamictons and well-sorted fluvial deposits. The sediments also  
385 seem to correspond to the proximal facies of terminoglacial fans described by  
386 Zieliński & van Loon (2000), which include boulder-rich diamictons and sandy  
387 gravels deposited by sheetflows and catastrophic hyperconcentrated flows. There is  
388 little evidence of fine matrix where boulder concentrations occur, but this could have  
389 been washed out of the surface material during or after deposition. The stratigraphy of  
390 the Illåe fan (Jotunheimen), where similar boulder deposits occur, revealed a variable  
391 content of matrix with alternating, crudely-sorted and generally indistinct clast-  
392 supported and matrix-supported layers (McEwen *et al.* 2020).

393

394 In order to achieve the high debris concentrations necessary for debris floods,  
395 with sufficient large subrounded to subangular boulders (Figs 10, 11), the catchment  
396 would have had to contain a suitable sediment source. This is likely have been a till  
397 cover, deposited prior to ~9.7 ka, when the catchment was completely glacierized. We  
398 argue below that all four catchments had a substantial till cover, which was exposed to  
399 subaerial processes following deglaciation. This till cover would have been readily  
400 eroded from the steep slopes of the catchments, and provides the likely source of the  
401 sediments in the debris-flood boulder deposits.

402

#### 403 *The chronology of events*

404

405 The chronology of boulder deposits from each fan (Fig. 8) shows that the earliest  
406 depositional events occurred shortly after deglaciation at ~9.7 ka. Indeed, the oldest  
407 SHD dates from Snøskreda (Sa 4) and Erikstølsdalen (En 5) are  $9480 \pm 765$  and  
408  $9215 \pm 720$  years, respectively which are statistically indistinguishable (along with  
409 several other SHD dates) from 9.7 ka. Taking account of the confidence intervals,  
410 both of these fans have a very high proportion of SHD ages  $>8.0$  ka, while all the  
411 boulder deposits on the Kvamsdalen fan have SHD ages  $>4.0$  ka and the Kupegjelet  
412 fan has a relatively high proportion between 8.0 and 6.0 ka. Three fans developed  
413 rapidly within two millennia of deglaciation while the fourth (Kupegjelet) appears to  
414 have started its rapid development about two millennia later than the others. All four  
415 fans therefore underwent major aggradation attributable to debris-flood activity during  
416 the Early Holocene.

417

418 Persistence of so many boulder deposits from the Early Holocene is indicative  
419 of the decline in the frequency of debris-flood events later in the Holocene. If the  
420 frequency of such events had remained high during the Middle and Late Holocene,  
421 fan aggradation in the form of debris-flood deposits would have continued into the  
422 Late Holocene and earlier deposits would have been buried by later ones. Instead, a  
423 small number of debris-flood deposits with ages <4 ka occur only at the  
424 Erikstølsdalen and Kupegjelet fans. At the Kvamsdalen and Snøskreda fans, debris-  
425 flood deposits are confined to distal and marginal parts of the fans. In proximal- and  
426 mid-fan locations, however, these two fans exhibit evidence of late-Holocene and  
427 modern aggradation from snow-avalanche and fluvial activity, which may have buried  
428 earlier boulder deposits.

429

430 The combined chronology from the four fans (Fig. 8) suggests a relatively  
431 steady decline in frequency of debris-flood events from a maximum at ~9.0–8.0 ka.  
432 However, the interpretation is complicated by wide confidence intervals for SHD age  
433 of the order of 700–900 years, the apparent absence of events between deglaciation  
434 and ~8.0 ka at the Kupegjelet fan where activity peaks at ~7.0 ka, and the possibility  
435 of centennial- to millennial-scale variations in aggradation in the Middle to Late  
436 Holocene.

437

#### 438 *Holocene development of the alluvial fans and their environmental controls*

439

440 The peak in debris-flood activity immediately following deglaciation at ~9.7 ka is  
441 clearly indicative of a paraglacial pattern of sediment deposition conditioned directly  
442 by glaciation (cf. Ryder 1971; Church & Ryder 1972; Ballantyne 2002a, 2013).  
443 During and immediately following deglaciation, the till deposits on the extremely  
444 steep slopes of these catchments would have been particularly susceptible to gully  
445 erosion triggered by rainstorms, and glacial and snow meltwater (Curry 2000). Being  
446 a diamicton, the till would have been a source of abundant large subangular to  
447 subrounded boulders and fine matrix, providing the high sediment concentrations for  
448 debris-floods. These flows would have been confined in the narrow, steep tributary  
449 valleys before they debouched onto the main valley floor where redeposition and fan  
450 aggradation occurred.

451

452           Although paraglacial processes are most effective in the unstable landscape  
453 that emerges during deglaciation, paraglacial effects may last for several millennia  
454 until the landscape stabilises or sediment sources are exhausted (Ballantyne & Benn  
455 1994; Curry 1999; Ballantyne 2002b). A steady decline in the frequency of debris-  
456 flood deposits over the first few millennia following deglaciation (Fig. 8) might  
457 therefore be accounted for simply in terms of paraglaciation. Furthermore, the  
458 exhaustion of accessible sediment sources is a distinct possibility on extremely steep  
459 slopes, particularly at relatively high altitudes in all four catchments where extensive  
460 bedrock exposure is evidence of a more-or-less completely eroded, former till cover.  
461

462           The Jostedalbreen ice-cap, along with the Kvitekoll ice cap and the other  
463 glaciers that directly affected the catchments of the alluvial fans, are inferred to have  
464 melted away completely by ~7.3 ka (Nesje & Kvamme 1991; Nesje *et al.* 1991, 2000,  
465 2001). This date coincides with the rapid development of the Kupegelet fan which,  
466 according to our SHD dates, occurred up to two millennia later than at the other three  
467 fans. A possible explanation for later development at Kupegelet is the survival of  
468 glacier ice for longer in its narrow catchment and in the steep, north-facing cirque-like  
469 extension to the valley head on its south side. In much the same way, the north-facing  
470 valley head of Røysedalen is currently occupied by the northern outlet glacier of the  
471 Kvitekoll ice cap (Fig. 2). A second possible explanation is that rapid fan  
472 development at Kupegelet was triggered by the paraperiglacial degradation of  
473 permafrost in the upper catchment: i.e. it was a conditional response to the transition  
474 from permafrost to seasonal-freezing regime (cf. Mercier 2008; Scarpozza 2016;  
475 Matthews *et al.* 2018). The lower altitudinal limit of discontinuous mountain  
476 permafrost currently lies at ~1600 m a.s.l. in the Jostedalbreen region, and could be  
477 lower in north-facing rock walls (Etzelmüller & Hagen 2005; Gislås *et al.* 2016;  
478 Steiger *et al.* 2016).

479  
480           Landscape stabilisation and hence reduced paraglacial aggradation on the fans  
481 are likely to have been accentuated by the spread of a dense tree cover onto the lower-  
482 altitude slopes of the catchments in the Early to Middle Holocene as a result of a  
483 warmer climate than today during the Holocene Thermal Maximum (HTM). Present-  
484 day tree lines attain altitudes of 850–1000 m at favourable locations within the four  
485 catchments (<https://www.norgeskart.no/>) and, based on pollen analyses from the

486 valleys around Jostedalsbreen, would have been at least 200 m higher during the  
487 HTM (Nesje & Kvamme 1991; Nesje et al. 1991; see also Wilford et al. 2005;  
488 Marston 2010; Pawlik 2013).

489

490 The SHD evidence indicates that paraglacial sedimentation was the dominant  
491 control on the Early Holocene development of all four fans. However, there was  
492 greater divergence in their evolution during the Late-Holocene: the Kvamsdalen fan  
493 seems to have become an essentially relict landform when paraglacial effects  
494 effectively ceased at ~4.0 ka; the Kupegelet fan experienced continuing deposition  
495 from debris floods at a much reduced level until at least ~2.0 ka; the Erikstølsdalen  
496 and Snøskreda fans appear to have been dominated by a different sedimentological  
497 and hydrological regime, which began sometime after ~8.0 ka and has continued to  
498 the present day. This new regime, which is attributed to the diminished sediment  
499 supply after the cessation of the debris floods of the paraglacial phase, was dominated  
500 by snow-avalanches and fluvial activity, and has left boulder deposits unburied at the  
501 margins of these fans. Evidence of the importance of snow avalanching at these sites  
502 includes the presence of extensive accumulations of snow on the fan apex and  
503 upstream, which are clearly visible on late-summer aerial photography  
504 (<https://www.norgebilder.no/>), isolated angular boulders scattered over the fan  
505 surface, and the names ‘Snøskreda’ (which means snow avalanche in Norwegian) and  
506 ‘Erikstølskreda’, which are established place names used on topographic maps.  
507 Fluvial activity is indicated by gravel deposits alongside the current stream, largely  
508 vegetated distributary channels, and the boulder-rich sediments deposited by the AD  
509 1979 flash flood.

510

511 Neoglaciation from ~6.1 ka and Late-Holocene glacier variations appear to  
512 have made significant contributions to the later phases of fan evolution, particularly at  
513 the Erikstølsdalen and Kupegelet sites. Based on moraines dated by historical  
514 evidence and lichenometric dating, it is well established that the main glaciers in this  
515 region, including Austerdalsbreen and Nystølsbreen (Fig. 2) attained their Late-  
516 Holocene maximum extent *c.* AD 1750, in the Little Ice Age (Bickerton & Matthews  
517 1993) and, in the case of Nystølsbreen, the glacier extended onto its fan (McEwen *et*  
518 *al.* 2011). Similar undated moraines in Røysedalen indicate that the northern outlet of  
519 the Kvitekoll ice cap expanded at this time (Fig. 2), and strongly suggest that both this

520 ice cap and Tverradalsbreen overflowed into the fan catchments during the Little Ice  
521 Age. Although there is insufficient evidence from this study to detect any century- to  
522 millennia-scale responses, the existence of small glaciers in these catchments during  
523 the Little Ice Age and earlier neoglacial glacier expansion episodes (cf. Nesje et al.  
524 2008; Nesje 2009; Matthews 2013) are likely to have affected meltwater discharge,  
525 slope processes and sediment loads, and hence variations in Late-Holocene fan  
526 aggradation (cf. McEwen *et al.* 2011; Laute & Beylich 2012, 2013). Similarly,  
527 changes in fan aggradation would be expected from any Late-Holocene variations in  
528 snow meltwater discharge and snow-avalanche frequency, the latter affecting the  
529 Erikstølsdalen and Snøskreda fans in particular.

530

531 *A regional model of alluvial fan evolution in recently-deglaciated mountains*

532

533 The evolution of alluvial fans in the SE Jostedalbreen region – including the four  
534 fans reported in this study and the Nystølen fan investigated by McEwen *et al.* (2011)  
535 – can be generalized as a regional conceptual model (Fig. 12A–D) that includes local  
536 variations in the timing of four main phases of fan development, the changing nature  
537 and intensity of aggradational processes, variations in glacier size, and changes in the  
538 climatic and hence hydrological regime during the Holocene. This model extends and  
539 refines a previous model of alluvial fan development in glacierized catchments  
540 presented by McEwen *et al.* (2020) and makes a broader contribution to the rather  
541 limited understanding of alluvial fans in alpine and subalpine environments from  
542 various perspectives (cf. Kostaschuk *et al.* 1986; Eyles & Kocsis 1988; Derbyshire &  
543 Owen 1990; Blair & McPherson 1994; Cavalli & Marchi 2008; Korup & Clagues  
544 2009; Schneuwly-Bollscheiler *et al.* 2013; Heiser *et al.* 2015; Tomczyk *et al.* 2019).

545

546 *Phase 1: Intense paraglacial aggradation (9.7–8.0 ka).* – The first phase begins  
547 immediately after deglaciation. Aggradation rapidly intensifies as gully propagation  
548 takes place in steep and initially unvegetated till-mantled catchment slopes. Peak  
549 paraglacial aggradation, on the basis of the frequency of dated debris-flood deposits  
550 from three fans (Kvamsdalen, Erikstølsdalen and Snøskreda), is placed at ~9.0 ka  
551 (Fig. 12A, B).

552



553           The start of this intense phase may be delayed by the late survival of glacier  
554 ice within the catchment (or by paraperiglacial permafrost degradation), as  
555 hypothesised for Kupegelet. Intense paraglacial aggradation takes place not only at a  
556 time of shrinking glaciers (Fig. 12C), but also in a climatic environment of high and  
557 rising temperatures and increasing precipitation (Fig. 12D). The hydrological effect of  
558 this is likely to contribute to relatively high discharges from both glacial and snow  
559 meltwater.

560

561 *Phase 2: Reduced paraglacial aggradation (8.0–4.0 ka).* – The transition to a phase of  
562 reduced paraglacial aggradation is considered, on the basis of the dating evidence  
563 from Kvamsdalen and Kupegelet, to occur no more than 2000 years after the start of  
564 the intense phase. The start of this second phase is therefore placed at ~8 ka for  
565 Kvamsdalen (Fig. 12A) and this date is also used in Fig. 12B (although this phase was  
566 delayed to ~6.0 ka at Kupegelet). The apparent absence of debris-floods for many  
567 millennia from ~8.0 ka at Erikstølsdalen and Snøskreda is attributed to their burial by  
568 later fluvial and snow-avalanche sedimentation.

569

570           Reduced aggradation after ~8.0 ka is primarily a response to the reduced  
571 availability of sediment and the possible eventual exhaustion of sediment sources  
572 within the catchment. Three other factors are seen as contributing to increasing  
573 stability within the catchment and the reduction of aggradation on the fans. First, with  
574 glaciers very small or absent from the catchments (Fig. 12C), the paraglacial sediment  
575 load of the rivers is supplemented to a negligible extent by glaciofluvial sediments  
576 direct from glaciers. Second, stabilization increases over time with the establishment  
577 of vegetation and, in particular, with the spread of trees at relatively low altitudes  
578 within the catchments. Third, temperatures remain high while precipitation is much  
579 reduced, at least until ~6.0 ka (Fig. 12D): the climatic regime therefore suggests  
580 reduced runoff from snowmelt at this time. Diminution of paraglacial aggradation is  
581 shown in Fig. 12B to continue until ~4.0 ka, though this must be regarded as an  
582 arbitrary point on the long-term declining trend.

583

584 *Phase 3: Fan surface stability (4.0–0 ka).* – A phase of near-zero aggradation on the  
585 fan surface is the logical outcome of the exhaustion of sediment supply within the  
586 catchment, and is recognised at Kupegelet from ~2.0 ka and at Kvamsdalen from

587 ~4.0 ka (Fig. 12A). Fan surface stability may also follow from flows with decreasing  
588 sediment concentrations resulting from an increase in discharge during Late-Holocene  
589 climatic deterioration and the early stages of neoglacial glacier growth. Judged in  
590 terms of the non-existence of dated debris-flood deposits, stabilization of fan surfaces  
591 did not take place before ~4.0 ka, but evidence of older stable phases could be buried  
592 by later aggradation.

593

594 The possibility of entrenchment introduces a further complication (cf.  
595 McEwen *et al.* 2020), which may itself be initiated in response to reduced sediment  
596 loads during the phase of reduced paraglacial aggradation. In this study, entrenchment  
597 is exhibited to some extent by the modern streams on the upper (proximal) parts of  
598 each fan (Fig. 4). This helps explain the tendency to asymmetrical development, at  
599 least during the later stages of fan evolution, and hence the persistence and survival of  
600 debris-flood deposits on the north side of each fan, as well as towards each fan toe.  
601 Each stream currently discharges to the south side of the fan, topographically-  
602 controlled avulsions having followed the slope of the fan (cf. De Haas *et al.* 2019),  
603 which is in turn influenced by the direction of the trunk valley, thus diverting flows  
604 away from the north side of the fans.

605

606 *Phase 4: Neoglacial re-activation (4.0–0 ka).* – Re-activation takes place in the Late-  
607 Holocene in response to climatic deterioration and glacier growth, provided that  
608 sufficient sediment sources are available and accessible within the catchment. The  
609 onset of this final phase is placed at ~4.0 ka on the basis of dated debris-flood deposits  
610 at Kupegjelet and Erikstølsdalen (Fig. 12A). Small glaciers regenerating as early as  
611 ~6.1 ka (Fig. 12C), and/or the associated climatic deterioration involving decreasing  
612 temperatures and increasing precipitation (Fig. 12D), are seen as unlikely to have had  
613 a major effect on aggradation initially. By ~4.0 ka, however, as neoglaciation  
614 intensifies, increasing discharge combined with greater potential for bedload  
615 generation and transport is consistent with renewed aggradation.

616

617 Re-activation greatly increases the potential for burial of older deposits, which  
618 is inferred to account for the apparent absence of debris-flood activity after ~8.0 ka at  
619 Erikstølsdalen and Snoskreda. This argument is supported by the confinement of the  
620 debris-flood deposits at the latter fan to its extreme distal fringe, the remainder of the

621 fan surface being affected by more recent water-flood and snow-avalanche deposits.  
622 Neoglacial re-activation associated with an increase in water-flood and snow-  
623 avalanches was even more effective at the Nystølen fan (Fig. 12A), where the whole  
624 of the fan surface dates from the Little Ice Age (Lewis & Birnie 2001; McEwen *et al.*  
625 2011).

626

627

## 628 Conclusions

629

- 630 • Boulder deposits from four subalpine alluvial fans in the SE Jostedalbreen  
631 region of southern Norway were dated using SHD, demonstrating the  
632 usefulness of this technique for establishing the exposure-age of surface  
633 boulders in the context of the evolution of alluvial fans. The 47 SHD ages  
634 were established with 95% confidence intervals of ~700–900 years and were  
635 sufficient in number to determine a chronology of aggradational events during  
636 the Holocene based on age-frequency distributions and probability density  
637 functions.
- 638
- 639 • SHD ages indicated that a major phase of alluvial fan aggradation commenced  
640 immediately following regional deglaciation at ~9.7 ka and peaked at ~9.0–8.0  
641 ka. This is attributed to paraglacial processes within unvegetated and only  
642 partially forested catchments. On three of the fans, later aggradation failed to  
643 bury the Early-Holocene deposits, which is consistent with a regional decline  
644 in the effectiveness of paraglacial processes through the Middle Holocene. The  
645 increase in glacierization of the catchments from ~6.0 ka (neoglaciation) and  
646 especially after ~4.0 ka, which accompanied climatic deterioration and  
647 culminated in the Little Ice Age of the last few centuries, accounts for the  
648 limited number of boulder deposits and reduced aggradation over the Late  
649 Holocene. Topography of the catchments, combined with differences in the  
650 timing and extent of glaciers in the catchments during deglaciation and later  
651 neoglacial glacierization, explains the local differences in fan evolution.

652

- 653 • Alluvial fan aggradation and boulder concentrations on fan surfaces are  
654 commonly attributed to fluvial activity (water floods) and/or debris flows. This  
655 study highlights the potential importance of debris floods, of which relatively  
656 little is known, especially in the context of alluvial fan evolution. The  
657 morphology of the boulder deposits on our fans is distinctive, consisting of  
658 broad, low ridges with distal splays but no evidence of the levées characteristic  
659 of debris flows. The degree of boulder rounding and crude sorting present in  
660 the boulder deposits, and the catchment characteristics also point to an  
661 intermediate flow-type between water flow and debris flow. Such flows  
662 require a debris concentration of 40-70% by weight, which we argue was  
663 attained during the paraglacial reworking of till deposits in these steep  
664 catchments.
- 665
- 666 • Our results have led to the development of a conceptual model of alluvial fan  
667 evolution for glacierized catchments and recently deglaciated mountains SE of  
668 the Jostedalsbreen ice cap (Fig. 12). A phase of ‘intense paraglacial  
669 aggradation’ is succeeded by phases of ‘reduced paraglacial aggradation’, ‘fan  
670 surface stability’ and ‘neoglacial re-activation’. The model incorporates the  
671 timing of deglaciation, subsequent glacier activity, catchment topography and  
672 vegetation cover, sediment sources and climatic changes linked to the  
673 hydrological regime, all of which are effective controls on fan aggradation.  
674 The model should be applicable to some degree in other recently deglaciated  
675 mountain regions with small, steep catchments, if only as a template for  
676 comparison.

677  
678

679 *Acknowledgements.* – Fieldwork was carried out on the Swansea University  
680 Jotunheimen Research Expeditions of 2013 and 2014. The authors are grateful to  
681 Amber Vater and Ross Pinnock for Fig. 3 and field assistance, Ole Jacob and Tove  
682 Grindvold (Leirvassbu) for continued logistical support, and Anna C. Ratcliffe for  
683 preparing the figures for publication. The data that support the findings of this study  
684 are available from the corresponding author upon reasonable request. This manuscript  
685 constitutes Jotunheimen Research Expeditions, Contribution No. 216 (see  
686 <http://jotunheimenresearch.wixsite.com/home>).

687  
688  
689

*Author contributions.* – Matthews, McEwen and Owen conceived and planned the study based on previous work and carried out the fieldwork; Matthews analysed the

690 Schmidt-hammer and lichenometry data, calculated the SHD ages and wrote the first  
691 draft of the paper; McEwen did the palaeohydrological calculations; Los carried out  
692 the probability density analyses. All authors contributed substantially to the final  
693 draft.

694

## 695 References

696

697 Aa, A. R. 1982: Ice movements and deglaciation in the area between Sogndal and  
698 Jostedalbreen, western Norway. *Norsk Geologisk Tidsskrift* 62, 179–190.

699

700 Aune, B. 1993: *Temperatur normaler: normalperiode 1961–90 (Rapport 02/93)*. 63 pp. Den  
701 Norske Meteorologiske Institutt, Oslo.

702

703 Ballantyne, C. K. & Benn, D. I. 1994: Paraglacial slope adjustment and resedimentation  
704 following glacier retreat, Fåbergstølsbreen, Norway. *Arctic & Alpine Research* 26, 255–269.

705

706 Ballantyne, C. K. 2002a: Paraglacial geomorphology. *Quaternary Science Reviews* 21, 1935–  
707 2017.

708

709 Ballantyne, C. K. 2002b: A general model of paraglacial landscape response. *The Holocene*  
710 12, 371–376.

711

712 Ballantyne, C. K. 2013: Paraglacial geomorphology. In Elias, S. (ed.): *Encyclopedia of*  
713 *Quaternary Science*. 553–565. Elsevier: Amsterdam.

714

715 Bickerton, R. J. & Matthews, J. A. 1992: On the accuracy of lichenometric dates: an  
716 assessment based on the ‘Little Ice Age’ moraine sequence of Nigardsbreen, southern  
717 Norway. *The Holocene* 2, 227–237.

718

719 Bickerton, R. W. & Matthews, J. A. 1993: ‘Little Ice Age’ variations of outlet glaciers from  
720 the Jostedalbreen ice cap, southern Norway: a regional lichenometric-dating study of ice-  
721 marginal moraine sequences and their climatic significance. *Journal of Quaternary Science* 8,  
722 45–66.

723

724 Blair, T. C. & McPherson, J. G. 1994: Alluvial fans and their natural distinction from rivers  
725 based on morphology, hydraulic processes, sedimentary processes and facies assemblages.  
726 *Journal of Sedimentary Research* 64A, 450–489.

727

728 Blikra, L. H. & Nemeč, W. 1998: Postglacial colluvium in western Norway: Depositional  
729 processes, facies and palaeoclimatic record. *Sedimentology* 45, 909–959.

730

731 Blikra, L. H. & Nesje, A. 1997: Holocene avalanche history in Western Norway:  
732 chronostratigraphy and palaeoclimatic implication. *Paläoklimaforschung* 19, 299–312.

733

734 Blikra, L. H. & Selvik, S. F. 1998: Climatic signals recorded in snow avalanchedominated  
735 colluvium in western Norway: Depositional facies successions and pollen records. *The*  
736 *Holocene* 8, 631–658.

737

738 Bøe, A.-G., Dahl, S. O., Lie, Ø. & Nesje, A. 2006: Holocene river floods in the upper  
739 Glomma catchment, southern Norway: a high-resolution multiproxy record from lacustrine  
740 sediments. *The Holocene* 16, 445–455.

741

742 Boulton, G. S. 1978: Boulder shape and grain-size distributions of debris as indicators of  
743 transport paths through a glacier and till genesis. *Sedimentology* 25, 773–799.

744  
745 Bowman, D. 2019: *Principles of Alluvial Fan Morphology*. 151 pp. Springer Nature,  
746 Dordrecht.  
747  
748 Calhoun, N. C. & Clague, J. J. 2018: Distinguishing between debris flows and  
749 hyperconcentrated flows: an example from the eastern Swiss Alps. *Earth Surface Processes*  
750 *and Landforms* 43, 1280-1294.  
751  
752 Cavalli, M. & Marchi, L. 2008: Characterisation of the surface morphology of an alpine  
753 alluvial fan using airborne LIDAR. *Natural Hazards and Earth System Science* 8, 323-333.  
754  
755 Church, M. & Ryder, J. M. 1972: Paraglacial sedimentation: a consideration of fluvial  
756 processes conditioned by glaciation. *Geological Society of America Bulletin* 83, 3059-3071.  
757  
758 Costa, J. E. 1984: Physical geomorphology of debris flows. In Costa, J. E. & Fleisher, P. J.  
759 (eds): *Developments and Applications of Geomorphology*, 268-317. Springer-Verlag, Berlin.  
760  
761 Crosta, G. B. & Frattini, P. 2004: Controls on modern alluvial fan processes in the Central  
762 Alps, northern Italy. *Earth Surface Processes and Landforms* 29, 267-293.  
763  
764 Curry, A. M. 1999: Paraglacial modification of slope form. *Earth Surface Processes &*  
765 *Landforms* 24, 1213-1228  
766  
767 Curry, A. M. 2000: Observations on the distribution of paraglacial reworking of glacial  
768 drift in western Norway. *Norsk Geografisk Tidsskrift* 54, 139-147.  
769  
770 Dahl, S. O., Nesje, A., Lie, Ø., Fjordheim, K. & Matthews, J. A. 2002: Timing, equilibrium-  
771 line altitudes and climatic implications of two early-Holocene glacier advances during the  
772 Erdalen Event at Jostedalbreen, western Norway. *The Holocene* 12, 17-25.  
773  
774 D'Agustino, V. 2013: Assessment of past torrential events through historical sources. In  
775 Schneuwly-Bollschweiler, M., Stoffel, M. & Rudolf-Miklau, F. (eds.): *Dating Torrential*  
776 *Processes on Fans and Cones*, 131-146. Springer, Dordrecht.  
777  
778 De Haas, T., Kleinhaus, M. G., Carbonneau, P. E., Rubensdotter, L. & Hauber, E. 2015:  
779 Morphology of fans in the high-arctic periglacial environment of Svalbard: controls and  
780 processes. *Earth-Science Reviews* 146, 163-182.  
781  
782 De Haas, T., Densmore, A. L., Stoffel, M., Suwa, H., Imaizuma, F., Ballesteros-Cánovas, J.  
783 A. & Wasklewicz, T. 2019: Avulsions and the spatio-temporal evolution of debris-flow fans.  
784 *Earth-Science Reviews* 177, 53-75.  
785  
786 Decaulne, A., Sæmundsson, Þ., Jónsson, H. P. & Sandberg, O. 2007: Changes in deposition  
787 on a colluvial fan during the upper Holocene in the Tindastóll mountain, Skagafjörður district,  
788 north Iceland: preliminary results. *Geografiska Annaler Series A (Physical Geography)* 89A,  
789 51-63.  
790  
791 Derbyshire, E. & Owen, L. A. 1990: Quaternary alluvial fans in the Karakorum Mountains. In  
792 Rachocki, A. H. & Church, M. (eds.): *Alluvial Fans: A Field Approach*, 27-53. Wiley,  
793 Chichester.  
794  
795 Drageset, T.-A., 2001: Flomberegning for Jostedøla. Norges Vassdrags- og Energidirektoret,  
796 Dokument 2001-1. 42 pp. Norwegian Water Resources and Energy Directorate, Oslo.  
797

798 Etzelmüller, B. & Hagen, J. O. 2005: Glacier–permafrost interaction in Arctic and alpine  
799 mountain environments with examples from southern Norway and Svalbard. In Harris, C. &  
800 Murton, J. B. (eds.): Cryospheric systems: glaciers and permafrost. Geological Society,  
801 London: Special Publications, 242, 11–27.  
802

803 Evans, D. J. A. & Benn, D.I. (eds.) 2004: *A Practical Guide to the Study of Glacial*  
804 *Sediments*. 266 pp. Arnold, London.  
805

806 Eyles, N. & Kocsis, S. 1988: Sedimentology and clast fabric of subaerial debris flow facies in  
807 a glacially-influenced alluvial fan. *Sedimentary Geology* 59, 15-28.  
808

809 Førland, E. J. 1993: *Nedbørnormaler, normalperiode 1961–90. (Rapport 39/93)*. 63 pp. Den  
810 Norske Meteorologiske Institutt, Oslo.  
811

812 Germain, D. & Ouellet, M.-A. 2014: Subaerial sediment-water flows on hillslopes: essential  
813 research questions and classification challenges. *Progress in Physical Geography* 37, 813–  
814 833.  
815

816 Gislås, K., Etzelmüller, B., Lussana, C., Hjort, J., Sannel, A. B. K., Isaksen, K., Westermann,  
817 S., Kuhry, P., Christiansen, H., Frampton, A. & Akerman, J. 2017: Permafrost map for  
818 Norway, Sweden and Finland. *Permafrost and Periglacial Processes* 28, 359–378.  
819

820 Gjessing, Y. T. & Wold, B. 1980: Flommen i Jostedalen 14-15 August 1979. *Været* 1, 29-34.  
821

822 Grove, J. M. 1988: *The Little Ice Age*. 498 pp. Methuen, London.  
823

824 Harvey, A. M. 2004: Alluvial fan. In Goudie, A. S. (ed.): *Encyclopedia of Geomorphology*,  
825 15-19. Routledge, London.  
826

827 Harvey, A. M. 2013: Processes of sediment supply to alluvial fans and debris cones. In  
828 Schnewly-Bollschweiler, M., Stoffel, M. & Rudolf-Miklau, F. (eds.): *Dating Torrential*  
829 *Processes on Fans and Cones*, 15–32. Springer, Dordrecht.  
830

831 Harvey, A. M., Mather, A.E. & Stokes, M. 2005: Alluvial fans: geomorphology,  
832 sedimentology, dynamics – introduction. A review of research. In Harvey, A.M., Mather,  
833 A.E. & Stokes, M. (eds.): *Alluvial Fans: Geomorphology, Sedimentology, Dynamics*,  
834 *Geological Society, London, Special Publications* 251. 1-17.  
835

836 Heiser, M., Scheidl, C., Eisl, J., Spangl, B. and Hübl, J. 2015: Process type identification in  
837 torrential catchments in the eastern Swiss Alps. *Geomorphology* 232, 239-247.  
838

839 Hilger, P. 2019: *Rock slope failures in Norway – temporal development and climatic*  
840 *conditioning*. PhD thesis. Oslo University, Oslo, 200 pp..  
841

842 Hungr, O., Evans, S. G., Bovis, M. J. & Hutchinson, J. N. 2001: A review of the classification  
843 of landslides of the flow type. *Environmental and Engineering Geoscience* 7, 221–238.  
844

845 Hungr, O., Leroueil, S. & Picarelli, L. 2014: The Varnes classification of landslide types, an  
846 update. *Landslides* 11, 167–194.  
847

848 Innes, J. L. 1985a: Lichenometric dating of debris-flow deposits on alpine colluvial fans in  
849 southwest Norway. *Earth Surface Processes and Landforms* 10, 519–524.  
850

851 Innes, J. L. 1985b: Magnitude-frequency relations of debris flows in northwest Europe.  
852 *Geografiska Annaler Series A (Physical Geography)* 67A, 23–32.

853  
854 Jomelli, V. 2013: Lichenometric dating of debris avalanche deposits with an example from  
855 the French Alps. In Schneuwly-Bollschweiler, M., Stoffel, M. & Rudolf-Miklau, F. (eds.):  
856 *Dating Torrential Processes on Fans and Cones*, 211–224. Springer, Dordrecht.  
857  
858 Korup, O. & Clague, J. J. 2009: Natural hazards, extreme events, and mountain topography.  
859 *Quaternary Science Reviews* 28, 977–990.  
860  
861 Kostaschuk, R., MacDonald, G. & Putnam, P. 1986: Depositional processes and alluvial fan-  
862 drainage basin morphometric relationships near Banff, Alberta, Canada. *Earth Surface*  
863 *Processes and Landforms* 11, 471–484.  
864  
865 Laute, K. & Beylich, A. A. 2012: Influences of the Little Ice Age glacier advance on hillslope  
866 morphometry and development in paraglacial valley systems around the Jostedalbreen ice  
867 cap in Western Norway. *Geomorphology* 167–168, 51–69.  
868  
869 Laute, K. & Beylich, A. A. 2013: Holocene hillslope development in glacially formed valley  
870 systems in Nordfjord, western Norway. *Geomorphology* 188, 12–30.:  
871  
872 Lewis, S. G. & Birnie, J. F. 2001: Little Ice Age alluvial fan development in Langedalen,  
873 western Norway. *Geografiska Annaler Series A (Physical Geography)* 83A, 179–190.  
874  
875 Lutro, O. & Tveten, E. 1996: *Geologisk kart over Norge, berggrunskart Årdal 1:250,000*.  
876 Norges Geologiske Undersøkelse, Trondheim.  
877  
878 Marston, R. A. 2010: Geomorphology and vegetation on hillslopes: interaction, dependencies  
879 and feedback loops. *Geomorphology* 116, 206–217.  
880  
881 Matthews, J. A. 1987: Regional variation in the composition of Neoglacial end moraines,  
882 Jotunheimen, Norway: an altitudinal gradient on clast roundness and its possible  
883 palaeoclimatic significance. *Boreas* 16, 173–188.  
884  
885 Matthews, J. A. 2013: Neoglaciation in Europe. In Elias, S.A. (ed.): *Encyclopedia of*  
886 *Quaternary Science*, 2, 257–268. Elsevier, Amsterdam.  
887  
888 Matthews, J. A. & McEwen L. J. 2013: High Precision Schmidt-hammer exposure-age dating  
889 (SHD) of flood berms, Vetlestølsdalen, alpine southern Norway: first application and some  
890 methodological issues. *Geografiska Annaler Series ,(Physical Geography)* 95, 185–194.  
891  
892 Matthews, J. A. & Owen, G. 2010: Schmidt-hammer exposure-age dating: developing linear  
893 age-calibration curves using Holocene bedrock surfaces from the Jotunheimen-Jostedalbreen  
894 regions of southern Norway. *Boreas* 39, 105–115.  
895  
896 Matthews, J. A. & Trenbirth, H. E. 2011: Growth rate of a very large crustose lichen  
897 (*Rhizocarpon* subgenus) and its implications for lichenometry. *Geografiska Annaler Series A*  
898 *(Physical Geography)* 93A, 27–39.  
899  
900 Matthews, J. A. & Wilson, P. 2015: Improved Schmidt-hammer exposure ages for active and  
901 relict pronival ramparts in southern Norway, and their palaeoenvironmental implications.  
902 *Geomorphology* 246, 7–21.  
903  
904 Matthews, J. A. & Winkler, S. 2011: Schmidt-hammer exposure-age dating (SHD):  
905 application to early Holocene moraines and a reappraisal of the reliability of terrestrial  
906 cosmogenic-nuclide dating (TCND) at Austanbotnbreen, Jotunheimen, Norway. *Boreas* 40,  
907 256–270.



908  
909 Matthews, J. A., Shakesby, R. A., McEwen, L. J., Berrisford, M. S., Owen, G. & Bevan, P.  
910 1999: Alpine debris flows in Leirdalen, Jotunheimen, Norway, with particular reference to  
911 distal fans, intermediate-type deposits and flow types. *Arctic, Antarctic and Alpine Research*  
912 *31*, 421–435.  
913  
914 Matthews, J. A., Dahl, S.-O., Dresser, P. Q., Berrisford, M. S., Lie, Ø., Nesje, A. & Owen, G.  
915 2009: Radiocarbon chronology of Holocene colluvial (debris-flow) activity at Sletthamn,  
916 Jotunheimen, southern Norway: a window on the changing frequency of extreme climatic  
917 events and their landscape impact. *The Holocene* *19*, 1107–1129.  
918  
919 Matthews, J. A., Nesje, A. & Linge, H. 2013: Relict talus-foot rock glaciers at Øyberget,  
920 upper Ottadalen, southern Norway: Schmidt hammer exposure ages and palaeoenvironmental  
921 implications. *Permafrost and Periglacial Processes* *24*, 336–346.  
922  
923 Matthews, J. A., Winkler, S. & Wilson, P. 2014: Age and origin of ice-cored moraines in  
924 Jotunheimen and Breheimen, southern Norway: insights from Schmidt-hammer exposure-age  
925 dating. *Geografiska Annaler Series A (Physical Geography)* *96*, 531–548.  
926  
927 Matthews, J. A., McEwen, L. J. & Owen, G. 2015: Schmidt-hammer exposure-age dating  
928 (SHD) of snow-avalanche impact ramparts in southern Norway: approaches, results and  
929 implications for landform age, dynamics and development. *Earth Surface Processes and*  
930 *Landforms* *40*, 1705–1718.  
931 .  
932 Matthews, J. A., Winkler, S., Wilson, P., Tomkins, M. D., Dortsch, J. M., Mourne, R. W.,  
933 Hill, J. L., Owen, G. & Vater, A. E. 2018: Small rock-slope failures conditioned by Holocene  
934 permafrost degradation: a new approach and conceptual model based on Schmidt-hammer  
935 exposure-age dating, Jotunheimen, southern Norway. *Boreas* *47*, 1144–1169.  
936  
937 Mauri, A., Davis, B. A. S., Collins, P. M. & Kaplan, J. O. 2015: The climate of Europe during  
938 the Holocene: a gridded pollen-based reconstruction and its multi-proxy evaluation.  
939 *Quaternary Science Reviews* *112*, 109–127.  
940  
941 Mayer, B., Stoffel, M., Bollschweiller, M., Hübl, J. & Rudolf-Miklau, F. 2010: Frequency and  
942 spread of debris floods on fans: a dendrogeomorphic case study from a dolomite catchment in  
943 the Austrian Alps. *Geomorphology* *118*, 199–206.  
944  
945 McCarroll, D. 1987: The Schmidt hammer in geomorphology: five sources of instrument  
946 error. *British Geomorphological Research Group. Technical Bulletin* *36*, 16–27.  
947  
948 McCarroll, D. 1994: The Schmidt hammer as a measure of degree of rock surface weathering  
949 and terrain age. In Beck, C. (ed.): *Dating in Exposed and Surface Contexts*, 29–45.  
950 University of New Mexico Press, Albuquerque.  
951  
952 McEwen, L. J., Owen, G., Matthews, J. A. & Hiemstra, J. F. 2011: Late Holocene  
953 development of a Norwegian alpine alluvial fan affected by proximal glacier variations,  
954 episodic distal undercutting and colluvial activity. *Geomorphology* *127*, 198–215.  
955  
956 McEwen, L. J., Matthews, J. A. & Owen, G. 2020: Development of a Holocene glacier-fed  
957 alluvial fan based on calibrated- and relative-age dating techniques: the Illåe fan,  
958 Jotunheimen, Norway. *Geomorphology* (in press).  
959  
960 Melton, M. A. 1965: The geomorphic and paleoclimatic significance of alluvial deposits in  
961 southern Arizona. *Journal of Geology* *73*, 1–38.  
962

- 963 Mercier, D. 2008: Paraglacial and paraperiglacial land systems: concepts, temporal scales and  
 964 spatial distribution. *Geomorphologie: Relief, Processus, Environnement* 14, 223–233.  
 965
- 966 Moscariello, A. 2018: Alluvial fans and fluvial fans at the margins of continental sedimentary  
 967 basins: geomorphic and sedimentological distinction for geo-energy exploration and  
 968 development. In Ventra, D. & Clarke, L. E. (eds.): *Geology and Geomorphology of Alluvial  
 969 and Fluvial Fans*. Geological Society, London, *Special Publications* 440, 215–243.  
 970
- 971 Mottershead, D. N & Collin, R. L. 1976: A study of Flandrian glacier fluctuations in  
 972 Tunsbergdalen, southern Norway. *Norsk Geologiske Tidsskrift* 56, 413–436.  
 973
- 974 Nesje, A. 2009: Latest Pleistocene and Holocene alpine glacier fluctuations in Scandinavia.  
 975 *Quaternary Science Reviews* 28, 2119–2136.  
 976
- 977 Nesje, A. & Kvamme, M. 1991: Holocene glacier and climatic variations in western Norway:  
 978 evidence for early Holocene glacier demise and multiple Neoglacial events. *Geology* 19, 610–  
 979 612.  
 980
- 981 Nesje, A., Kvamme, M., Rye, N. & Løvlie, R. 1991: Holocene glacial and climatic history of  
 982 the Jostedalbreen region, western Norway: evidence from lake sediments and terrestrial  
 983 deposits. *Quaternary Science Reviews* 10, 97–114.  
 984
- 985 Nesje, A., Dahl, S. O., Andersson, C. & Matthews, J. A. 2000: The lacustrine sedimentary  
 986 sequence in Synesgardvatnet, western Norway: a continuous high-resolution record of the  
 987 Jostedalbreen ice cap during the Holocene. *Quaternary Science Reviews* 19, 1047–1065.  
 988
- 989 Nesje, A., Matthews, J. A., Dahl, S. O., Berrisford, M. S. & Andersson, C. 2001: Holocene  
 990 glacier fluctuations of Flatebreen and winter precipitation changes in the Jostedalbreen  
 991 region, western Norway, based on glaciolacustrine records. *The Holocene* 11, 267–280.  
 992
- 993 Nesje, A., Bakke, J., Dahl, S. O., Lie, Ø. & Matthews, J. A. 2008: Norwegian mountain  
 994 glaciers in the past, present and future. *Global and Planetary Change* 60, 10–27.  
 995
- 996 Ouellet, M.-A. & Germain, D. 2014: Hyperconcentrated flows on a forested alluvial fan of  
 997 eastern Canada: geomorphic characteristics, return period, and triggering scenarios. *Earth  
 998 Surface Processes and Landforms* 39, 1876–1887.  
 999
- 1000 Olsen, T., Borella, J. & Stahl, T. 2020: Clast transport history influences Schmidt hammer  
 1001 rebound values. *Earth Surface Processes and Landforms* 45, 1392–1400.  
 1002
- 1003 Owen, G. 2014: Alluvial fan. In Matthews, J. A. (ed.): *Encyclopedia of Environmental  
 1004 Change*, 33–34. Sage, London.  
 1005
- 1006 Pawlik, L. 2013: The role of trees in the geomorphic system of forested hillslopes – a review.  
 1007 *Earth-Science Reviews* 126, 250–265.  
 1008
- 1009 Pierson, T. C. 2005: Hyperconcentrated flow – transitional process between water flow and  
 1010 debris flow. In Jakob, M. & Hungr. O. (eds.): *Debris Flows/Avalanches*, 1-12. Geological  
 1011 Society of America, Boulder CO.
- 1012 Powers, M. C. 1953: A new roundness scale for sedimentary particles. *Journal of  
 1013 Sedimentary Petrology* 23, 117–119.  
 1014
- 1015 Proceq 2004: Operating instructions. Betonprüfhammer N/NR-L/LR. 21 pp. Proceq SA,  
 1016 Schwerzenbach.

1017  
1018 R Core Team 2019: R: A language and environment for statistical computing. R Foundation  
1019 for Statistical Computing, Vienna, Austria., (<https://www.R-project.org/>).  
1020  
1021 Rachocki, A. H. & Church, M. (eds.): 1990: *Alluvial Fans: a Field Approach*. 391 pp. Wiley,  
1022 Chichester.  
1023  
1024 Ryder, J. 1971: The stratigraphy and morphology of para-glacial alluvial fans in south-central  
1025 British Columbia, Canada. *Canadian Journal of Earth Sciences* 8, 279–298.  
1026  
1027 Scarpozza, C. 2016: Evidence of paraglacial and paraperiglacial crisis in Alpine sediment  
1028 transfer since the last glaciation (Tincino, Switzerland). *Quaternaire* 27, 139–155.  
1029  
1030 Schneuwly-Bollschweiler, M. & Stoffel, M. 2013: Dendrogeomorphology – tracking past  
1031 events with tree rings. In Schneuwly-Bollschweiler, M., Stoffel, M. & Rudolf-Miklau, F.  
1032 (eds.): *Dating Torrential Processes on Fans and Cones*, 165–178. Springer, Dordrecht.  
1033  
1034 Schneuwly-Bollschweiler, M., Stoffel, M. & Rudolf-Miklau, F. (eds.) 2013: *Dating*  
1035 *Torrential Processes on Fans and Cones*. 423 pp. Springer. Dordrecht.  
1036  
1037 Schürch, P., Densmore, A. L., Ivy-Ochs, S., Rosser N. J., Kober, F., Schlunegger, F.,  
1038 McArdeil, B. & Alifimov, V. 2016: Quantitative reconstruction of Late Holocene  
1039 surface evolution of an alpine debris-flow fan. *Geomorphology* 275, 46–57.  
1040  
1041 Shakesby, R. A., Matthews, J. A. & Owen, G. 2006: The Schmidt hammer as a relative-age  
1042 dating tool and its potential for calibrated-age dating in Holocene glaciated environments.  
1043 *Quaternary Science Reviews* 25, 2846–2867.  
1044  
1045 Shakesby, R. A., Matthews, J. A., Karlén, W. & Los, S. 2011: The Schmidt hammer as a  
1046 Holocene calibrated-age dating technique: testing the form of the R-value–age relationship  
1047 and defining predicted errors. *The Holocene* 21, 615–628.  
1048  
1049 Slaymaker, O. 1988: The distinctive attributes of debris torrents. *Hydrological Sciences*  
1050 *Journal* 33, 567–573.  
1051  
1052 Sletten, K. & Blikra, L. H. 2007: Holocene colluvial (debris flow and water flow) processes in  
1053 eastern Norway: stratigraphy, chronology and palaeoenvironmental implications. *Journal of*  
1054 *Quaternary Science* 22, 619–635.  
1055  
1056 Sletten, K., Blikra, L. H., Ballantyne, C. K., Nesje, A. & Dahl, S. O. 2003: Holocene debris  
1057 flows recognized in a lacustrine sedimentary succession: Sedimentology, chronostratigraphy  
1058 and cause of triggering. *The Holocene* 13, 907–920.  
1059  
1060 Stahl, T., Winkler, S., Quigley, M., Babington, M., Duffy, B. & Duke, D. 2013: Schmidt  
1061 hammer exposure-age dating (SHD) of late Quaternary fluvial terraces in New Zealand. *Earth*  
1062 *Surface Processes and Landforms* 38, 1838–1850.  
1063  
1064 Steiger, C., Etzelmüller, B., Westermann, S. & Myhra, K. S. 2016: Modelling the permafrost  
1065 distribution in steep rockwalls in Norway. *Norwegian Journal of Geology* 96, 329–341.  
1066  
1067 Stoffel, M. 2013: Tree-ring based record of debris-flow dynamics and triggering rainstorms at  
1068 Ritigraben (Swiss Alps) since AD 1570. In Schneuwly-Bollschweiler, M., Stoffel, M. &  
1069 Rudolf-Miklau, F. (eds.): *Dating Torrential Processes on Fans and Cones*, 179–185. Springer,  
1070 Dordrecht.

- 1071  
1072 Tomczyk, A. M., Ewertowski, M. W., Stawska, M. & Rachlewicz, G. 2019: Detailed alluvial  
1073 fan geomorphology in a high-arctic periglacial environment, Svalbard: application of  
1074 unmanned aerial (UAV) surveys. *Journal of Maps* 15, 460-473.  
1075  
1076 Tomkins, M. D., Dortch, J. M. & Hughes, P. D. 2016: Schmidt hammer exposure dating  
1077 (SHED): establishment and implications for the retreat of the last British Ice Sheet.  
1078 *Quaternary Geochronology* 33, 46–60.  
1079  
1080 Tomkins, M. D., Dortch, J. M., Hughes, P. D., Huck, J. J., Stimson, A. G., Delmas, M.,  
1081 Calvet, M. & Pallas, R. 2018: Schmidt hammer exposure dating (SHED): rapid age  
1082 assessment of glacial landforms in the Pyrenees. *Quaternary Research* 90, 26–37.  
1083  
1084 Vasskog, K., Nesje, A., Støren, E. N., Waldmann, N., Chapron, E. & Ariztegui, D. 2011: A  
1085 Holocene record of snow-avalanche and flood activity reconstructed from a  
1086 lacustrine sedimentary sequence at Oldevatnet, western Norway. *The Holocene* 21, 597–614.  
1087  
1088 Ventra, D. & Clarke, L. E. 2018: Geology and geomorphology of alluvial and fluvial fans:  
1089 current progress and research perspectives. In Ventra, D. & Clarke, L. E. (eds.): *Geology and*  
1090 *geomorphology of alluvial and fluvial fans*. Geological Society, London, Special Publications  
1091 440, 1–21.  
1092  
1093 Viles, H., Goudie, A., Grabb, S. & Lalley, J. 2011: The use of the Schmidt hammer and  
1094 Equotip for rock hardness assessment in geomorphology and heritage science: a comparative  
1095 analysis. *Earth Surface Processes and Landforms* 36, 320–333.  
1096  
1097 Wells S. G. & Harvey, A. M. 1987: Sedimentologic and geomorphic variations in storm  
1098 generated alluvial fans, Howgill Fells, northwest England. *Geological Society of America*  
1099 *Bulletin* 98, 182–198.  
1100  
1101 Wilford, D. J., Sakals, M. E., Innes, J. L., Sidle, R. C. & Bergerud, W. A. 2004: Recognition  
1102 of debris flow, debris flood and flood hazard through watershed morphometrics. *Landslides* 1,  
1103 61–66.  
1104  
1105 Wilford, D. J., Sakals, M. E., Innes, J. L. & Sidle, R. C. 2005: Fans with forests:  
1106 contemporary hydrogeomorphic processes on fans with forests in west central British  
1107 Columbia, Canada. In Harvey, A. M., Mather, A. E. & Stokes, M. (eds.): *Alluvial Fans:*  
1108 *Geomorphology, Sedimentology, Dynamics* Geological Society, London, Special  
1109 *Publications* 251, 25–40.  
1110  
1111 Williams, G. 1983: Paleohydrological methods and some examples from Swedish fluvial  
1112 environments: I. Cobble and boulder deposits. *Geografiska Annaler Series A (Physical*  
1113 *Geography)* 65A, 224–243.  
1114  
1115 Wilson, P. & Matthews, J. A. 2016: Age assessment and implications of late Quaternary  
1116 periglacial and paraglacial landforms on Muckish Mountain, northwest Ireland, based on  
1117 Schmidt-hammer exposure age dating (SHD). *Geomorphology* 270, 134–144.  
1118  
1119 Wilson, P., Matthews, J. A. & Mourne, R. W. 2017: Relict blockstreams at Insteheia,  
1120 Valldalen-Tafjorden, southern Norway: their nature and Schmidt-hammer exposure age.  
1121 *Permafrost and Periglacial Processes* 28, 286–297.  
1122  
1123 Wilson, P., Linge, H., Matthews, J. A., Mourne, R. W. & Olsen, J. 2019: Comparative  
1124 numerical surface-age dating (<sup>10</sup>Be and Schmidt hammer) of an early-Holocene rock

1125 avalanche at Alnesfjellet, Valldalen, southern Norway. *Geografiska Annaler Series A*  
 1126 (*Physical Geography*) 101, 293–309.  
 1127  
 1128 Winkler, S. 2009: First attempt to combine terrestrial cosmogenic nuclide ( $^{10}\text{Be}$ ) and Schmidt  
 1129 hammer relative-age dating: Strauchon Glacier, Southern Alps, New Zealand. *Central*  
 1130 *European Journal of Geosciences* 1, 274–290.  
 1131  
 1132 Winkler, S., Matthews, J. A., Mourné, R.W. & Wilson, P. 2016: Schmidt hammer  
 1133 exposure ages from periglacial patterned ground (sorted circles) in Jotunheimen, Norway, and  
 1134 their interpretive problems. *Geografiska Annaler Series A (Physical Geography)* 98A, 265-  
 1135 285.  
 1136  
 1137 Zieliński, T. & van Loon, A. J. 2000: Subaerial terminoglacial fans III: overview of  
 1138 sedimentary characteristics and depositional model. *Geologie en Mijnbouw/Netherlands*  
 1139 *Journal of Geosciences* 79, 93-107.

1140

1141

1142

1143

#### 1144 **FIGURE CAPTIONS**

1145

1146 Fig. 1. Location of the study area SE of the Jostedalbreen ice cap in southern  
 1147 Norway.

1148

1149 Fig. 2. The study area, the alluvial fans, their catchments, and the location of sites  
 1150 used as ‘young’ (Y) and ‘old’ (O) control surfaces for SHD dating. Note also the  
 1151 location of Fig. 4 A, B.

1152

1153 Fig. 3. Kupegjelet fan, Austerdalen, from opposite valley side. Note extensive boulder  
 1154 deposits on the fan surface, and the rugged catchment with exposed bedrock on its  
 1155 upper slopes.

1156

1157 Fig. 4. Aerial photographs of the alluvial fans flown in 2017 showing numbered  
 1158 boulder deposits and fan outlines. A. Kvamsdalen and Erikstølsdalen. B. Snøskreda  
 1159 and Kupegjelet (source: <https://www.norgebilder.no/>).

1160

1161 Fig. 5. Boulder deposits on fan surfaces. A. Typical wide boulder ridge (Kupegjellet  
 1162 17, up-slope view). B. Narrow boulder ridge (Kupegjellet 13, down-slope view). C.  
 1163 Typical boulder splay at the distal end of a ridge (Kupegjellet 16). D Boulder ‘fingers’  
 1164 extending from the distal end of a boulder splay (Kupegjellet 10). E. Steep-fronted  
 1165 (lobe-like) boulder splay (Kupegjellet 8). F. Boulder ‘finger’ extending from a steep-  
 1166 fronted boulder splay (Snøskreda 1).

1167

1168 Fig. 6. A. Frequency distributions of Schmidt hammer R-values for control points of  
 1169 ‘old’ (unshaded) and ‘young’ (shaded) control points. B. The calibration equation and  
 1170 calibration curve. (A) and (B) are linked by the dashed vertical lines representing the  
 1171 mean R-values of the control points.

1172

1173 Fig. 7. Schmidt hammer R-value distributions for 47 boulder deposits from four  
 1174 alluvial fans: Kvamsdalen (Kn), Erikstølsdalen (En), Kupegjelet (Kt) and Snøskreda  
 1175 (Sa). Vertical lines indicate the mean R-values from the ‘old’ and ‘young’ control

1176 points. Sample size ( $n$ ) was 100 boulders for each boulder deposit, except for Kn 1-4  
1177 where  $n = 150$ .

1178

1179 Fig. 8. SHD ages and probability density distributions for boulder deposits from each  
1180 fan: Kvamsdalen (A), Erikstølsdalen (B), Kupegjelet(C) and Snøskreda (D). SHD age  
1181 for each boulder deposit is represented by the mean boulder exposure age (circled)  
1182 and  $2\sigma$  confidence interval (horizontal line). The probability density function for each  
1183 boulder deposit is shown as a normal distribution; combined probability density  
1184 distributions are also shown for each fan (thick black lines). In (E), the frequency of  
1185 SHD ages in 500-year intervals for the combined data set is shown together with the  
1186 regional probability density distribution (thick black line). Regional deglaciation  
1187 followed the Erdalen Event (10.2-9.7 ka), which is shown by the shaded vertical band  
1188 across all parts of the figure.

1189

1190 Fig. 9. Lichen size (mean of the five largest lichens) and lichenometric age for 47  
1191 boulder deposits from the four fans. Lichenometric age uses the 5.1 calibration  
1192 equation of Bickerton & Matthews (1991, 1992).

1193

1194 Fig. 10. Mean roundness (mean of 25 boulders) for 37 boulder deposits from the four  
1195 fans. Mean roundness values for sub-angular (SA, 3.0) and sub-rounded (SR, 4.0)  
1196 clasts are indicated.

1197

1198 Fig. 11. The largest boulders for 47 boulder deposits from the four fans. A. Maximum  
1199 boulder size B. Median size ( $D_{50}$ ) of the 10 largest boulders.

1200

1201 Fig. 12. Regional conceptual model of alluvial fan evolution in recently-deglaciated  
1202 mountains related to Holocene glacier and climatic variations. A. Phases of fan  
1203 development since deglaciation in the Jostedalsbreen region. B. Schematic intensity of  
1204 paraglacial and neoglacial drivers of aggradation. C. Generalized size of the  
1205 Jostedalsbreen ice cap (based on Nesje 2001) and D. Smoothed mean annual air  
1206 temperature (MAAT) and annual precipitation (AP) anomalies for the normal period  
1207 AD 1961-1990 in western Norway (based on Mauri et al. 2015; Hilger 2019).

1208

1209