

1 **Microbial mechanisms of carbon and nitrogen acquisition in**
2 **contrasting urban soils**

3
4 Henry W.G. Birt¹, Samuel A.F Bonnett^{1*}

5
6 *Corresponding author and current address:

7 ¹Department of Applied Sciences

8 Faculty of Health and Life Sciences

9 University of the West of England

10 Frenchay Campus

11 Coldharbour Lane

12 Bristol BS16 1QY

13 Tel: +44 (0)117 32 87157

14 Email: sam.bonnett@uwe.ac.uk

15

16

17 **Key Words:** Urban soil; nitrogen; organic matter; extracellular enzymes; substrate-
18 induced respiration.

19

20

21

22

23

24

25

26 **Abstract**

27 Urban soils play an essential role in delivering ecosystem services due to soil
28 microbial functions but there is limited evidence of the role of urban soils in the
29 global carbon cycle. Inorganic nitrogen (N) reduces microbial respiration of soil
30 organic matter (SOM) in pristine and managed forest soils but there is less evidence
31 available on the extent to which this occurs in contrasting urban soils. This study
32 examined the ephemeral effect of inorganic N and SOM (woodland versus grassland
33 urban soil) on microbial functions represented by extracellular enzyme activities and
34 microbial respiration of added substrates of contrasting quality. It was hypothesized
35 that inorganic N stimulates extracellular enzyme activities and microbial respiratory
36 responses to the addition of substrates varying in SOM quantity or quality. Results
37 showed significantly higher SOM content, DOC and dissolved phenolic compounds
38 in the woodland compared to grassland soil. In the woodland soil only, N addition
39 increased β -glucosidase and *N*-acetyl-glucosaminidase enzyme activities and
40 decreased microbial respiration responses to substrates. This suggests a microbial
41 requirement for C acquisition dependent on N availability that reduced overflow
42 respiration of the microbial community due to the composition of the woodland SOM
43 pool. In conclusion, urban soils that contrast in vegetation types and hence OM
44 content will likely differ mechanistically in response to increased N deposition and
45 climate change altering their potential ability to store soil C in the future.

46

47

48

49

50

51 **1. Introduction**

52 In urban ecosystems, soils play an essential role in delivering ecosystem
53 services via the soil microbial community. However, the microbial community is not
54 generally taken into consideration in the sustainable development of urban
55 ecosystems [1]. Urban land-use change has been identified as one of the major
56 components of environmental change because of its effects on climate, water,
57 biodiversity, carbon (C), and nutrients across large areas of the globe [2]. Despite the
58 growing body of literature [3] many aspects of urban ecosystem services have not
59 been studied conclusively and empirical evidence is still scarce [4].

60 There is considerable interest in understanding the biological processes that
61 determine C storage in soils in order to better understand mechanisms to limit
62 anthropogenic climate change [5]. Soil organic matter (SOM) provides C to a range of
63 soil organisms, being utilised as the basis for a range of organic molecules and is
64 essential for providing the energy at the base of food webs [6,7]. Decomposition of
65 SOM and ultimately carbon dioxide (CO₂) release depend on the combined response
66 of extracellular and intracellular (microbial), enzymatically mediated reactions [8].
67 Extracellular enzymes catalyze the initial hydrolysis of a variety of complex
68 polysaccharides in soil to simple monomers that can be transported actively and
69 passively into microbial cells and catabolized by intracellular enzymes producing CO₂
70 [8]. The quantity and quality of SOM is known to affect soil enzyme activities,
71 microbial respiration, and microbial biomass, and these in turn will impact on soil C
72 storage via greenhouse gas production [5,9,10]. Edmondson et al [11] have shown
73 that organic C storage may be significantly greater in urban soil than in regional
74 agricultural land at equivalent depths. However, the microbial mechanisms explaining
75 C storage in contrasting urban soils are not fully understood.

76 The nitrogen (N) cycle has been perturbed since post-industrial times through
77 enhanced reactive N in the form of anthropogenic sources such as fossil fuel burning
78 and agricultural fertilizers with implications for microbial functions and C dynamics
79 within urban soils [12,13,14]. It has been argued that N deposition is a fundamental
80 driver of increased C sequestration in forest ecosystems, significantly affecting the C
81 balance of temperate and boreal forests [15]. Janssens et al. [16] proposed that the
82 mechanism for this increased C sequestration has occurred from N stimulating woody
83 biomass at the expense of below ground C allocation. As a result, two simultaneous
84 effects occur: (1) an increase of recalcitrant sources of C into forest soil ecosystems as
85 the additional woody biomass enters soil C pools as leaf litter and (2) decreasing
86 labile C through root exudation. This results in a decrease of microbial biomass and
87 respiration that increase soil C storage.

88 Kleber [17] and Schmidt et al [18] recently discussed in detail how the
89 molecular structure (recalcitrance) of SOM alone does not control SOM stability.
90 Labile organic carbon has been defined [19] as being both chemically degradable and
91 physically accessible by soil microbes whilst availability of SOC has been defined
92 [20] as the biochemical recalcitrance of organic compounds, that is, their
93 susceptibility to enzymic degradation with further uptake of reactive products by soil
94 microorganisms. Soil microbial communities require C and nutrients to synthesize
95 extracellular enzymes to breakdown recalcitrant SOM [21]. Therefore, nutrients such
96 as N can become a limiting factor for growth of microorganisms reliant on recalcitrant
97 sources of SOM [22]. However, the production of extracellular enzymes by microbes
98 represents an energetic cost [14,23] and enzymes will only be synthesized when
99 available nutrients in simple forms are scarce resulting in the utilisation of more
100 complex and stable forms [14]. In C rich soils, N availability can limit extracellular

101 enzyme synthesis that is required to breakdown complex C polymers to simple forms
102 that subsequently enhance microbial activity and growth [24]. However, N addition to
103 soil has been found to decrease microbial decomposition and respiration [25,26]
104 especially for the SOM pools that cycle slowly [27,28]. Spohn [29] suggested three
105 mechanisms to explain these changes: (i) microorganisms ‘mine’ litter for N, burning
106 readily available (or labile) C in order to gain energy to acquire N from more
107 recalcitrant forms of SOM containing a higher C/N ratio [29,30,31] – microbial C
108 limitation will be regulated by the return on investment in extracellular enzymes that
109 depends on the availability of N [14]; (ii) microorganisms uncouple respiration from
110 energy production and only respire easily available C to dispose of it via ‘overflow
111 respiration’ to maintain the stoichiometric ratio of C/N [32] – when growing on N-
112 poor substrate, microorganisms do not have enough N to build up as much biomass as
113 the C concentration would allow due to stoichiometry [29]; and (iii) the activity of
114 oxidative enzymes involved in the degradation of aromatic compounds decreases with
115 increasing N concentration [33] suggesting that lignin degradation is a mechanism of
116 N acquisition by mining. As urban environments are characterized by high levels of N
117 deposition [13], this raises the question of whether direct N deposition to soil (i.e.
118 mineral versus organic N), leading to decreased litter C/N ratios, might control C
119 storage by driving microbial mechanisms of C and N acquisition in specific urban
120 soils.

121 This study examined the ephemeral effect of inorganic N addition on
122 extracellular enzyme activities and microbial respiration by adding substrates of
123 varying quality in two urban soils (woodland and grassland) located in close
124 proximity and characterized by contrasting SOM pools (Table 1). Woodland soils
125 typically have sparse understorey vegetation and C accumulates from dead leaf litter

126 and detritus that are decomposed to produce humified recalcitrant SOM [34]. By
127 contrast, grassland plant communities have been shown to positively affect the supply
128 of root exudates suggesting rhizospheric microbes dominate with limited requirement
129 for the production of C degrading enzymes [35]. Thus, woodland soils generally
130 consist of higher amounts of SOM with recalcitrant C (i.e. aromatic phenolics) whilst
131 grassland has a greater availability of faster cycling labile C reflecting differences in
132 SOM quantity and quality. We compared the activities of the extracellular enzymes β -
133 glucosidase, *N*-acetyl-glucosaminidase and phenol oxidase, as these enzymes are
134 involved in the decomposition of cellulose (a major type of complex C compounds in
135 soil), chitin (a significant fraction of humus-bound N in soil) and polyphenolic
136 substances (slowly decomposing complex aromatic compounds). The substrates
137 chosen for microbial respiration using the MicroResp™ method were
138 carbohydrates/complex organic polymers (D-glucose, D-arabinose, D-galactose,
139 fructose, D-trehalose, sucrose, cellulose, lignin) or amino/carboxylic acids (L-
140 arginine, L-alanine, glycine, γ -aminobutyric acid, L-malic acid, citric acid)
141 representing a range of important exudates and labile or recalcitrant organic substrates
142 in soil to investigate the microbial respiration response and substrate preference by
143 the microbial community. It was hypothesized that inorganic N would stimulate
144 extracellular enzyme activities and microbial respiration of added labile C substrates
145 but that this response would be dependent on soil characteristics of the grass and
146 woodland soil (i.e. SOM content, phenolics).

147

148

149

150

151

152 **2. Materials and Methods**

153 *2.1. Site selection and soil sampling*

154 The study area, Stoke Park Estate, is an urban, public area, located in close
155 proximity to the M32 motorway in the north of Bristol, UK (Figure 1). Soil was
156 collected from a seasonally wet semi-natural grassland pasture (*Arrhenatherum*
157 *elatius*) measuring approximately 16 ha and ancient lowland mixed broadleaf
158 woodland (*Quercus robur*, *Fraxinus excelsior*, *Fagus sylvatica*, *Aesculus*
159 *hippocastanum*) measuring approximately 8 ha (UK grid reference: 51.494827, -
160 2.553233). The grassland area was likely originally part of the woodland in the past
161 as can be seen from the regular shape of the grassland. These two habitats were
162 chosen as they are located in close proximity that minimizes variability in
163 confounding factors such as weather and physicochemistry. Despite the differences in
164 aboveground vegetation and hence SOM that was of interest as a factor, the sites were
165 also chosen as the soil types are both defined as the Denchworth vegetation type
166 (712b) (Stagni-Vertic Cambisol under FAO classification). This soil type is defined as
167 slowly permeable, seasonally waterlogged, clayey soils with similar fine loamy over
168 clayey soils [36,37,38].

169 On the 6th of November 2014, surface leaf litter, detritus and/or grass
170 leaves/roots were removed from a representative area in each habitat measuring
171 approximately 10 m². Soil samples were pooled from 4 spatial replicates to 15 cm
172 below the O horizon. These sites were chosen due to the likelihood of differences in
173 SOM quantity (i.e. loss on ignition) and quality (i.e. phenolics, SOM / inorganic N)
174 appropriate for this laboratory study rather than as a comparison of habitat types.
175 Sixteen samples of 220 grams of soil (wet weight) were homogenised through a 1 mm

176 sieve and placed in 16 x 1 litre containers (8 x woodland and 8 x grassland soils). The
177 soils were prepared within 24 h and incubated at field temperature (13 °C). Within
178 each vegetation type, at week 1 and week 3, four randomly selected replicates were
179 treated with 30 ml of deionised water and the remaining four treated with 30 ml of
180 0.125 M NH₄NO₃. This amount was chosen according to DEFRA fertilizer guideline
181 application rates [39]. The applications of N were split in two applications to prevent
182 osmotic shock and applied uniformly across the soil surface. The experimental design
183 was fully factorial with SOM (site) x NH₄NO₃ treatment allowing for interaction
184 effects.

185 Gravimetric water content was adjusted to 65% in all jars until week 4 when
186 moisture was reduced to 44% for the MicroResp™ assay [40]. Soil moisture was
187 determined every 72 h for moisture loss by weight and moisture was replaced by
188 equivalent amounts of deionised water. Jars were open to prevent CO₂ build up and
189 incubated for five weeks in total. Soil moisture content was determined by drying in
190 an oven at 105 °C for 24 h. SOM was determined by loss on ignition in a furnace at
191 500 °C for 24 h. The following soil physicochemical characteristics were determined
192 using soil extracts prepared by dissolving 2 g of wet soil in 50 ml deionised water.
193 Relative soil pH was measured using a standard electrode. A proxy for aromatic
194 dissolved organic carbon (DOC) was determined by centrifuging a sample at 12,000
195 rpm for 10 minutes and measuring UV absorbance at 254 nm [41] and also dissolved
196 phenolics by colorimetric analysis at 750 nm following reaction with Folin-Ciocalteu
197 reagent [42]. Nitrate and ammonium were determined using 1 g wet soil in 50 ml of 2
198 M KCL solution using the methods of Griffin et al 1995 [43].

199

200 *2.2. Extracellular enzyme activities*

201 Hydrolytic enzyme activities were determined by the method of DeForest
202 [44]. Fluorogenic methylumbelliferyl (MUF) substrates: MUF-glucopyranoside and
203 MUF-N-acetylglucosaminide are substrates for the determination of β -glucosidase
204 and N-acetyl-glucosaminidase enzyme activities respectively. MUF substrates (1 mM
205 – saturating concentration) and MUF standard (100 μ M) were pre-dissolved for 24 h
206 in 1ml ethylene glycol monomethyl ether. A buffer of sodium acetate trihydrate/acetic
207 acid was prepared to maintain a pH of 5.5 in the soil suspensions. 1.39 g of soil (dry
208 weight) from each experimental replicate were placed in 50 ml of buffer and 150 μ l of
209 each replicate substrate pipetted into a 300 μ l well. 100 μ l of substrate were added in
210 a specific order on a black 96 well plate with blanks, controls and quench standards.
211 The plates were incubated for one hour at 13 °C and 50 μ l of NaOH added to
212 terminate the reactions and maximise fluorescence [44]. As substrate fluorescence can
213 diverge with NaOH and time, plates were read immediately. The plates were read at
214 365 nm excitation and 450 nm emission wavelengths on a SpectraMax M2 Micro-
215 mode Microplate reader (Molecular Devices). The calculations of DeForest [44] were
216 used to calculate enzyme activity as nmol MUF g⁻¹ dry weight soil h⁻¹.

217 Phenol oxidase enzyme activities were determined by adding 0.75 ml of a 10
218 mM solution of L-3,4-dihydroxyphenylalanine (L-DOPA) to 0.75 ml of the 1.39 g
219 dry weight per 50 ml buffered soil extract in 1.5 ml centrifuge tubes [45]. Deionised
220 water was used in place of the L-DOPA solution as a blank. Final activity was
221 determined after 1 h incubation at 460 nm on a SpectraMax M2 Micro-mode
222 Microplate reader (Molecular Devices). Phenol oxidase activity (minus the blank) was
223 determined as nmol dicq g⁻¹ dry weight soil h⁻¹.

224

225 *2.3. Microbial respiration*

226 The MicroResp™ method as outlined by Campbell et al [40] was used to
227 assess respiration of added C substrates. One week prior to the assay, soils were dried
228 to 44% moisture. Deep well plates were filled with 6.9 g per 16 wells to an equal level
229 to give approximately 0.3 g of soil dry weight per well. Soil was collected from each
230 soil container using a corer as to take a representative sample over the depth of each
231 jar (10 cm). The deep well plates were incubated for 24 h to minimise disturbance
232 effects. Detection plates were prepared by pipetting 150 µl of 1% agar and cresol red
233 indicator solution at a ratio of 1:2 into clear 96 well plates. Plates were placed in a
234 dark desiccator at room temperature (20 °C) and equilibrated for 48 h. Plates were
235 read at 570 nm prior to the assay. Fifteen different substrates were pipetted in 25 µl
236 aliquots delivering 30 mg of substrate per gram soil water to the 96 deep well plates.
237 The substrates were carbohydrates/complex organic polymers (D-glucose, D-
238 arabinose, D-galactose, fructose, D-trehalose, sucrose, cellulose, lignin) or
239 amino/carboxylic acids (L-arginine, L-alanine, glycine, γ-aminobutyric acid, L-malic
240 acid, citric acid) representing a range of important labile and recalcitrant organic
241 substrates. After the addition of substrates, detection plates and deep well plates were
242 clamped together with an airtight seal and incubated at 13 °C for 6 h. Detection plates
243 were then removed and read at 570 nm for colour change on a SpectraMax M2 Micro-
244 mode Microplate reader (Molecular Devices). Measured values were then converted
245 to % CO₂ by curve fitting experimental data relating absorbance to % CO₂ where %
246 CO₂ = -0.2265 - 1.606 / (1 - 6.771 abs). The data were converted to a flux of µg CO₂
247 g⁻¹ h⁻¹ and normalized as outlined by Campbell et al. [40]. Shannon's diversity index
248 ($H' = -\sum p_i \log_2 p_i$) was computed as a measure of functional diversity [46] where p_i is
249 the ratio of respiration rate of each single C-substrate specific substrate to the sum of
250 all SIR.

251

252 2.4. Statistical analyses

253 IBM SPSS® Statistics (Version 23) and Minitab (Minitab ® 17.2.1) software
254 were used for all statistical analyses. Data were tested for normality using the
255 Anderson-Darling test and equality of variance using an *F* test. Two-way ANOVA
256 using the General Linear model with interaction terms were used to determine
257 differences between Site (woodland versus grassland) and N addition (control versus
258 NH₄NO₃) treatments in soil physicochemistry, enzymes and MicroResp™ data.
259 Community Level Physiological Profiles (CLPPs) were compared using MANOVA
260 with all substrates, carbohydrates/complex organic polymers or amino/carboxylic
261 acids. Post-hoc pairwise comparisons were used to determine specific differences
262 between treatments using Bonferroni tests.

263

264

265 3. Results

266 The woodland soil was significantly higher in SOM ($P < 0.001$ – 24 vs 17%),
267 showing higher UV absorbance at 254 nm as a proxy for aromatic DOC ($P < 0.001$ –
268 0.29 vs 0.03 abs units) and dissolved phenolic compounds ($P < 0.01$ – 9.3 vs 5.1 $\mu\text{g L}^{-1}$)
269 compared to the grassland soil (Table 1). There were no significant differences in
270 nitrate, ammonium, pH or gravimetric moisture content between the incubated grass
271 and woodland soils (Table 1). β -glucosidase activity was significantly higher in the
272 woodland soil ($P < 0.001$) and enhanced by inorganic N in the woodland soil only
273 (Bonferroni $P < 0.01$) (Fig 2). Site and N had a significant interactive effect on *N*-
274 acetyl-glucosaminidase activity ($P < 0.01$) (Fig. 2). Nitrogen addition significantly
275 reduced the SOC / inorganic N ratio ($P < 0.001$) in both woodland (2.93 to 0.30) and

276 grassland (4.80 to 0.25) but there were no significant effects of site or N on phenol
277 oxidase activity.

278 MANOVA showed distinctions in respiration when categorised into
279 carbohydrates/complex organic polymers or amino acids/carboxylic acids. The
280 respiratory response to carbohydrates showed significant main effects of Site and N
281 ($P < 0.001$ and $P < 0.01$; Fig. 3) except for an insignificant effect of N on respiration
282 of sucrose. Bonferroni post-hoc results for each carbohydrate showed distinct
283 univariate effects of the treatments on specific respiration - the interactive effect of
284 Site and N on D-arabinose ($P < 0.001$) and lignin ($P < 0.05$) (Fig. 4). MANOVA for
285 the respiratory response to amino acids/carboxylic acids showed a significant effect of
286 Site ($P < 0.01$). As with the carbohydrates, this multivariate main effect was due to
287 distinct univariate effects of Site and N on the amino acids/carboxylic acids.
288 Bonferroni post-hoc results for each carbohydrate are shown in Fig. 5. Overall, these
289 results suggest an inhibitive effect of inorganic N on respiration of substrates in the
290 woodland soil but not the grassland soil (except for fructose and γ -aminobutyric
291 acid). There were no significant differences in microbial respiration patterns using
292 MANOVA between woodland and grassland soils using the 13 substrates and no
293 significant difference between treatments in functional diversity represented by the
294 Shannon diversity index.

295

296 **4. Discussion**

297 This study suggests that rhizospheric soil systems such as grasslands may
298 show a distinct response in microbial mechanisms of C and N acquisition relative to
299 more complex SOM systems such as woodland soils that have higher detrital inputs
300 from leaf fall. Results from natural and agricultural systems have reported increased

301 enzyme activities upon the addition of N [31,47] and this effect may depend on SOM
302 quantity and/or quality (e.g. C/N ratio, phenolics). The increase in hydrolase activities
303 likely occurred as the addition of N induced the microbial production of enzymes as
304 'cost effective' in terms of return in C and nutrients [23]. This was to be expected as
305 microbes use labile sources of C and shift to complex or stable forms that require
306 catabolism by extracellular enzymes [48]. While the rate at which SOM is processed
307 is strongly controlled by the quality of the material, the extent of C limitation to the
308 microorganisms is controlled by the dynamics of exoenzymes [22], as opposed to
309 SOM quality [49] or soil microbial dynamics [50]. Allison and Vitousek [14] showed
310 that the addition of simple nutrients on their own does not necessarily affect soil
311 enzyme activities, attributed to the prerequisite of C and N for microbial enzyme
312 synthesis. The significant increase in activities in only the woodland soil suggests
313 limited accessibility to simple C (i.e. higher root exudation of labile C in grassland or
314 higher phenolic concentration in woodland), hence the microbial synthesis of
315 enzymes in response to N availability. β -glucosidase and *N*-acetyl-glucosaminidase
316 catalyse the hydrolytic release of glucose from the most abundant soil
317 polysaccharides, cellulose and chitin, so activities are closely related to soil C content
318 [51] and are induced by the availability of their substrate in excess [14]. However,
319 despite the significant difference in SOM quantity between woodland and grassland,
320 which is known to influence soil microbial functioning [7], the levels of SOM were
321 both elevated given the ecosystem types (24% and 17% respectively). Therefore, the
322 driver for enzyme production in this case was most likely SOM quality (i.e.
323 phenolics) or limited root exudation although we acknowledge that this requires
324 further investigation.

325 Phenol oxidase is a key enzyme in the degradation of lignin [52]. Lignin has
326 been shown to be highly stable and a complex substrate to degrade [53]. Therefore,
327 less complex or lower molecular weight OM was primarily decomposed by hydrolase
328 enzymes despite potentially higher phenolic content. We only examined a single time
329 event and so it is likely that a reduction in phenol oxidase activity by N addition may
330 occur in later stages as the microbial community shifts from labile to recalcitrant
331 decomposition and microbial biomass increases from total enzyme investment. Both
332 cellulolytic [54] and lignolytic microbes [55] induce exoenzyme production under
333 conditions of C limitation although we suggest that the woodland was also N limited
334 as described by the model of Schimel and Weintraub [22]. Phenol oxidase can be
335 down-regulated upon additions of low molecular weight N [33,56,57]. This is likely
336 to arise from inefficient C use and slow growth of lignolytic fungi that results from
337 competition with species that can thrive in abundant N supply [58]. These results for
338 this woodland urban soil are consistent with others [33,57] that show that inorganic N
339 accelerates the extracellular decomposition of cellulose and chitin but has no effect on
340 decomposition of lignin and polyphenolics.

341 The negative effect of inorganic N on microbial respiration and the positive
342 effect of N on extracellular enzyme activities supports model [19] and experimental
343 results [14,29,30,47,56,59]. For example, Dalmonech et al. [47] found that after 55
344 days of incubation a woodland soil treated with 75 kg N ha⁻¹ application of NH₄NO₃
345 had increased enzyme activities and microbial biomass but suppressed microbial
346 respiration. Given that we observed no change in phenol oxidase, overflow respiration
347 may be the most parsimonious explanation. Meyer et al [59] found evidence
348 suggesting that “stoichiometric decomposition” and “microbial N mining” do not
349 mutually exclude each other but operate at different temporal scales. In addition,

350 Meyer et al [59] reported that SOC-derived CO₂ was higher in unfertilized than in
351 fertilized treatments that supports Qiao et al [60] that high N supply suppresses the
352 priming effect. Both availability and lability of SOC may therefore explain the
353 contrast in enzyme activities between the grassland and woodland soils. Limited
354 access to C in the woodland compared to grassland soil may occur due to physical
355 protection of SOM, complexation with soil minerals, chemical recalcitrance (i.e.
356 phenolics) and/or low availability of low molecular weight C (i.e. rhizodeposition)
357 [17,18]. The insignificant investment in lignin degrading enzymes is supportive of
358 physicochemical mechanisms of SOM stabilisation over different rates of turnover of
359 SOM pools [61]. These results suggest ephemeral investment in hydrolase enzymes
360 due to N addition was to increase the availability of C as a microbial energy source
361 and that this C would have reduced overflow respiration.

362

363 **Conclusion**

364 These results support observations from natural and agricultural systems that
365 inorganic N addition may stimulate an increase in extracellular enzyme activities for
366 C acquisition resulting in reduced overflow respiration of the microbial community
367 due to the physicochemical nature of the SOM pool. Further long-term research is
368 required to confirm linkages between vegetation types, rhizodeposition rates, SOM
369 inputs and chemistry effects on microbial functioning in urban soils. In particular,
370 there is a need to determine how priming effects via plant exudation and shifts in the
371 microbial community structure and function may regulate the decomposition of SOM
372 in contrasting urban habitats exposed to elevated N deposition. In conclusion, urban
373 soils that contrast in vegetation types and hence OM content will likely differ

374 mechanistically in response to increased N deposition and climate change altering
375 their potential ability to store soil C in the future.

376

377

378 **Acknowledgements**

379 Support for this research was provided by the University of the West of England. We
380 thank the technical staff at UWE for their assistance and anonymous reviewers for
381 their useful comments and suggestions.

382 **References**

383 [1] J.L. Morel, C. Chenu, K. Lorenz Ecosystem services provided by soils of urban,
384 industrial, traffic, mining and military areas (SUITMAs), *J. Soils Sediments* 15 (2015)
385 1659-1666.

386 [2] N.B.F. Grimm, H. Stanley, N.E. Golubiewski, C.L. Redman, J. Wu, X. Bai, J.M.
387 Briggs, Global change and the ecology of cities, *Science* 319 (2008) 756-760.

388 [3] E. Gómez-Baggethun, Å. Gren, D.N. Barton, J. Langemeyer, T. McPhearson, P.
389 O'Farrell, E. Andersson, Z. Hamstead, P. Kremer,. Urban ecosystem services. In:
390 Elmqvist, T., et al. (Eds.), *Urbanization, Biodiversity and Ecosystem Services:*
391 *Challenges and Opportunities*. Springer, Dordrecht, Netherlands, 2013, pp. 175–180.

392 [4] D.E. Pataki, M.M. Carreiro, J. Cherrier, N.E. Grulke, V. Jennings, S. Pincetl, R.V.
393 Pouyat, T.H. Whitlow, W.C. Zipperer, Coupling biogeochemical cycles in urban
394 environments: ecosystem services, green solutions, and misconceptions, *Front. Ecol.*
395 *Environ.* 9 (2011) 27–30.

396 [5] R.D. Bardgett, C. Freeman, N. Ostle, Microbial contributions to climate change
397 through carbon cycle feedbacks, *ISME* 2 (2008) 805-814.

398 [6] U. Pascual, M. Termansen, K. Hedlund, L. Brussaard, J.H. Faber, S. Foudi, P.
399 Lemanceau, S.L. Jørgensen, On the value of soil biodiversity and ecosystem services,
400 *Ecosyst. Serv.* 15 (2015) 11-18.

401 [7] S. Zhao, K. Li, W. Zhou, S. Qiu, S. Huang, P. He, Changes in soil microbial
402 community, enzyme activities and organic matter fractions under long-term straw
403 return in north-central China, *Agri. Ecosys. Environ.* 216 (2016) 82-88.

404 [8] E. Blagodatskaya, S. Blagodatsky, N. Khomyakov, O. Mychina, Y. Kauzyakov,
405 Temperature sensitivity and enzymatic mechanisms of soil organic matter
406 decomposition along an altitudinal gradient on Mount Kilimanjaro, *Scientific Reports*
407 6 (2016) 22240.

408 [9] D.S. Powlson, P.C. Prookes, Measurement of soil microbial biomass provides an
409 early indication of changes in total soil organic matter due to straw incorporation, *Soil*
410 *Biol. Biochem.* 19 (1987) 159-164.

411 [10] F. Grosso, F. Temussi, F. De Nicola, Water-extractable organic matter and
412 enzyme activity in three forest soils of the Mediterranean area, *Euro J. Soil Biol.* 64
413 (2014) 15-22.

414 [11] J.L. Edmondson, Z.G. Davies, N. McHugh, K.J. Gaston, J.R. Leake, Organic
415 carbon hidden in urban ecosystems, *Scientific Reports* 2 (2012) 963 | DOI:
416 10.1038/srep00963.

417 [12] E.A. Davidson, The contribution of manure and fertilizer nitrogen to atmospheric
418 nitrous oxide since 1860, *Nat. Geosci.* 2 (2009) 659-662.

419 [13] P.M. Vitousek, H.A. Mooney, J. Lubchenco, J.M. Melillo, Human domination of
420 Earth's ecosystems, *Science* 277 (1997) 494-499.

421 [14] S.D. Allison, P. Vitousek, Responses of extracellular enzymes to simple and
422 complex nutrient inputs, *Soil Biol. Biochem.* 37 (2005) 937-944.

423 [15] F. Magnani, M. Mencuccini, M. Borghetti, P. Berbigier, F. Berninger, S. Delzon,
424 A. Grellle, P. Hari, P.G. Jarvis, P. Kolari, A.S. Kowalski, H. Lankreijer, B.E. Law, A.
425 Lindroth, D. Loustau, G. Manca, J.B. Moncrieff, M. Rayment, V. Tedeschi, R.
426 Valentini, J. Grace, The human footprint in the carbon cycle of temperate and boreal
427 forests, *Nature* 447 (2007) 849-851.

428 [16] I. Janssens, W. Dieleman, S. Luyssaert, J.-A. Subke, M. Reichstein, R.
429 Ceulemans, P. Ciais, A.J. Dolman, J. Grace, G. Mateucci, D. Papale, S.L. Piao, E-D.
430 Schulze, J. Tang, B.E. Law, Reduction of forest soil respiration in response to
431 nitrogen deposition, *Nat. Geosci.* 3 (2010) 315-322.

432 [17] M. Kleber, What is recalcitrant organic matter? *Environ Chem* 7 (2010) 1363-
433 1371.

434 [18] M.W. Schmidt et al, Persistence of soil organic matter as an ecosystem property.
435 *Nature* 478 (2011) 49-56.

436 [19] X.M. Zou, H.H. Ruan, Y. Fu, X.D. Yang, L.Q. Sha, Estimating soil labile
437 organic carbon and potential turnover rates using a sequential fumigation-incubation
438 procedure. *Soil Biol. Biochem.* 37 (2005) 1923-1928.

439 [20] R. Chen, M. Senbayram, S. Blagodatsky, O. Myachina, K. Dittert, X. Lin, E.
440 Blagodatskaya, Y. Kuzyakov, Soil C and N availability determine the priming effect:
441 Microbial N mining and stoichiometric decomposition theories. *Glob. Chan. Biol.* 20
442 (2014) 2356-2367.

443 [21] Z. Veres, Z. Kotroczó, I. Fekete, J. Tóth, K. Lajtha, K. Townsend, K., B.
444 Tóthmérész, Soil extracellular enzyme activities are sensitive indicators of detrital
445 inputs and carbon availability, *Appl. Soil Ecol.* 19 (2015) 18-23.

446 [22] J.P. Schimel, M.N. Weintraub, The implications of exoenzyme activity on
447 microbial carbon and nitrogen limitation in soil: a theoretical model, *Soil Biol.*
448 *Biochem.* 35 (2003) 549-563.

449 [23] A.L. Koch, The macroeconomics of bacterial growth, in: M. Fletcher, G.D.
450 Floodgate (Eds.), *Bacteria in their Natural Environments*, Academic Press, London,
451 1985, pp. 1-42.

452 [24] Y. Kuzyakov, J.K. Friedel, K. Stahr, Review of mechanisms and quantification
453 of priming effects, *Soil Biol. Biochem.* 32 (2000) 1485-1498.

454 [25] K.S. Ramirez, J.M. Craine, N. Fierer, Consistent effects of nitrogen amendments
455 on soil microbial communities and processes across biomes, *Glob. Chan. Biol.* 18
456 (2012) 1918-1927.

457 [26] L. Zhou, X. Xhou, B. Zhang, M. Lu, Y. Luo, L. Liu, B. Li, Different response of
458 soil respiration and its components to nitrogen addition among biomes: a meta-
459 analysis, *Glob. Chan. Biol.* 20 (2014) 2332-2343.

460 [27] S.E. Hobbie, W.C. Eddy, C.R. Buyarski, E.C. Adair, M.L. Ogdahl, P.
461 Weisenhorn, Response of decomposing litter and its microbial community to multiple
462 forms of nitrogen enrichment, *Ecol. Monogr.* 82 (2012) 389-405.

463 [28] C.E. Riggs, S.E. Hobbie, E.M. Bach, K.S. Hofmockel, C.E. Kazanski, Nitrogen
464 addition changes grassland soil organic matter decomposition, *Biogeochemistry* 125
465 (2015) 203-219.

466 [29] M. Spohn, Microbial respiration per unit microbial biomass depends on litter
467 layer carbon-to-nitrogen ratio, *Biogeosciences* 12 (2015) 817-823.

468 [30] J.M. Craine, C. Morrow, N. Fierer, Microbial nitrogen limitation increases
469 decomposition, *Ecology*, 88 (2007) 2105-2113.

470 [31] D.L. Moorhead, R.L. Sinsabaugh, A theoretical model of litter decay and
471 microbial interaction, *Ecol. Monogr.* 76 (2006) 151-174.

472 [32] S. Mazoni, J.A. Trofymow, R.B. Jackson, A. Porporato, Stoichiometric controls
473 on carbon, nitrogen and phosphorus dynamics in decomposing litter, *Ecol. Monogr.*
474 80 (2010) 89-106.

475 [33] K.R. Saiya-Cork, R.L. Sinsabaugh, D.R. Zak, The effects of long term nitrogen
476 deposition on extracellular enzyme activity in an *Acer saccharum* forest soil, *Soil*
477 *Biol. Biochem.* 34 (2002) 1309-1315.

478 [34] T. Boutton, J. Liao, T.R. Filley, S.R. Archer, Belowground carbon storage and
479 dynamics accompanying woody plant encroachment in a subtropical savanna, in: R.
480 Lal, R. Follett (Eds.), *Soil Carbon Sequestration and the Greenhouse Effect*, 2nd ed.,
481 Soil Science Society of America, Madison, WI, 2009, pp. 181-205.

482 [35] R. Bardgett, J. Mawdsley, S. Edwards, P. Hobbs, J. Rodwell, W. Davies, Plant
483 species and nitrogen effects on soil biological properties of temperate upland
484 grasslands, *Funct. Ecol.* 13 (1999) 650-660.

485 [36] National Soil Resources Institute (2016) Academic Soils Site Report for location
486 361776E, 177669N, 1km x 1km, National Soil Resources Institute, Cranfield
487 University. Accessed via <https://www.landis.org.uk/sitereporter/>.

488 [37] B.W. Avery, *Soil classification for England and Wales*. Soil Survey Technical
489 Monograph No.14, Harpenden, UK 1980.

490 [38] B. Clayden, J.M. Hollis, *Criteria for Differentiating Soil Series*. Soil Survey
491 Technical Monograph No.17, pp159. Harpenden, UK 1984.

492 [39] DEFRA, *Fertiliser Manual*, eighth ed., TSO, 2010.

493 [40] C.D. Campbell, S.J. Champman, C.M. Cameron, M.S. Davidson, J.M. Potts, A
494 Rapid Microtiter Plate Method To Measure Carbon Dioxide Evolved from Carbon
495 Substrate Amendments so as To Determine the Physiological Profiles of Soil
496 Microbial Communities by Using Whole Soil, *Appl. Environ. Microbiol.* 69 (2003)
497 3593-3599.

498 [41] J.L. Weishaar, G.R. Aiken, B.A. Bergamaschi, M.S. Fram, R. Fujii, K. Mopper,
499 Evaluation of specific ultraviolet absorbance as an indicator of the chemical

500 composition and reactivity of dissolved organic carbon, *Environ. Sci. Technol.* 37
501 (2003) 4702-4708.

502 [42] J.D. Box, Investigation of the Folin-Cioalteau phenol reagent for the
503 determination of polyphenolic substances in natural waters, *Wat. Res.* 17 (1983) 249-
504 261.

505 [43] G. Griffin, W. Jokela, D. Ross, Recommended Soil Nitrate-N Tests. University
506 of Delaware Agricultural Experiment Station Bulletin, 493, (1995) 22-29.

507 [44] J. DeForest, The influence of time, storage temperature, and substrate age on
508 potential soil enzyme activity in acidic forest soils using MUB-linked substrates and
509 l-DOPA, *Soil Biol. Biochem.* 41 (2009) 1180-1186.

510 [45] A. Pind, C. Freeman, M.A. Lock, Enzymic degradation of phenolic materials in
511 peatlands – measurement of phenol oxidase activity, *Plant Soil* 159 (1994) 227-231.

512 [46] G.D. Bending, M.K. Turner, F. Rayns, M. Marx, M. Wood, Microbial and
513 biochemical soil quality indicators and their potential for differentiating areas under
514 contrasting agricultural management regimes, *Soil Biol. Biochem.* 36 (2004) 1785-
515 1792.

516 [47] D. Dalmonech, A. Lagomarsino, M.C. Moscatelli, T. Chiti, R. Valentini,
517 Microbial performance under increasing nitrogen availability in a Mediterranean
518 forest soil, *Soil Biol. Biochem.* 42 (2010) 1596-1606.

519 [48] H. Glanville, J. Rousk, P. Golyshin, D.L. Jones, Mineralization of low molecular
520 weight carbon substrates in soil solution under laboratory and field conditions, *Soil*
521 *Biol. Biochem.* 48 (2012) 88–95.

522 [49] J. Dalenberg, G. Jager, Priming effect of some organic additions to ¹⁴C-labelled
523 soil, *Soil Biol. Biochem.* 21 (1989) 443-448.

524 [50] S. Weintraub, W. Wieder, C. Cleveland, A. Townsend, Organic matter inputs
525 shift soil enzyme activity and allocation patterns in a wet tropical forest,
526 *Biogeochemistry*, 114(1) (2013) 313-326.

527 [51] W.A. Dick, B. Thavamani, S. Conley, R. Blaisdell, A. Sengupta, Prediction of β -
528 glucosidase and β -glucosaminidase activities, soil organic C, and amino sugar N in a
529 diverse population of soils using near infrared reflectance spectroscopy. *Soil Biol.*
530 *Biochem.* 56 (2013) 99-104.

531 [52] R.L. Sinsabaugh, Phenol oxidase, peroxidase and organic matter dynamics of
532 soil. *Soil Biol. Biochem.* 42 (2010) 391-404.

533 [53] A. Hoffman, A. Heim, B.T. Christensen, A. Miltner, M. Gehre, M.W.I. Schmidt,
534 Lignin dynamics in two ¹³C-labelled arable soils during 18 years, *Euro. J. Soil Sci.*
535 60 (2009) 250-257.

536 [54] M.D. Romero, J. Aguado, L. González, M. Ladero, Cellulase production by
537 *Neurospora crassa* on wheat straw, *Enzy. Microbial Tech.* 25 (1999) 244-250.

538 [55] T.W. Jeffries, S. Choi, T.K. Kirk, Nutritional regulation of lignin degradation by
539 *Phanerochaete chrysosporium*. *Appl. Env. Microbiol.* 24 (1981) 290-296.

540 [56] S. Frey, M. Knorr, J. Parrent, R. Simpson, Chronic nitrogen enrichment affects
541 the structure and function of the soil microbial community in temperate hardwood and
542 pine forests, *For. Ecol. Manag.* 196 (2004) 159-171.

543 [57] B.L. Keeler, S.E. Hobbie, L.E. Kellogg, Effects of long-term nitrogen addition
544 on microbial enzyme activity in eight forested and grassland sites: Implications for
545 litter and soil organic matter decomposition, *Ecosystems*, 12 (2009) 1-15.

546 [58] J.L. DeForest, D.R. Zak, K.S. Pregitzer, A.J. Burton, Atmospheric nitrate
547 deposition and the microbial degradation of cellobiose and vanillin in a northern
548 hardwood forest, *Soil Biol. Biochem.* 36 (2004) 965-971.

549 [59] N. Meyer, G. Welp, L. Bornemann, W. Amelung, Microbial nitrogen mining
550 affects spatio-temporal patterns of substrate-induced respiration during seven years of
551 bare fallow, *Soil Biol. Biochem.* 104 (2017) 175-184.

552 [60] N. Qiao, X. Xu, Y. Hu, E. Blagodatskya, Y. Liu, D. Schaefer, Y. Kuzyakov,
553 Carbon and nitrogen additions induce distinct priming effects along an organic-matter
554 decay continuum, *Scientific Reports* 6 (2016) 1-8.

555 [61] J.A.J. Dungait, D.W. Hopkins, A.S. Gregory, A.P. Whitmore, Soil organic matter
556 turnover is governed by accessibility not recalcitrance, *Glob. Chang. Biol.* 18 (2012)
557 1781-1796.